

All random variables considered here are given on a probability space  $(\Omega, \mathcal{F}, P)$ .

1. (25 pts) Let  $X_1, X_2, \dots$  be a sequence of independent, identically distributed (i.i.d.) random variables taking only positive values. Let  $S_n = \sum_{j=1}^n X_j$  and  $N_t = \sup\{n : S_n \leq t\}$  for  $t > 0$ .

(a) Suppose  $EX_1 = \mu \in (0, \infty]$ . Prove that  $\lim_{t \rightarrow \infty} \frac{N_t}{t} = \frac{1}{\mu}$  almost surely (a.s.), where  $1/\infty = 0$ .

(b) Suppose  $\mu < \infty$  and  $\sigma^2 = \text{Var}(X_1) < \infty$ . Using the fact that the convergence in central limit theorem is uniform, show that  $\frac{N_t - t\mu^{-1}}{\sqrt{t\sigma^2\mu^{-3}}}$  converges in distribution as  $t \rightarrow \infty$  to  $N(0, 1)$ , the normal distribution with mean 0 and variance 1.

2. (25 pts)

(a) Suppose that random variables  $X_n \rightarrow X$  in probability and  $f$  is a continuous function. Show that  $f(X_n) \rightarrow f(X)$  in probability.

(b) Let  $X_1, X_2, \dots$  be a sequence of i.i.d. random variables with finite mean  $\mu$  and finite variance. Show that

$$\lim_{n \rightarrow \infty} \binom{n}{2}^{-1} \sum_{1 \leq i < j \leq n} X_i X_j = \mu^2 \quad \text{in probability.}$$

3. (25 pts) Let  $X_n, n \geq 0$ , be a martingale and  $H_n, n \geq 1$ , be a predictable sequence with respect to the filtration  $\mathcal{F}_n, n \geq 0$ . Show that

$$(H \cdot X)_n = \sum_{j=1}^n H_j (X_j - X_{j-1})$$

is a martingale provided  $E[H_n^2(X_n^2 - X_{n-1}^2)] < \infty$  for every  $n$ .

4. (25 pts) Let  $X_n$  be a martingale with respect to the filtration  $\mathcal{F}_n, n \geq 1$ , and  $S < T$  be two stopping times. Consider the relation

$$E[X_T | \mathcal{F}_S] = X_S. \tag{1}$$

(a) Give an example to show that (1) is not true in general.

(b) Show that (1) is valid when either  $T$  is bounded or  $|X_n|$  is uniformly bounded, i.e. either  $T \leq k \in \mathbb{N}$  a.s. or  $|X_n| \leq c \in \mathbb{R}_+$  a.s. for all  $n$ .