

國立臺灣大學數學系
九十五學年度博士班資格考試試題
科目：實分析

2007.06.01

1. (20 pt) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be Riemann integrable over \mathbb{R} . Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be Lebesgue integrable over \mathbb{R} .

(1) Must f be Lebesgue integrable over \mathbb{R} ? (10 pt)

(2) Must g be Riemann integrable over \mathbb{R} ? (10 pt)

Prove or disprove your answers.

2. (20 pt) Let $u \in L^6(\Omega)$ and $v \in L^4(\mathbb{R}) \cap L^6(\mathbb{R})$, where Ω is a bounded domain in $\mathbb{R}^n, n \geq 1$.

(1) Can there exist C_1 a positive constant independent of u such that

$$\left(\int_{\Omega} u^6 \right)^2 \leq C_1 \left(\int_{\Omega} u^4 \right)^3 ?$$

(10 pt)

(2) Can there exist C_2 a positive constant independent of v such that

$$\left(\int_{\mathbb{R}} v^4 \right)^3 \leq C_2 \left(\int_{\mathbb{R}} v^6 \right)^2 ?$$

(10 pt)

Prove or disprove your answers.

3. (20 pt) Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable almost everywhere and satisfy $w, w' \in L^2(\mathbb{R})$, where w' is the derivative of w . Can there exist C a positive constant independent of w such that

$$\int_{\mathbb{R}} (w')^2 + w^2 \geq C \left(\int_{\mathbb{R}} w^4 \right)^{1/2} ?$$

Prove or disprove all your answers.

4. (20 pt) Let $u : B_1 \rightarrow \mathbb{R}$ be a smooth function satisfying $u(x) = 0$, for $|x| = 1$, where B_1 is the unit ball in \mathbb{R}^2 with center at origin. Can there exist a positive constant C independent of u such that

$$\int_{B_1} u^2 dx \leq C \int_{B_1} u_r^2 dx$$

hold? Here (r, θ) is the polar coordinate and u_r is the associated partial derivative. Prove or disprove your answer.

5. (20 pt) Let $\{\nu_j\}_{j=1}^{\infty}$ be a sequence of Radon measures satisfying

$$\|\nu_j\|_{\infty} \leq M, \quad j = 1, 2, 3, \dots,$$

where M is a positive constant independent of j , and the norm $\|\cdot\|_{\infty}$ is defined by

$$\|\nu\|_{\infty} = \sup_{f \in C_0^{\infty}(\mathbb{R}^n)} \frac{\int_{\mathbb{R}^n} f d\nu}{\int_{\mathbb{R}^n} f d\mu}.$$

Here μ is the standard Lebesgue measure, and $C_0^{\infty}(\mathbb{R}^n)$ is the collection of smooth functions with compact support. Can there exist ν_* a Radon measure such that $\nu_j \rightarrow \nu_*$ i.e.

$$\int_{\mathbb{R}^n} f d\nu_j \rightarrow \int_{\mathbb{R}^n} f d\nu_*, \quad \forall f \in C_0^{\infty}(\mathbb{R}^n)$$

(up to a subsequence) as $j \rightarrow \infty$? Prove or disprove your answer.