


2. (a)



$$f(x) = x^2 - x \text{ and } \Delta x = \frac{2-0}{4} = 0.5 \Rightarrow$$

$$R_4 = 0.5f(0.5) + 0.5f(1) + 0.5f(1.5) + 0.5f(2)$$

$$= 0.5(-0.25 + 0 + 0.75 + 2) = 1.25$$

The Riemann sum represents the sum of the areas of the two rectangles above the x -axis minus the area of the rectangle below the x -axis. (The second rectangle vanishes.)

$$(b) \int_0^2 (x^2 - x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x \quad [\Delta x = 2/n \text{ and } x_i = 2i/n]$$


$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{4i^2}{n^2} - \frac{2i}{n} \right) \left(\frac{2}{n} \right) = \lim_{n \rightarrow \infty} \frac{2}{n} \left[\frac{4}{n^2} \sum_{i=1}^n i^2 - \frac{2}{n} \sum_{i=1}^n i \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{8}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} - \frac{4}{n^2} \cdot \frac{n(n+1)}{2} \right] = \lim_{n \rightarrow \infty} \left[\frac{4}{3} \cdot \frac{n+1}{n} \cdot \frac{2n+1}{n} - 2 \cdot \frac{n+1}{n} \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{4}{3} \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right) - 2 \left(1 + \frac{1}{n} \right) \right] = \frac{4}{3} \cdot 1 \cdot 2 - 2 \cdot 1 = \frac{2}{3}$$

$$(c) \int_0^2 (x^2 - x) dx = \left[\frac{1}{3}x^3 - \frac{1}{2}x^2 \right]_0^2 = \left(\frac{8}{3} - 2 \right) = \frac{2}{3}$$

(d) $\int_0^2 (x^2 - x) dx = A_1 - A_2$, where A_1 and A_2 are the areas shown in the diagram.



7. First note that either a or b must be the graph of $\int_0^x f(t) dt$, since $\int_0^0 f(t) dt = 0$, and $c(0) \neq 0$. Now notice that $b > 0$ when c is increasing, and that $c > 0$ when a is increasing. It follows that c is the graph of $f(x)$, b is the graph of $f'(x)$, and a is the graph of $\int_0^x f(t) dt$.

15. Let $u = y^2 + 1$, so $du = 2y dy$ and $y dy = \frac{1}{2} du$. When $y = 0$, $u = 1$; when $y = 1$, $u = 2$. Thus,

$$\int_0^1 y(y^2 + 1)^5 dy = \int_1^2 u^5 \left(\frac{1}{2} du \right) = \frac{1}{2} \left[\frac{1}{6} u^6 \right]_1^2 = \frac{1}{12} (64 - 1) = \frac{63}{12} = \frac{21}{4}.$$

21. $\int_{-\pi/4}^{\pi/4} \frac{t^4 \tan t}{2 + \cos t} dt = 0$ by Theorem 5.5.6(b), since $f(t) = \frac{t^4 \tan t}{2 + \cos t}$ is an odd function.

24. Let $u = \cos x$. Then $du = -\sin x dx$, so $\int \sin x \cos(\cos x) dx = -\int \cos u du = -\sin u + C = -\sin(\cos x) + C$.

27. Since $x^2 - 4 < 0$ for $0 \leq x < 2$ and $x^2 - 4 > 0$ for $2 < x \leq 3$, we have $|x^2 - 4| = -(x^2 - 4) = 4 - x^2$ for $0 \leq x < 2$ and $|x^2 - 4| = x^2 - 4$ for $2 < x \leq 3$. Thus,

$$\begin{aligned} \int_0^3 |x^2 - 4| dx &= \int_0^2 (4 - x^2) dx + \int_2^3 (x^2 - 4) dx = \left[4x - \frac{x^3}{3} \right]_0^2 + \left[\frac{x^3}{3} - 4x \right]_2^3 \\ &= \left(8 - \frac{8}{3} \right) - 0 + (9 - 12) - \left(\frac{8}{3} - 8 \right) = \frac{16}{3} - 3 + \frac{16}{3} = \frac{32}{3} - \frac{9}{3} = \frac{23}{3} \end{aligned}$$

$$37. y = \int_{\sqrt{x}}^x \frac{\cos \theta}{\theta} d\theta = \int_1^x \frac{\cos \theta}{\theta} d\theta + \int_{\sqrt{x}}^1 \frac{\cos \theta}{\theta} d\theta = \int_1^x \frac{\cos \theta}{\theta} d\theta - \int_1^{\sqrt{x}} \frac{\cos \theta}{\theta} d\theta \Rightarrow$$

$$y' = \frac{\cos x}{x} - \frac{\cos \sqrt{x}}{\sqrt{x}} \frac{1}{2\sqrt{x}} = \frac{2 \cos x - \cos \sqrt{x}}{2x}$$

$$41. 0 \leq x \leq 1 \Rightarrow 0 \leq \cos x \leq 1 \Rightarrow x^2 \cos x \leq x^2 \Rightarrow \int_0^1 x^2 \cos x dx \leq \int_0^1 x^2 dx = \frac{1}{3} [x^3]_0^1 = \frac{1}{3} \quad [\text{Property 7}].$$

49. Let $u = 2 \sin \theta$. Then $du = 2 \cos \theta d\theta$ and when $\theta = 0$, $u = 0$; when $\theta = \frac{\pi}{2}$, $u = 2$. Thus,

$$\int_0^{\pi/2} f(2 \sin \theta) \cos \theta d\theta = \int_0^2 f(u) \left(\frac{1}{2} du\right) = \frac{1}{2} \int_0^2 f(u) du = \frac{1}{2} \int_0^2 f(x) dx = \frac{1}{2}(6) = 3.$$

54. Let $F(x) = \int_2^x \sqrt{1+t^3} dt$. Then $F'(2) = \lim_{h \rightarrow 0} \frac{F(2+h) - F(2)}{h} = \lim_{h \rightarrow 0} \frac{1}{h} \int_2^{2+h} \sqrt{1+t^3} dt$, and $F'(x) = \sqrt{1+x^3}$, so

$$\lim_{h \rightarrow 0} \frac{1}{h} \int_2^{2+h} \sqrt{1+t^3} dt = F'(2) = \sqrt{1+2^3} = \sqrt{9} = 3.$$

$$56. \lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(\frac{1}{n}\right)^9 + \left(\frac{2}{n}\right)^9 + \left(\frac{3}{n}\right)^9 + \cdots + \left(\frac{n}{n}\right)^9 \right] = \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{i=1}^n \left(\frac{i}{n}\right)^9 = \int_0^1 x^9 dx = \left[\frac{x^{10}}{10}\right]_0^1 = \frac{1}{10}$$

The limit is based on Riemann sums using right endpoints and subintervals of equal length.