

BESSEL PERIODS AND ANTICYCLOTOMIC p -ADIC SPINOR L -FUNCTIONS

MING-LUN HSIEH AND SHUNSUKE YAMANA

ABSTRACT. We construct the anticyclotomic p -adic L -function that interpolates a square root of central values of twisted spinor L -functions of a quadratic base change of a Siegel cusp form of genus 2 with respect to a paramodular group of square-free level.

CONTENTS

1.	Introduction	1
2.	The basic setting	8
3.	Local Bessel integrals for GSp_4	14
4.	Explicit calculations of Bessel integrals I: new vectors	19
5.	Explicit calculations of Bessel integrals II: degenerate vectors	27
6.	Bessel periods on principal series representations	32
7.	The Iwahori Hecke algebras and the ordinary projector e_{ord}^0	39
8.	Explicit calculations of Bessel integrals III: ordinary vectors	42
9.	Global Bessel periods for GU_2^D	45
10.	Theta elements and p -adic L -functions	51
	References	59

1. INTRODUCTION

The purpose of this article is to carry out the first step towards the analytic side of anticyclotomic Iwasawa theory for Siegel cusp forms by generalizing the works [4, 8, 21] for elliptic cusp forms. Namely, we construct anticyclotomic p -adic L -functions for scalar valued Siegel cusp forms of genus two and weight greater than one with respect to paramodular groups of square-free level and establish the explicit interpolation formulae.

1.1. Anticyclotomic Iwasawa main conjecture. Let $\pi \simeq \otimes'_v \pi_v$ be a unitary irreducible cuspidal automorphic representation of $\mathrm{PGSp}_4(\mathbb{A})$ generated by a scalar valued degree two Siegel cuspidal Hecke eigenform f of weight $\kappa \geq 2$ and paramodular level N , where \mathbb{A} denotes the rational adèle ring. Let

Date: February 12, 2024, 19:29:10.

1991 Mathematics Subject Classification. 11F46, 11F67, 11R23.

Key words and phrases. Böcherer conjecture, spinor L -functions, p -adic L -functions.

\mathfrak{S} be the set of ramified places of the representation π . Fix a prime number $p \notin \mathfrak{S}$ and embeddings $\iota_\infty : \bar{\mathbb{Q}} \hookrightarrow \mathbb{C}$ and $\iota_p : \bar{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$. Let E/\mathbb{Q}_p be a finite extension containing Hecke eigenvalues of f . Thanks to the work of many people (Chai-Faltings, Laumon, Shimura, Taylor and Weissauer), there exists a geometric p -adic Galois representation $\rho_{f,p} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GSp}_4(E)$ such that $\rho_{f,p}$ is unramified outside $\mathfrak{S} \cup \{p\}$ and

$$\det(\mathbf{1}_4 - \iota_\infty \iota_p^{-1} \rho_{f,p}(\text{Frob}_\ell) \ell^{-s})^{-1} = L\left(s - \kappa + \frac{3}{2}, \text{Spn}(\pi_\ell)\right)$$

for $\ell \notin \mathfrak{S} \cup \{p\}$ at least if $\kappa > 2$, where Frob_ℓ is the geometric Frobenius and the right-hand side is the spinor L -factor of π_ℓ . See [25] and [38] for the complete result. Denote by ε_{cyc} the p -adic cyclotomic character. We are interested in the central critical twist

$$\rho_{f,p}^* := \rho_{f,p} \otimes \varepsilon_{\text{cyc}}^{\kappa-1} : \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GSp}_4(E).$$

Let $V_f = E^4$ be the representation space of $\rho_{f,p}^*$. Then V_f is self-dual in the sense that $V_f^\vee(1) \simeq V_f$. We further assume the V_f satisfies the following *Panchishkin condition*: there exists a rank two $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ -invariant subspace $\text{Fil}_p^+ V_f$ of V_f such that $\text{Fil}_p^+ V_f$ has positive Hodge-Tate weights $(\kappa - 1, 1)$ while the quotient $V_f/\text{Fil}_p^+ V_f$ has non-positive Hodge-Tate weights $(0, 2 - \kappa)^1$. Let \mathfrak{o}_E be the ring of integers of E . We shall fix a $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -stable \mathfrak{o}_E -lattice $T_f \subset V_f$ once and for all. Let $A_f = V_f/T_f$ and let $\text{Fil}_p^+ A_f$ be the image of $\text{Fil}_p^+ V_f$ in A_f . For any algebraic extension L over \mathbb{Q} , we consider the (minimal) Selmer group defined by

$$\text{Sel}(A_f/L) := \ker \left\{ \text{H}^1(L, A_f) \rightarrow \prod_{v \nmid p} \text{H}^1(L_v, A_f) \times \prod_{\mathfrak{p}|p} \text{H}^1(L_{\mathfrak{p}}, A_f/\text{Fil}_p^+ A_f) \right\}.$$

Let K be an imaginary quadratic field of discriminant $-\Delta_K < 0$ with integer ring \mathfrak{o}_K and adèle ring \mathbb{A}_K . Denote by K^{ab} the maximal abelian extension over K and by \mathfrak{K}_∞ the composition of all the \mathbb{Z}_p -extensions of K . Take the decomposition $\text{Gal}(\mathfrak{K}_\infty/K) \simeq \Gamma^+ \oplus \Gamma^-$ so that the non-trivial element of $\text{Gal}(K/\mathbb{Q})$ acts on $\Gamma^\pm \simeq \mathbb{Z}_p$ by ± 1 . Let K_∞^\pm be the subfield of \mathfrak{K}_∞ with $\text{Gal}(K_\infty^\pm/K) = \Gamma^\pm$. The \mathbb{Z}_p -extension K_∞^-/K is called anticyclotomic. We consider Iwasawa theory for f over K_∞^- . On the algebraic side, one considers the Pontryagin dual $\text{Sel}(A_f/K_\infty^-)^\vee$ of the Selmer group $\text{Sel}(A_f/K_\infty^-)$, which is known to be a finitely generated $\mathfrak{o}_E[[\Gamma^-]]$ -module. On the analytic side, one expects the existence of the p -adic L -function $L_p(f/K_\infty^-) \in \mathfrak{o}_E[[\Gamma^-]]$ attached to f which interpolates the central values of L -functions associated with $\rho_{f,p}$ twisted by characters of Γ^- , and then one could make the following anticyclotomic Iwasawa main conjecture for Siegel cusp forms.

Conjecture 1.1. The characteristic ideal $\text{char}_{\mathfrak{o}_E[[\Gamma^-]]} \text{Sel}(A_f/K_\infty^-)^\vee$ is generated by $L_p(f/K_\infty^-)$.

¹Here $\mathbb{Q}_p(1)$ has Hodge-Tate weight 1 in our convention.

The main result of this paper is the construction of $L_p(f/K_\infty)$ when f is a paramodular newform of square-free level. Actually, we will construct a square root Θ_f of the anticyclotomic p -adic L -function.

1.2. Paramodular Siegel cusp forms. The paramodular group of level N is defined by

$$\mathbf{K}(N) = \mathrm{Sp}_4(\mathbb{Q}) \cap \begin{pmatrix} \mathbb{Z} & \mathbb{Z} & N^{-1}\mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & N\mathbb{Z} \\ N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \end{pmatrix}.$$

This subgroup is a good analogue of the congruence subgroup $\Gamma_0(N)$ underlying the newform theory for GL_2 (cf. [31]). Put

$$\mathrm{Sym}_g = \{z \in \mathbb{M}_g \mid {}^t z = z\}, \quad \mathfrak{H}_g = \{Z \in \mathrm{Sym}_g(\mathbb{C}) \mid \Im Z > 0\}.$$

Throughout this paper we require N to be square-free. Let π be an irreducible cuspidal automorphic representation of $\mathrm{PGSp}_4(\mathbb{A})$ generated by a paramodular Siegel cuspidal Hecke eigenform

$$f(Z) = \sum_B \mathbf{c}_B(f) e^{2\pi\sqrt{-1}\mathrm{tr}(BZ)}, \quad Z \in \mathfrak{H}_2.$$

of genus 2 and weight κ with respect to $\mathbf{K}(N)$. For each prime $\ell \nmid N$ we write $t_{\ell,1}$ and $t_{\ell,2}$ for the respective eigenvalues of the Hecke operators

$$\ell^{\kappa-3}[\mathbf{K}(N)\mathrm{diag}[1, 1, \ell, \ell]\mathbf{K}(N)], \quad \ell^{2(\kappa-3)}[\mathbf{K}(N)\mathrm{diag}[1, \ell, \ell^2, \ell]\mathbf{K}(N)]$$

acting on f . Let

$$Q_\ell(X) = 1 - t_{1,\ell}X + (\ell t_{2,\ell} + (\ell^3 + \ell)\ell^{2\kappa-6})X^2 - \ell^{2\kappa-3}t_{1,\ell}X^3 + \ell^{4\kappa-6}X^4$$

be the Hecke polynomial of f at ℓ . Then we have

$$Q_\ell(\ell^{-s}) = L\left(s - \kappa + \frac{3}{2}, \mathrm{Spn}(\pi_\ell)\right).$$

We write $\mathrm{Spn}(\pi)$ for the strong lift of π to an automorphic representation of $\mathrm{GL}_4(\mathbb{A})$ and $\mathrm{Spn}(\pi)_K$ for the base change of $\mathrm{Spn}(\pi)$ to $\mathrm{GL}_4(\mathbb{A}_K)$. We consider its L -function twisted by Hecke characters ν

$$L(s, \mathrm{Spn}(\pi)_K \otimes \nu) = \prod_\ell L(s, \mathrm{Spn}(\pi_\ell)_{K_\ell} \otimes \nu_\ell).$$

When ℓ does not divide N and the conductor of ν , the local L -factor is given as follows: If $\ell = \mathfrak{l}$ is inert in K , then

$$L(s, \mathrm{Spn}(\pi_\ell)_{K_\ell} \otimes \nu_\ell) = Q_\ell(\lambda_{\mathfrak{l}}\ell^{3/2-\kappa-s})Q_\ell(-\lambda_{\mathfrak{l}}\ell^{3/2-\kappa-s}),$$

where $\lambda_{\mathfrak{l}}^2 = \nu_\ell(\mathfrak{l})$, if $\ell = \mathfrak{l}^2$ is ramified in K , then

$$L(s, \mathrm{Spn}(\pi_\ell)_{K_\ell} \otimes \nu_\ell) = Q_\ell(\lambda_{\mathfrak{l}}\ell^{3/2-\kappa-s}),$$

where $\lambda_{\mathfrak{l}} = \nu_\ell(\mathfrak{l})$, and if $\ell = \mathfrak{l}_1\mathfrak{l}_2$ is split in K , then

$$L(s, \mathrm{Spn}(\pi_\ell)_{K_\ell} \otimes \nu_\ell) = Q_\ell(\lambda_{\mathfrak{l}_1}\ell^{3/2-\kappa-s})Q_\ell(\lambda_{\mathfrak{l}_2}\ell^{3/2-\kappa-s}).$$

where $\lambda_{\mathfrak{l}_i} = \nu_{\ell}(\mathfrak{l}_i)$ for $i = 1, 2$. The L -factors at prime factors of N are given in (4.5). It is conjectured that there is a bijection between isogeny classes of abelian surfaces A/\mathbb{Q} of conductor N with $\text{End}_{\mathbb{Q}}A = \mathbb{Z}$ and such cusp forms f with rational eigenvalues, up to scalar multiplication (see [42, 6]). In this case T_f is the Tate module $\varprojlim A[p^n]$ and the generalized BSD conjecture predicts that the vanishing order of $L(s, \text{Spn}(\pi)_K \otimes \nu)$ at the center $s = \frac{1}{2}$ coincides the dimension $\dim_{\mathbb{C}}(A(K_{\infty}^-) \otimes \mathbb{C})^{\nu}$ of the ν -eigenspace of the Mordell-Weil group of A for an anticyclotomic character ν of Γ^- .

1.3. The Böcherer conjecture. We construct the anticyclotomic p -adic L -function attached to π over K with explicit evaluation formula for anticyclotomic characters of finite order. The key ingredient of our construction is the *Böcherer conjecture* [5], which is a special case of the refined Gross-Prasad conjecture formulated by Yifeng Liu [26] in full generality.

Let S be a positive definite half-integral symmetric matrix of size 2 with determinant $\frac{\Delta_K}{4}$. The Böcherer conjecture relates the central L -values of π to a square of the Bessel period defined by

$$B_S^{\nu}(\phi) = \int_{K^{\times} \mathbb{A}^{\times} \backslash \mathbb{A}_K^{\times}} \int \phi \left(\begin{pmatrix} t & tz \\ 0 & (\det t) t^{-1} \end{pmatrix} \right) \overline{\mathbf{e}(\text{tr}(Sz))\nu(t)} dz dt,$$

where z is integrated over symmetric matrices of size 2 over \mathbb{A}/\mathbb{Q} , \mathbf{e} denotes the standard additive character on \mathbb{A}/\mathbb{Q} and K^{\times} is identified with the subgroup $\{t \in \text{GL}_2(\mathbb{Q}) \mid {}^t S t = (\det t) S\}$. Let $\alpha_{S, \nu_v}^{\natural}$ be a normalized local Bessel integral. Furusawa and Morimoto [15, Theorem 1.2] have recently proved the Böcherer conjecture.

Theorem 1.2 (Furusawa-Morimoto). *Assume that π is tempered. For every anticyclotomic Hecke character $\nu : K^{\times} \mathbb{C}^{\times} \mathbb{A}^{\times} \backslash \mathbb{A}_K^{\times} \rightarrow \mathbb{C}^{\times}$ of p -power conductor and every nonzero cusp form $\phi = \otimes_v \phi_v \in \pi^D$*

$$\frac{|B_S^{\nu}(\phi)|^2}{(\phi, \phi)} = \xi_{\mathbb{Q}}(2) \xi_{\mathbb{Q}}(4) \frac{\Lambda(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \nu)}{2^{s_{\pi}} \Lambda(1, \pi, \text{ad})} \prod_v \frac{\alpha_{S, \nu_v}^{\natural}(\phi_v, \phi_v)}{(\phi_v, \phi_v)},$$

where $s_{\pi} = 2$ or 1 according as π is endoscopic or not.

1.4. The ordinary hypothesis. The Hecke polynomial $Q_p(X)$ of f at p can be factorized into

$$Q_p(X) = (1 - \alpha_p X)(1 - \beta_p X)(1 - p^{2\kappa-3} \alpha_p^{-1} X)(1 - p^{2\kappa-3} \beta_p^{-1} X)$$

such that

$$0 \leq \text{ord}_p(\iota_p(\alpha_p)) \leq \text{ord}_p(\iota_p(\beta_p)) \leq \kappa - \frac{3}{2}.$$

Let $\alpha_{\mathcal{Q}} := p^{2-\kappa} \alpha_p \beta_p$. In view of [36, Théorème 1], the Panchishkin hypothesis is equivalent to the the following *Klingen p -ordinary hypothesis*:

$$(\mathcal{Q}\text{-ord}) \quad \text{ord}_p(\iota_p(\alpha_{\mathcal{Q}})) = 0.$$

Remark 1.3. The Siegel p -ordinary hypothesis is

$$(\mathcal{P}\text{-ord}) \quad \text{ord}_p \iota_p(\alpha_{\mathcal{P}}) = 0.$$

The condition $(\mathcal{Q}) + (\mathcal{P})$ is referred to as the p -ordinary assumption relative to the Borel subgroup (cf. Remark 10.11).

1.5. Main theorem. Let $\text{rec}_K : K^\times \backslash \mathbb{A}_K^\times \rightarrow \text{Gal}(K^{\text{ab}}/K)$ denote the geometrically normalized reciprocity law map. Put $w_K = \sharp \mathfrak{o}_K^\times$. We view a character $\hat{\nu} : \Gamma^- \rightarrow \bar{\mathbb{Q}}_p^\times$ as a character of the Galois group of K and associate a Hecke character

$$\nu = \iota_\infty \circ \iota_p^{-1} \circ \hat{\nu} \circ \text{rec}_K : \mathbb{A}_K^\times / K^\times \mathbb{A}^\times \rightarrow \mathbb{C}^\times.$$

A character ν as above is usually referred to be anticyclotomic in the sense that ν is trivial on \mathbb{A}^\times . We write $c(\nu)$ for the smallest non-negative integer n such that ν_p is trivial on $\mathfrak{o}_{K_p}^\times \cap (1 + p^n \mathfrak{o}_{K_p})$.

Theorem 1.4. *Assume that N is square-free and that K is split at each prime factor of N . Fix a decomposition $N\mathfrak{o}_K = \mathfrak{N}_0^+ \overline{\mathfrak{N}_0^+}$. If π is tempered and $(\mathcal{Q}\text{-ord})$ holds, then there is an element $\Theta_f \in \mathfrak{o}_E[[\Gamma^-]]$ such that for every finite-order character $\hat{\nu} : \Gamma^- \rightarrow \bar{\mathbb{Q}}_p^\times$ we have the following interpolation formula:*

$$\begin{aligned} \hat{\nu}(\Theta_f)^2 &= \frac{\Lambda(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \nu)}{\Omega_{\pi,1}} \cdot e(\pi_p, \nu_p)^2 \\ &\quad \times \nu^{-1}(\mathfrak{N}_0^+) \cdot \alpha_{\mathcal{P}}^6 \cdot 2^{2\kappa-4} \cdot w_K^2 \Delta_K^{\kappa-1} \cdot N^{-1}, \end{aligned}$$

where $e(\pi_p, \nu_p)$ is the p -adic multiplier defined by

$$\begin{aligned} (c(\nu) > 0) \quad & e(\pi_p, \nu_p) = (p^{\kappa-1} \alpha_{\mathcal{Q}}^{-1})^{c(\nu)}, \\ (c(\nu) = 0, p = \mathfrak{p}_1 \mathfrak{p}_2) \quad & e(\pi_p, \nu_p) = \prod_{i=1}^2 (1 - \alpha_{\mathcal{P}}^{-1} \lambda_{\mathfrak{p}_i} p^{\kappa-2}) (1 - \beta_{\mathcal{P}}^{-1} \lambda_{\mathfrak{p}_i} p^{\kappa-2}), \\ (c(\nu) = 0, p = \mathfrak{p}) \quad & e(\pi_p, \nu_p) = (1 - \alpha_{\mathcal{P}}^{-2} p^{2\kappa-4}) (1 - \beta_{\mathcal{P}}^{-2} p^{2\kappa-4}), \\ (c(\nu) = 0, p = \mathfrak{p}^2) \quad & e(\pi_p, \nu_p) = (1 - \alpha_{\mathcal{P}}^{-1} \lambda_{\mathfrak{p}} p^{\kappa-2}) (1 - \beta_{\mathcal{P}}^{-1} \lambda_{\mathfrak{p}} p^{\kappa-2}) \end{aligned}$$

and the complex number $\Omega_{\pi,1}$ is defined by

$$\Omega_{\pi,1} = \frac{\Lambda(1, \pi, \text{ad})}{\langle f, f \rangle_{\mathbb{K}(N)}}, \quad \langle f, f \rangle_{\mathbb{K}(N)} = \int_{\mathbb{K}(N) \backslash \mathfrak{H}_2} |f(Z)|^2 (\det Y)^{\kappa-3} dZ.$$

Here f is normalized so that all the Fourier coefficients of f are real and contained in E via $\iota_p : \bar{\mathbb{Q}}_p \simeq \mathbb{C}$ and some Fourier coefficient of f is non-vanishing modulo the maximal ideal of \mathfrak{o}_E .

Remark 1.5. (1) The imaginary quadratic field K uniquely determines a factorization $N = N^+ N^-$ with N^+ divisible only by primes that are split in K and N^- divisible only by primes that are inert or

ramified in K . In Theorem 2.5, we actually constructs the theta element $\Theta_f \in \mathfrak{o}_E[[\Gamma^-]]$ under the following *Heegner hypothesis*:

(Heeg) N^- is the product of an even number of primes.

If (Heeg) is not true, then $L(\frac{1}{2}, \text{Spn}(\pi)_{K \otimes \nu}) = 0$ (see Remark 2.6(2)). Theorem 1.4 is its special case where $N = N^+$ and $N^- = 1$.

- (2) Since endoscopic Siegel cusp forms of degree 2 are never paramodular (cf. the proof of Proposition 12.3 of [39]), $s_\pi = 1$.
- (3) If $\kappa > 2$ and π is not a Saito-Kurokawa lift, then π is tempered by Proposition 8.1 of [15].
- (4) The modified Euler factor $e(\pi_p, \nu_p)$ is compatible with the conjectural shape of p -adic L -functions due to Coates and Perrin-Riou. Indeed, let M be the motive over \mathbb{Q} associated with $\rho_{f,p}|_{\text{Gal}(\bar{K}/K)} \otimes \nu$. Then $e(\pi_p, \nu_p)^2$ is the ratio between $\mathcal{L}_p^{(\rho)}(M)$ defined in [9, (18), p. 109] and $L_p(M) = L(\frac{1}{2}, \text{Spn}(\pi_p)_{K_p} \otimes \nu_p)$.

1.6. The construction of Θ_f . We sketch the construction of the theta element Θ_f . Define $K(N, p)$ to be the subgroup which consists of matrices $(b_{ij}) \in K(N)$ such that $b_{21}, b_{31}, b_{32}, b_{34}, b_{41}, b_{42}$ are divisible by p . We define Hecke operators on $S_\kappa(K(N, p))$ by

$$\begin{aligned} [\mathbf{U}_p^{\mathcal{P}} h](Z) &= \sum_B \mathbf{c}_{pB}(h) e^{2\pi\sqrt{-1}\text{ptr}(BZ)}, \\ [\mathbf{U}_p^{\mathcal{Q}} h](Z) &= \sum_{x=1}^p \sum_B \mathbf{c}_{t_{u_p(x)} B u_p(x)}(h) e^{2\pi\sqrt{-1}\text{tr}(BZ)}, \end{aligned}$$

where B runs over positive definite symmetric half-integral matrices of size 2 and $u_p(x) = \begin{pmatrix} p & x \\ 0 & 1 \end{pmatrix}$. We define the p -stabilization $f^\ddagger \in S_\kappa(K(N, p))$ of f with respect to $\alpha_{\mathcal{Q}}$ and $\alpha_{\mathcal{P}}$ by

$$f^\ddagger = \alpha_{\mathcal{P}}^{-3} \alpha_{\mathcal{Q}}^{-1} (\mathbf{U}_p^{\mathcal{Q}} - p^{\kappa-1} \alpha_{\mathcal{P}} \beta_{\mathcal{P}}^{-1}) (\mathbf{U}_p^{\mathcal{P}} - p^{2\kappa-3} \alpha_{\mathcal{P}}^{-1}) (\mathbf{U}_p^{\mathcal{P}} - p^{2\kappa-3} \beta_{\mathcal{P}}^{-1}) (\mathbf{U}_p^{\mathcal{P}} - \beta_{\mathcal{P}}) f.$$

This form f^\ddagger is an eigenform of the operators $\mathbf{U}_p^{\mathcal{Q}}$ and $\mathbf{U}_p^{\mathcal{P}}$ with eigenvalues $\alpha_{\mathcal{Q}}$ and $\alpha_{\mathcal{P}}$, respectively.

We begin with some notation. Let $\mathcal{R} = \begin{pmatrix} \mathbb{Z} & \mathbb{Z} \\ pN\mathbb{Z} & \mathbb{Z} \end{pmatrix}$ be the standard Eichler order of level pN in $D := M_2(\mathbb{Q})$. The group \mathcal{R}^\times acts on the set $\Psi \in \text{Hom}(K, D)$ by $\Psi \cdot r \mapsto r^{-1}\Psi r$. Let

$$\text{CM}(K, D) := \text{Hom}(K, D) / \mathcal{R}^\times$$

be the set of \mathcal{R}^\times -conjugacy classes of field homomorphisms from K to D . For $\Psi \in \text{Hom}(K, D)$, denote by $[\Psi]$ the \mathcal{R}^\times -conjugacy class of Ψ in $\text{CM}(K, D)$. For a positive integer c , let \mathcal{O}_c be the order of K of conductor c and K_c the associated ring class field. The conductor of a homomorphism $\Psi \in \text{Hom}(K, D)$ is the unique positive integer c such that $\Psi^{-1}(\mathcal{R}) = \mathcal{O}_c$. Let $\text{CM}(\mathcal{O}_c, R)$ be

the set of \mathcal{R}^\times -conjugacy classes of homomorphisms $\Psi \in \text{Hom}(K, D)$ of conductor c , which we call the set of CM points of conductor c .

The Galois group $\mathcal{G}_c := \text{Gal}(K_c/K)$ acts on $\text{CM}(\mathcal{O}_c, \mathcal{R})$ in the following manner ([18, p.133]): for $\sigma \in \mathcal{G}_c$ and $[\Psi] \in \text{CM}(\mathcal{O}_c, \mathcal{R})$, write $\sigma = \text{rec}_K(a)$ for some $a \in \widehat{K}^\times$ and decompose $\Psi(a) = \gamma \cdot u$ for some $u \in \widehat{\mathcal{R}}^\times$ and $\gamma \in \text{GL}_2(\mathbb{Q})$ with $\det \gamma > 0$ by strong approximation. The action $[\Psi]^\sigma$ is defined by

$$[\Psi]^\sigma := [\gamma^{-1}\Psi\gamma].$$

To each $\Psi \in \text{Hom}(K, D)$ of conductor c , we associate a unique half-integral symmetric positive definite matrix S_Ψ defined by

$$\pm S_\Psi = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \cdot \Psi(c\sqrt{-\Delta_K}/2).$$

Thus $[\Psi] \mapsto [S_\Psi]$ gives a map from $\text{CM}(\mathcal{O}_c, \mathcal{R})$ to \mathcal{R}^\times -conjugacy classes of primitive half-integral symmetric positive definite matrices. For each non-negative integer n , we will choose special CM points $[\Psi_n] \in \text{CM}(\mathcal{O}_{p^n}, \mathcal{R})$ of conductor p^n , and define the n -th theta element $\tilde{\Theta}_n \in \mathfrak{o}_E[\mathcal{G}_{p^n}]$ by

$$\tilde{\Theta}_n = \alpha_{\mathbb{Q}}^{-n} \sum_{\sigma \in \mathcal{G}_{p^n}} \mathbf{c}_{S_{\Psi_n}^\sigma}(f^\ddagger) \cdot \sigma.$$

The fact that f^\ddagger is an $\mathbf{U}_p^{\mathbb{Q}}$ -eigenform allows us to make a good choice of CM points $\{[\Psi_n]\}_{n=1}^\infty$ such that $\tilde{\Theta}_n$ is norm-compatible, i.e.

$$\Pi_n^{n+1}(\tilde{\Theta}_{n+1}) = \tilde{\Theta}_n$$

under the quotient map $\Pi_n^{n+1} : \mathcal{G}_{p^{n+1}} \rightarrow \mathcal{G}_{p^n}$, and hence we obtain the element $\tilde{\Theta}_\infty := \varprojlim_n \tilde{\Theta}_n \in \mathfrak{o}_E[[\text{Gal}(K_{p^\infty}/K)]]$. The theta element Θ_f is defined

by $\Theta_f = \Pi_{K_\infty^-}^{K_{p^\infty}}(\tilde{\Theta}_\infty)$ via the quotient map $\Pi_{K_\infty^-}^{K_{p^\infty}} : \text{Gal}(K_{p^\infty}/K) \rightarrow \Gamma^-$. For each anticyclotomic character ν of conductor p^n , the interpolation $\hat{\nu}(\Theta_f)$ is essentially the global Bessel period of f^\ddagger with respect to S_n and ν , and the square of the global Bessel period is a product of the central L -value and local Bessel integrals by the Böcherer conjecture.

1.7. The interpolation formula. To obtain the precise interpolation formula of Θ_f , we have to evaluate the local Bessel integrals explicitly. To this end, we will construct a $\text{GSp}_4(\mathbb{Q}_v)$ -equivariant isomorphism $M_v : \pi_v \simeq \pi_v^S$. Choose an element $\mathbf{J} \in \text{GL}_2(\mathbb{Q})$ which satisfies ${}^t\mathbf{J}\mathbf{S}\mathbf{J} = S$ and $\det \mathbf{J} = -1$. Then

$$[\phi_v \otimes \phi'_v \mapsto \alpha_{S, \nu_v}^{\natural}(\phi_v, M_v(\pi_v(\mathbf{t}(\mathbf{J}))\phi'_v))] \in \text{Hom}_{R_S \times R_S}(\pi_v \boxtimes \pi_v, \nu_v^S \boxtimes \nu_v^S),$$

where we put $\mathbf{t}(\mathbf{J}) = \text{diag}[\mathbf{J}, -{}^t\mathbf{J}^{-1}]$ and define the character ν_v^S of $R_S = K_v^\times \rtimes \text{Sym}_2(\mathbb{Q}_v)$ by $\nu_v^S(t, z) = \nu_v(t)\mathbf{e}_v(\text{tr}(Sz))$. The Böcherer conjecture relates the square $B_S^\nu(\phi)^2$ to the product of the central L -value and

$$\alpha_{S, \nu_v}^{\natural}(\phi_v^0, M_v(\pi_v(\mathbf{t}(\mathbf{J}))\phi_v^0)).$$

Our main task is to compute this local factor for a nice test vector $\phi_v^0 \in \pi_v$. If $v \neq p$ and both π_v and ν_v are unramified, then ϕ_v^0 is an unramified vector and the local Bessel integral has been calculated in [26, 10]. If v divides N , then ϕ_v^0 is a paramodular new vector of π_v and its Bessel period will be computed in §4.2. The quaternion case is discussed in §5.3. If $v = \infty$, then ϕ_∞^0 is a lowest weight vector and the computation is done in §5.4. When $\kappa > 2$, the archimedean Bessel integral has been computed in [10] by a method suggested by Kazuki Morimoto. Our computation is different and includes the case $\kappa = 2$.

The calculation of the local integral at $v = p$ is one of the main novelties in this paper. We first construct an ordinary projector $e_{\text{ord},p}^0$ in Section 7 and compute the Bessel integral of $e_{\text{ord},p}^0 \phi_p^0$ in Section 8. To that end, we will construct a local Bessel period $\mathbf{B}_{S,\nu_p}^{\mathcal{W},\pi_p} \in \text{Hom}_{R_S}(\pi_p, \nu_p^S)$ so that $\mathbf{B}_{S,\nu_p}^{\mathcal{W},\pi_p}(e_{\text{ord},p}^0 \phi_p^0)$ is computable. By uniqueness we are led to a functional equation and a factorization

$$\mathbf{B}_{S,\nu_p}^{\mathcal{W},\pi_p^\vee} \circ M_p = c(\pi_p, \nu_p) \mathbf{B}_{S,\nu_p}^{\mathcal{W},\pi_p}, \quad \alpha_{S,\nu_p}^\natural = c'(\pi_p, \nu_p) \mathbf{B}_{S,\nu_p}^{\mathcal{W},\pi_p} \otimes \mathbf{B}_{-S,\nu_p^{-1}}^{\mathcal{W},\pi_p^\vee}.$$

We determine the proportionality constants in Propositions 3.4, 6.5 and 6.6.

Acknowledgement. Yamana is partially supported by JSPS Grant-in-Aid for Scientific Research (C) 18K03210, (C)23K03055 and (B) 19H01778. He would like to thank Michael Harris for very stimulating discussions and Kazuki Morimoto and Hiraku Atobe for helpful discussions. During the preparation of the article, Hsieh is partially supported by NSTC grants 103-2115-M-002-012-MY5, 108-2628-M-001-009-MY4 and 110-2628-M-001-004. This work was done when Yamana stayed in the Max Planck Institute for Mathematics, and he would like to thank the institute for the excellent working environment. This work was partly supported by MEXT Promotion of Distinctive Joint Research Center Program JPMXP0723833165.

2. THE BASIC SETTING

2.1. Notation. Besides the standard symbols $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}, \mathbb{Z}_p, \mathbb{Q}_p$ we denote by \mathbb{N} the set of positive integers, by \mathbb{R}_+^\times the group of strictly positive real numbers and by $\mathbb{C}_p = \widehat{\mathbb{Q}_p}$ the completion of an algebraic closure of the p -adic field \mathbb{Q}_p . If x is a real number, then we put $[x] = \max\{i \in \mathbb{Z} \mid i \leq x\}$. For any finite set A we denote by $\sharp A$ the number of elements in A . For any set X we denote by $\mathbb{1}_X$ the characteristic function of X . When G is a topological group, we write G° for its connected component of the identity. When G is locally compact and abelian, we denote the group of quasi-characters of G by $\Omega(G)$ and the subgroup of unitary characters of G by $\Omega^1(G)$.

Let F be a local field of characteristic zero with normalized absolute value $\omega_F = |\cdot|_F$. We often simply write $|x| = |x|_F$ and $\omega(x) = \omega_F(x)$ for $x \in F^\times$ if its meaning is clear from the context without possible confusion. The group

$\Omega(F^\times)^\circ$ (resp. $\Omega^1(F^\times)^\circ$) consists of homomorphisms of the form ω_F^s with $s \in \mathbb{C}$ (resp. $s \in \sqrt{-1}\mathbb{R}$). Let $\sigma \in \Omega(F^\times)$. Define $\Re\sigma$ as the unique real number such that $\sigma\omega_F^{-\Re\sigma} \in \Omega^1(F^\times)$.

Let F be nonarchimedean. We denote the integer ring of F by \mathfrak{o}_F , the maximal ideal of \mathfrak{o}_F by \mathfrak{p} and the order of the residue field $\mathfrak{o}_F/\mathfrak{p}$ by q and the different of F by \mathfrak{d}_F . Fix a prime element ϖ of \mathfrak{o}_F . When $\sigma \in \Omega(F^\times)^\circ$, we put $L(s, \sigma) = \frac{1}{1-\sigma(\varpi)q^{-s}}$. Otherwise we put $L(s, \sigma) = 1$. We extend $|\cdot|_F$ to fractional ideals of \mathfrak{o}_F by $|\varpi^i|_F = q^{-i}$. Fix a generator d_F of \mathfrak{d}_F and a nontrivial additive character ψ on F . Put $\zeta(s) = \zeta_F(s) = \frac{1}{1-q^{-s}}$. In our later discussion we mostly let ψ be trivial on \mathfrak{o}_F but not trivial on \mathfrak{p}^{-1} . When the residual characteristic of F is p , we define the character ψ^F of F by $\psi^F(x) = e^{-2\pi\sqrt{-1}y}$ with $y \in \mathbb{Q}$ such that $\text{Tr}_{F/\mathbb{Q}_p}(x) - y \in \mathbb{Z}_p$. Let dx be the self-dual Haar measure on F with respect to the pairing $(x, y) \mapsto \psi^F(xy)$. This measure gives \mathfrak{o}_F the volume $|\mathfrak{d}_F|^{1/2}$. The Haar measure $d^\times x$ of F^\times is normalized by $d^\times x = \zeta(1) \frac{dx}{|x|_F}$. When K is a quadratic étale algebra over F , let dt be the quotient measure of the Haar measures of K^\times and F^\times .

For an admissible representation (π, V) of a reductive group G over F we will write π^\vee for its contragredient representation. We occasionally identify the space V with π itself when there is no danger of confusion. When $G = \text{GL}_n(F)$ and $\mu \in \Omega(F^\times)$, we define the representation $\pi \otimes \mu$ on the same space V_π by $(\pi \otimes \mu)(g) = \mu(\det g)\pi(g)$. When Π is an irreducible admissible representation of $\text{GL}_n(F)$, we write Π^K for its base change to $\text{GL}_n(K)$ and write $L(s, \Pi)$ for its Godement-Jacquet L -factor. Given an irreducible admissible representation π of $\text{GSp}_4(F)$, we denote its transfer to $\text{GL}_4(F)$ by $\text{Spn}(\pi)$ and its adjoint L -factor by $L(s, \pi, \text{ad})$. When π is not supercuspidal, these L -parameter and degree 10 L -factor are explicitly computed in Table A.7 of [31] and [3], respectively.

2.2. Quaternion unitary groups. For any ring R we denote by $M_{i,j}(R)$ the set of $i \times j$ -matrices with entries in R and write $M_g(R)$ in place of $M_{g,g}(R)$. The group of all invertible elements of $M_g(R)$ and the set of symmetric matrices of size g with entries in R are denoted by $\text{GL}_g(R)$ and $\text{Sym}_g(R)$, respectively. We sometimes write $R^\times = \text{GL}_1(R)$. The subgroup $B_g(R)$ consists of upper triangular matrices in $\text{GL}_g(R)$. For matrices $B \in \text{Sym}_g(R)$ and $G \in M_{g,m}(R)$ we use the abbreviation $B[G] = {}^tGBG$, where tG is the transpose of G . If A_1, \dots, A_r are square matrices, then $\text{diag}[A_1, \dots, A_r]$ denotes the matrix with A_1, \dots, A_r in the diagonal blocks and 0 in all other blocks. Let $\mathbf{1}_g$ be the identity matrix of degree g . When G is a reductive algebraic group and Z is its center, we write PG for the adjoint group G/Z .

Let D be a quaternion algebra over a field F . We denote by $x \mapsto \bar{x}$ the main involution of D and by ${}^t\bar{A}$ the conjugate transpose of a matrix $A \in M_n(D)$. Let $N_F^D(x) = x\bar{x}$ and $\text{Tr}_F^D(x) = x + \bar{x}$ denote the reduced norm and the reduced trace of $x \in D$. Put $D_- = \{z \in D \mid \bar{z} = -z\}$. We frequently

regard D as an algebraic variety over F and consider the algebraic group GU_2^D which associates to any F -algebra R the group

$$\mathrm{GU}_2^D(R) = \left\{ h \in \mathrm{GL}_2(D \otimes_F R) \mid h \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} {}^t \bar{h} = \lambda(h) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \lambda(h) \in R^\times \right\},$$

where λ is called the similitude character of GU_2^D . We define homomorphisms $\mathbf{m}, \mathbf{t} : D^\times \rightarrow \mathrm{GU}_2^D$, $\mathbf{n} : D_- \rightarrow \mathrm{GU}_2^D$ and $\mathbf{d} : F^\times \rightarrow \mathrm{GU}_2^D$ by

$$\mathbf{m}(A) = \begin{pmatrix} A & 0 \\ 0 & A^{-1} \end{pmatrix}, \quad \mathbf{t}(A) = \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}, \quad \mathbf{n}(z) = \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix}, \quad \mathbf{d}(\lambda) = \begin{pmatrix} 1 & 0 \\ 0 & \lambda \end{pmatrix}$$

and denote the parabolic subgroup of GU_2^D with a Levi factor $\mathbf{d}(F^\times)\mathbf{m}(D^\times)$ and the unipotent radical $\mathbf{n}(D_-)$ by \mathcal{P} .

Fix $S \in D_-$ with $d_0 = S^2 \neq 0$. Put $K = F + FS \subset D$. We choose an element $\mathbf{J} \in D_-$ such that $\mathbf{J}t\mathbf{J}^{-1} = \bar{t}$ for $t \in K$. Then $K \simeq F(\sqrt{d_0})$ and $D = K + K\mathbf{J}$. Let $R_S = \mathbf{t}(K^\times)\mathbf{n}(D_-) \simeq K^\times \ltimes D_-$ be a subgroup of \mathcal{P} .

Let GSp_{2g} be the symplectic similitude group of rank g defined by

$$\mathrm{GSp}_{2g} = \{h \in \mathrm{GL}_{2g} \mid hJ_g {}^t h = \lambda_g(h)J_g, \lambda_g(h) \in \mathrm{GL}_1\}, \quad J_g = \begin{pmatrix} 0 & \mathbf{1}_g \\ -\mathbf{1}_g & 0 \end{pmatrix}.$$

Put $U_2^D = \ker \lambda$ and $\mathrm{Sp}_g = \ker \lambda_g$. We define the homomorphisms

$$\mathbf{m} : \mathrm{GL}_g \times \mathrm{GL}_1 \rightarrow \mathrm{GSp}_{2g}, \quad \mathbf{n} : \mathrm{Sym}_g \rightarrow \mathrm{GSp}_{2g}$$

similarly by

$$\mathbf{m}(A, \lambda) = \begin{pmatrix} A & 0 \\ 0 & \lambda {}^t A^{-1} \end{pmatrix}, \quad \mathbf{n}(z) = \begin{pmatrix} \mathbf{1}_g & z \\ 0 & \mathbf{1}_g \end{pmatrix}.$$

We write

$$\mathbf{m}(A) = \mathbf{m}(A, 1), \quad \mathbf{t}(A) = \mathbf{m}(A, \det A), \quad \mathbf{d}(\lambda) = \mathbf{m}(\mathbf{1}_g, \lambda).$$

Define a maximal parabolic subgroup $\mathcal{P}_g = \mathcal{M}_g N_g$ of GSp_{2g} by

$$\mathcal{M}_g = \{\mathbf{m}(A, \lambda) \mid A \in \mathrm{GL}_g, \lambda \in \mathrm{GL}_1\}, \quad N_g = \{\mathbf{n}(z) \mid z \in \mathrm{Sym}_g\}$$

and a Borel subgroup of GSp_{2g} by

$$\mathcal{B}_g = \{\mathbf{m}(A, \lambda)\mathbf{n}(z) \mid A \in B_g, \lambda \in \mathrm{GL}_1, z \in \mathrm{Sym}_g\}.$$

Note that

$$\mathrm{PGSp}_2 \simeq \mathrm{SO}(2, 1), \quad \mathrm{PGSp}_4 \simeq \mathrm{SO}(3, 2), \quad \mathrm{PGU}_2^D \simeq \mathrm{SO}(4, 1), \quad \mathcal{B}_1 \simeq B_2.$$

We write U_g and \mathcal{U}_g for the unipotent radicals of B_g and \mathcal{B}_g , respectively.

We include the case in which D is the matrix algebra $M_2(F)$. In this case

$$\mathrm{GU}_2^D \simeq \mathrm{GSp}_4, \quad D_- \simeq \mathrm{Sym}_2, \quad \mathcal{P} \simeq \mathcal{P}_2.$$

2.3. Abstract Bessel integrals for GU_2^D . Let F be a local field and (π, V) an irreducible admissible representation of $\mathrm{PGU}_2^D(F)$. Since $\pi \simeq \pi^\vee$, we have a $\mathrm{GU}_2^D(F)$ -invariant bilinear perfect pairing $b : V \times V \rightarrow \mathbb{C}$. Given a pair $\phi_1, \phi_2 \in V$, we define the matrix coefficient $\Phi_{\phi_1, \phi_2} : \mathrm{GU}_2^D(F) \rightarrow \mathbb{C}$ by

$$\Phi_{\phi_1, \phi_2}(g) = \Phi_{\phi_1, \phi_2}^{\mathbf{J}}(g) = b(\pi(g)\phi_1, \pi(\mathbf{m}(\mathbf{J}, -1))\phi_2).$$

Definition 2.1. Let \mathcal{U} be a unipotent algebraic group over a \mathfrak{p} -adic field F and f a smooth function on $\mathcal{U}(F)$. We say that f has a stable integral over $\mathcal{U}(F)$ if there is a compact open subgroup U of $\mathcal{U}(F)$ such that for any open compact subgroup U' containing U

$$\int_{U'} f(z) dz = \int_U f(z) dz.$$

In this case we write $\int_{\mathcal{U}(F)}^{\mathrm{st}} f(z) dz = \int_U f(z) dz$.

We associate to $S \in D_-$ and Λ the character Λ^S of R_S by $\Lambda^S(\mathbf{t}(t)\mathbf{n}(z)) = \Lambda(t)\psi(\mathrm{Tr}_F^D(Sz))$. By [23, 26, 11] the following stable integral of a matrix coefficient exists for each $t \in K^\times$:

$$B_S^\psi(\phi_1, \phi_2, t) = \int_{D_-}^{\mathrm{st}} \Phi_{\phi_1, \phi_2}(\mathbf{n}(z)\mathbf{t}(t)) \overline{\psi(\mathrm{Tr}_F^D(Sz))} dz.$$

Definition 2.2 (abstract Bessel integrals relative to S and Λ). We define

$$B_S^\Lambda(\phi_1, \phi_2) = \int_{F^\times \backslash K^\times} \int_{D_-}^{\mathrm{st}} \Phi_{\phi_1, \phi_2}(\mathbf{n}(z)\mathbf{t}(t)) \overline{\psi(\mathrm{Tr}_F^D(Sz))} \Lambda(t)^{-1} dz dt$$

for $\Lambda \in \Omega(F^\times \backslash K^\times)$ and $\phi_1, \phi_2 \in V$ whenever the integral above converges.

When π is tempered and Λ is unitary, the iterated integral on the right-hand side converges. We give a direct proof for representations of our interest in Lemma 3.3 below. We know that

$$\dim_{\mathbb{C}} \mathrm{Hom}_{R_S}(\pi, \Lambda^S) \leq 1$$

by Corollary 15.3 of [16]. It is important to note that

$$\mathrm{Hom}_{R_S \times R_S}(\pi \boxtimes \pi, \Lambda^S \boxtimes \Lambda^S) = \mathbb{C} B_S^\Lambda,$$

i.e., B_S^Λ is a basis vector of this zero or one-dimensional space.

Remark 2.3. When π is not square-integrable, the Bessel integral may diverge and is defined via regularization. Yifeng Liu [26] has constructed a regularization of the archimedean Bessel integral in general. In §5.4 we will regularize the Bessel integrals of matrix coefficients of lowest weight representations of $\mathrm{GSp}_4(\mathbb{R})$ of scalar weight via a different way.

When $D = \mathrm{M}_2(F)$, we associate to $S \in \mathrm{Sym}_2(F)$ with $\det S \neq 0$ the Bessel integral B_S^Λ in a similar manner.

Remark 2.4. Let $A \in \mathrm{GL}_2(F)$, $\lambda \in F^\times$ and $t \in T$. Put

$$S' = \lambda^{-1}S[A], \quad T' = A^{-1}TA, \quad t' = A^{-1}tA, \quad \mathbf{J}' = A^{-1}\mathbf{J}A.$$

We define $\Lambda' \in \Omega(T')$ by $\Lambda'(t') = \Lambda(At'A^{-1})$. Since

$$\Phi_{\pi(\mathbf{m}(A,\lambda))\phi_1, \pi(\mathbf{m}(A,\lambda))\phi_2}^{\mathbf{J}}(\mathbf{n}(z)\mathbf{t}(t)) = \Phi_{\phi_1, \phi_2}^{\mathbf{J}'}(\mathbf{n}(\lambda A^{-1}z^t A^{-1})\mathbf{t}(t')),$$

it follows that

$$B_S^\Lambda(\pi(\mathbf{m}(A, \lambda))\phi_1, \pi(\mathbf{m}(A, \lambda))\phi_2, t) = |\lambda|^{-3} |\det A|^3 B_{S'}^\Lambda(\phi_1, \phi_2, t').$$

In particular, if we define the additive character ψ_λ by $\psi_\lambda(x) = \psi(\lambda x)$, then

$$\begin{aligned} B_S^\psi(\pi(\mathbf{d}(\lambda))\phi_1, \pi(\mathbf{d}(\lambda))\phi_2, t) &= |\lambda|^{-3} B_S^{\psi\lambda^{-1}}(\phi_1, \phi_2, t), \\ B_S^\Lambda(\pi(\mathbf{m}(A, \lambda))\phi_1, \pi(\mathbf{m}(A, \lambda))\phi_2) &= |\lambda|^{-3} |\det A|^3 B_{S'}^{\Lambda'}(\phi_1, \phi_2). \end{aligned}$$

2.4. Hypotheses. We switch to the global setting. Let $\pi \simeq \otimes'_v \pi_v$ be a unitary irreducible cuspidal tempered automorphic representation of $\mathrm{PGSp}_4(\mathbb{A})$ generated by a scalar valued degree two Siegel cuspidal Hecke eigenform f of weight $\kappa \geq 2$ and square-free paramodular level N .

Fix an imaginary quadratic field K . We decompose N as $N = N^+ N^-$, where each prime factor of N^+ is split in K and each prime factor of N^- is inert or ramified in K . We assume the following *Heegner hypothesis*:

(Heeg) N^- is the product of an even number of primes.

Then there is an indefinite quaternion algebra D that is ramified precisely at the prime factors of N^- . We consider the following inner form of GSp_4 :

$$\mathrm{GU}_2^D = \left\{ g \in \mathrm{GL}_2(D) \mid g \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \bar{t}g = \lambda(g) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\},$$

where $\bar{\cdot}$ denotes the main involution of D . Furusawa and Morimoto [15] have established the Böcherer conjecture more generally for GU_2^D (see Theorem 9.1).

When ρ is a discrete series representation of $\mathrm{GL}_2(\mathbb{Q}_q)$, we write ρ^D for its Jacquet-Langlands lift to $D^\times(\mathbb{Q}_q)$. We define the representation $\pi^D \simeq \otimes'_v \pi_v^D$ of $\mathrm{PGU}_2^D(\mathbb{A})$ by $\pi_v^D \simeq \pi_v$ for $v \nmid N^-$ and by $\pi_q^D \simeq \rho_q^D \rtimes \sigma_q$ for prime factors q of N^- , where we write $\pi_q \simeq \rho_q \rtimes \sigma_q$ (cf. Remark 4.3). When $N^- \neq 1$, we will make use of the inner form transfer on PGSp_4 established in [39, Theorem 11.4], namely *Jacquet-Langlands correspondence* between PGSp_4 and PGU_2^D (cf. [39, Proposition 12.3] and Remarks 4.3 and 5.1):

(JL) The representation π^D occurs in the space of cusp forms on $\mathrm{PGU}_2^D(\mathbb{A})$ with multiplicity one.

2.5. Main theorem. Put $w_K = \sharp \mathfrak{o}_K^\times$. To each character $\hat{\nu} : \Gamma^- \rightarrow \bar{\mathbb{Q}}_p^\times$ we associate a Hecke character $\nu = \iota_\infty \circ \iota_p^{-1} \circ \hat{\nu} \circ \text{rec}_K$ of K , where $\text{rec}_K : K^\times \backslash \mathbb{A}_K^\times \rightarrow \text{Gal}(K^{\text{ab}}/K)$ is the geometrically normalized reciprocity law map. We write $c(\nu)$ for the smallest non-negative integer n such that ν_p is trivial on $\mathfrak{o}_{K_p}^\times \cap (1 + p^n \mathfrak{o}_{K_p})$. Fix a decomposition $N^+ \mathfrak{o}_K = \mathfrak{N}_0^+ \overline{\mathfrak{N}_0^+}$. For each prime factor ℓ of N we write $\epsilon_\ell(f) = \varepsilon(\frac{1}{2}, \text{Spn}(\pi_\ell))$ for the eigenvalue of the Atkin-Lehner involution at ℓ . Put $\epsilon_{N^-}(f) = \prod_{\ell|N^-} \epsilon_\ell(f)$.

Theorem 2.5. *Assume the hypotheses (Heeg) and (\mathcal{Q} -ord) are true for π , K and p . Then there exist an explicitly given complex number $\Omega_{\pi, N^-} \in \mathbb{C}^\times$ and an element $\Theta_f \in \mathfrak{o}_E[[\Gamma^-]]$ with the following interpolation formula*

$$\begin{aligned} \hat{\nu}(\Theta_f)^2 &= \frac{L(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \nu)}{\Omega_{\pi, N^-}} \cdot e(\pi_p, \nu_p)^2 \cdot \nu^{-1}(\mathfrak{N}_0^+) \cdot \alpha_p^6 \\ &\quad \times 2^{2\kappa-3-\ell(\pi)} \cdot w_K^2 \Delta_K^{\kappa-1} \epsilon_{N^-}(f) \cdot N^{-1} \prod_{\ell|(N^-, \Delta_K)} (1 - \epsilon_\ell(f)) \end{aligned}$$

for every finite-order character $\hat{\nu} : \Gamma^- \rightarrow \bar{\mathbb{Q}}_p^\times$, where $e(\pi_p, \nu_p)$ is the p -adic multiplier defined by

$$\begin{aligned} (c(\nu) > 0) \quad & e(\pi_p, \nu_p) = (p^{\kappa-1} \alpha_Q^{-1})^{c(\nu)}, \\ (c(\nu) = 0, p = \mathfrak{p}_1 \mathfrak{p}_2) \quad & e(\pi_p, \nu_p) = \prod_{i=1}^2 (1 - \alpha_p^{-1} \lambda_{\mathfrak{p}_i} p^{\kappa-2}) (1 - \beta_p^{-1} \lambda_{\mathfrak{p}_i} p^{\kappa-2}), \\ (c(\nu) = 0, p = \mathfrak{p}) \quad & e(\pi_p, \nu_p) = (1 - \alpha_p^{-2} p^{2\kappa-4}) (1 - \beta_p^{-2} p^{2\kappa-4}), \\ (c(\nu) = 0, p = \mathfrak{p}^2) \quad & e(\pi_p, \nu_p) = (1 - \alpha_p^{-1} \lambda_{\mathfrak{p}} p^{\kappa-2}) (1 - \beta_p^{-1} \lambda_{\mathfrak{p}} p^{\kappa-2}). \end{aligned}$$

- Remark 2.6.*
- (1) The complex number Ω_{π, N^-} is given in Definition 10.13.
 - (2) Assume that $\epsilon_\ell(f) = -1$ for every prime factor ℓ of (D_K, N^-) . Then $\varepsilon(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \nu) = (-1)^{t(N^-)}$, where $t(N^-)$ is the number of prime factors of N^- . In particular, if (Heeg) is not true, then $L(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \nu) = 0$.
 - (3) We will construct the element $\Theta_f \in E[[\Gamma^-]]$ with the interpolation property without (\mathcal{Q} -ord) more generally for Hilbert-Siegel cusp forms.
 - (4) If ν has infinite order, then so does ν_∞ and so by Theorem 3.10 of [30] $B_S^\nu(\phi) = \alpha_{S, \nu_\infty}^\natural(\varphi, \varphi') = 0$ for all $\phi \in \pi^D$ and $\varphi, \varphi' \in \pi_\infty^D$.
 - (5) If Conjectures 9.4.2 and 9.5.4 of [1] hold for GU_2^D , then $2^{\ell(\pi)}$ coincides with the order of S -group of the Arthur parameter of π^D .

3. LOCAL BESSEL INTEGRALS FOR GSp_4

3.1. Explicit Bessel integrals. Fix $S = \begin{pmatrix} a_0 & \frac{b_0}{2} \\ \frac{b_0}{2} & c_0 \end{pmatrix} \in \mathrm{Sym}_2(F)$. Assume that $\det S \neq 0$. Put

$$\begin{aligned} d_0 &= -4 \det S = b_0^2 - 4a_0c_0, \\ K &= K_S = F(\sqrt{d_0}), \\ T &= T_S = \{A \in \mathrm{GL}_2(F) \mid {}^tASA = (\det A)S\}. \end{aligned}$$

We denote the nontrivial automorphism of K over F by $t \mapsto \bar{t}$. Put $\mathrm{Tr}(t) = \mathrm{Tr}_F^K(t) = t + \bar{t}$ and $\mathrm{N}(t) = \mathrm{N}_F^K(t) = t\bar{t}$ for $t \in K$. We write \mathfrak{r} for the maximal order of K and $\tau_{K/F} : F^\times \rightarrow \{\pm 1\}$ for the character of F^\times whose kernel is $\mathrm{N}(K^\times)$. One can verify that

$$T = \left\{ \begin{pmatrix} x - y\frac{b_0}{2} & -yc_0 \\ ya_0 & x + y\frac{b_0}{2} \end{pmatrix} \mid x, y \in F, x^2 - \frac{d_0}{4}y^2 \neq 0 \right\}.$$

We identify T with K^\times via the map

$$x + \frac{y}{2}\sqrt{d_0} \mapsto \begin{pmatrix} x - y\frac{b_0}{2} & -yc_0 \\ ya_0 & x + y\frac{b_0}{2} \end{pmatrix}.$$

We regard characters of K^\times as those of T .

Let π_0 be an irreducible admissible unitary infinite dimensional representation of $\mathrm{PGL}_2(F)$. It possesses a Whittaker model, i.e., there is a functional

$$0 \neq \mathcal{W} \in \mathrm{Hom}_{U_2}(\pi_0, \psi) = \mathrm{Hom}_{\mathrm{GL}_2(F)}(\pi_0, \mathrm{Ind}_{U_2}^{\mathrm{GL}_2(F)} \psi),$$

which is unique up to scalar multiple. Let $\mathcal{W}(\pi_0, \psi)$ be the space of functions of the form $\mathcal{W}_f(g) = \mathcal{W}(\pi_0(g)f)$ with $f \in \pi_0$. We sometimes identify V_π with $\mathcal{W}(\pi_0, \psi)$. Since $\pi_0 \simeq \pi_0^\vee$, we can define the $\mathrm{GL}_2(F)$ -invariant pairing

$$b_{\mathcal{W}} : \pi_0 \times \pi_0 \rightarrow \mathbb{C}, \quad b_{\mathcal{W}}(f, f') = \int_{F^\times} \mathcal{W}_f(\mathbf{t}(a)) \mathcal{W}_{f'}(\mathbf{t}(-a)) d^\times a.$$

For $\sigma \in \Omega(F^\times)$ we consider the induced representation

$$\pi = I(\pi_0, \sigma) := \mathrm{Ind}_{\mathcal{P}_2}^{\mathrm{GSp}_4(F)}(\pi_0 \otimes \sigma^{-1}) \boxtimes \sigma = \pi_0 \otimes \sigma^{-1} \rtimes \sigma.$$

It is noteworthy that π has trivial central character and can be viewed as a representation of $\mathrm{PGSp}_4(F) \simeq \mathrm{SO}(2, 3)$.

We normalize the Haar measure dz on $\mathrm{Sym}_2(F)$ so that $\mathrm{Sym}_2(\mathfrak{o}_F)$ has volume 1. Choose a right invariant measure dg on $\mathcal{P}_2 \backslash \mathrm{GSp}_4(F)$ such that

$$(3.1) \quad \int_{\mathcal{P}_2 \backslash \mathrm{GSp}_4(F)} f(g) dg = \int_{\mathrm{Sym}_2(F)} f(w_s \mathbf{n}(z)) dz$$

for all $f \in \mathrm{Ind}_{\mathcal{P}_2}^{\mathrm{GSp}_4(F)} \delta_{\mathcal{P}_2}^{1/2}$. We associate to any $\mathrm{GL}_2(F)$ -invariant pairing

$$\mathcal{B} : \pi_0 \times \pi_0 \rightarrow \mathbb{C}$$

the $\mathrm{GSp}_4(F)$ -invariant pairing $\mathcal{B}^\sharp : I(\pi_0, \sigma) \times I(\pi_0, \sigma^{-1}) \rightarrow \mathbb{C}$ by

$$(3.2) \quad \mathcal{B}^\sharp(\phi, \phi') = \int_{\mathcal{P}_2 \backslash \mathrm{GSp}_4(F)} \mathcal{B}(\phi(g), \phi'(g)) \, dg.$$

We use $b_{\mathcal{W}}^\sharp$ to identify $I(\pi_0, \sigma^{-1})$ with the contragredient representation π^\vee .

We shall study the Bessel integral for the representations of the form $I(\pi_0, \sigma)$, and in addition, we will explicitly factorize it into a product of two appropriate local Bessel periods when K is split or π_0 is a principal series representation. In what follows, we fix $D' \in \mathfrak{o}_F$ and put $\theta = \frac{D' + \sqrt{d_0}}{2} \in \mathfrak{r}$. Fix an element $\mathbf{J} \in \mathrm{GL}_2(F)$ such that $t\mathbf{J} = \mathbf{J}\bar{t}$ for $t \in T$. There is no loss of generality by letting

$$(3.3) \quad S = \begin{pmatrix} 1 & -\frac{\mathrm{Tr}(\theta)}{2} \\ -\frac{\mathrm{Tr}(\theta)}{2} & \mathrm{N}(\theta) \end{pmatrix}, \quad \mathbf{J} = \begin{pmatrix} -1 & \mathrm{Tr}(\theta) \\ 0 & 1 \end{pmatrix}$$

thanks to Remark 2.4. The embedding $\iota : K \hookrightarrow \mathrm{M}_2(F)$ attached to this S is

$$(3.4) \quad t = a\theta + b \mapsto \iota(t) = \begin{pmatrix} b + a\mathrm{Tr}(\theta) & -aN(\theta) \\ a & b \end{pmatrix} \quad (a, b \in F).$$

We introduce the intertwining operator

$$M(\pi_0, \sigma) : I(\pi_0, \sigma) \rightarrow I(\pi_0, \sigma^{-1}),$$

defined for $\Re\sigma \ll 0$ by the convergent integral

$$[M(\pi_0, \sigma)\phi](g) = \int_{\mathrm{Sym}_2(F)} \phi(J_2\mathbf{n}(z)g) \, dz,$$

and by meromorphic continuation otherwise. A normalized intertwining operator is defined by setting

$$M^*(\pi_0, \sigma) = \gamma(0, \pi_0 \otimes \sigma^{-1}, \psi)\gamma(0, \sigma^{-2}, \psi)M(\pi_0, \sigma).$$

The character $\psi_S : \mathrm{Sym}_2(F) \rightarrow \mathbb{C}^\times$ is defined by $\psi_S(\mathbf{n}(z)) := \psi(\mathrm{tr}(Sz))$.

Definition 3.1 (explicit Bessel integrals relative to S and Λ). We define

$$J_{S,\Lambda}^{\mathcal{W}} \in \mathrm{Hom}_{R_S \times R_S}(I(\pi_0, \sigma) \boxtimes I(\pi_0, \sigma^{-1}), \Lambda^S \boxtimes \Lambda^S)$$

as in Definition 2.2 by the integral

$$J_{S,\Lambda}^{\mathcal{W}}(\phi, \phi') = \int_{F^\times \backslash K^\times} \int_{\mathrm{Sym}_2(F)}^{\mathrm{st}} b_{\mathcal{W}}^\sharp(\pi(\mathbf{t}(t)\mathbf{n}(z))\phi, \pi^\vee(\mathbf{t}(\mathbf{J}))\phi') \overline{\psi_S(z)\Lambda(t)} \, dz dt$$

for $\phi \in I(\pi_0, \sigma)$ and $\phi' \in I(\pi_0, \sigma^{-1})$. Furthermore we define

$$B_{S,\Lambda}^{\mathcal{W}} \in \mathrm{Hom}_{R_S \times R_S}(I(\pi_0, \sigma) \boxtimes I(\pi_0, \sigma), \Lambda^S \boxtimes \Lambda^S)$$

by

$$B_{S,\Lambda}^{\mathcal{W}}(\phi_1, \phi_2) := J_{S,\Lambda}^{\mathcal{W}}(\phi_1, M^*(\pi_0, \sigma)\phi_2), \quad \phi_1, \phi_2 \in I(\pi_0, \sigma).$$

Clearly, Definition 3.1 is independent of the choice of \mathbf{J} .

3.2. Bessel periods. We have introduced the symmetric matrix S in (3.3). Define matrices $\varsigma \in \mathrm{GL}_2(F)$ and $S' \in \mathrm{Sym}_2(F)$ by

$$\varsigma = \begin{pmatrix} 1 & -\bar{\theta} \\ -1 & \theta \end{pmatrix}, \quad S' = \begin{pmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix}$$

Then

$$\iota_\varsigma(t) := \varsigma t \varsigma^{-1} = \mathrm{diag}[t, \bar{t}], \quad {}^t \varsigma S' \varsigma = S.$$

If K/F is not split, then we set

$$\varsigma = \mathbf{1}_2, \quad S' = S, \quad \iota_\varsigma = \iota.$$

Fix a ψ -Whittaker functional \mathcal{W} on π_0 . In order to investigate the Bessel integral $J_{S, \Lambda}^{\mathcal{W}}$ we will explicitly construct toric and Bessel periods

$$\mathbf{T}_\Lambda^{\mathcal{W}} \in \mathrm{Hom}_{\varsigma T \varsigma^{-1}}(\pi_0, \Lambda), \quad \mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma} \in \mathrm{Hom}_{R_{S'}}(I(\pi_0, \sigma), \Lambda_{S'}).$$

We define the toric period of $f \in \pi_0$ by

$$\mathbf{T}_\Lambda^{\mathcal{W}}(f) = \int_{F^\times \backslash K^\times} \mathcal{W}(\pi_0(\iota_\varsigma(t))f) \Lambda(t)^{-1} dt.$$

This integral is absolutely convergent and gives rise to a nonzero K^\times -invariant functional on π_0 for any unitary generic representation π_0 .

Let w_s be the Weyl element given by

$$(3.5) \quad w_s = \left(\begin{array}{cc|cc} & & 0 & 1 \\ & & -1 & 0 \\ \hline 0 & -1 & & \\ 1 & 0 & & \end{array} \right) = \mathbf{m} \left(\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right) s_2 s_1 s_2 \in \mathrm{GSp}_4(F).$$

It is important to note that

$$w_s \mathbf{t}(A) = \mathbf{t}(A) w_s.$$

Definition 3.2. We define the Bessel period of $\phi \in I(\pi_0, \sigma)$ by

$$\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}(\phi) = \int_{F^\times \backslash K^\times} \int_{\mathrm{Sym}_2(F)}^{\mathrm{st}} \mathcal{W}(\phi(w_s \mathbf{n}(z) \mathbf{t}(\iota_\varsigma(t)))) \psi_{S'}(-z) \Lambda(t)^{-1} dz dt.$$

For $i \in \mathbb{N}$ we put

$$\mathrm{Sym}_2^i = \left\{ \begin{pmatrix} x & y \\ y & w \end{pmatrix} \mid x, w \in F, y \in \mathfrak{p}^{-i} \right\}.$$

Lemma 3.3. *Assume that K/F is split. Write $\Lambda = (\Lambda_0, \Lambda_0^{-1})$. Let $\phi \in \pi$. Take $i \in \mathbb{N}$ such that $\pi(k)\phi = \phi$ for elements $k \in \mathbb{I}$ which satisfy $k - \mathbf{1}_4 \in \mathrm{M}_4(\mathfrak{p}^i)$. Suppose that ψ has order 0. If $\Re \sigma < \frac{1}{2}$, then the double integral*

$$\int_{F^\times} \int_{\mathrm{Sym}_2^i} \mathcal{W}(\phi(w_s \mathbf{n}(z) \mathbf{t}(\mathrm{diag}[a, 1]))) \psi_{S'}(-z) \Lambda_0(a)^{-1} dz d^\times a$$

is absolutely convergent and equal to $\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}(\phi)$.

Proof. Conjugating ϕ by $\mathbf{d}(\lambda)$ with $\lambda \in 1 + \mathfrak{p}^i$ and making a change of variables, we see that $\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}(\phi)$ is equal to

$$\int_{F^\times} \int_{F^3}^{\text{st}} \mathcal{W} \left(\phi \left(w_s \mathbf{n} \left(\begin{pmatrix} x & y \\ y & w \end{pmatrix} \right) \mathbf{t}(\text{diag}[a, 1]) \right) \right) \psi(\lambda y) \Lambda_0(a)^{-1} dx dy dw da.$$

Integrating both sides of this equality over $\lambda \in 1 + \mathfrak{p}^i$, we get

$$\int_{F^\times} \int_{\mathfrak{p}^{-i}} \int_{F^2}^{\text{st}} \mathcal{W} \left(\phi \left(w_s \mathbf{n} \left(\begin{pmatrix} x & y \\ y & w \end{pmatrix} \right) \mathbf{t}(\text{diag}[a, 1]) \right) \right) \frac{\psi(y)}{\Lambda_0(a)} dx dw dy d^\times a.$$

The set Sym_2^i is stable under the action of elements $\text{diag}[a, 1]$. It suffices to check that the double integral

$$\int_{F^2} \int_{F^\times} \left| \mathcal{W} \left(\phi \left(\mathbf{t}(\text{diag}[a, 1]) w_s \mathbf{n} \left(\begin{pmatrix} x & y \\ y & w \end{pmatrix} \right) \right) \right) \right| d^\times a dx dw$$

is convergent for every $y \in \mathfrak{p}^{-i}$.

Observe that

$$\mathbf{n}(z) = \begin{pmatrix} \mathbf{1}_2 & \\ z^{-1} & \mathbf{1}_2 \end{pmatrix} \begin{pmatrix} z & \\ & z^{-1} \end{pmatrix} \begin{pmatrix} & \mathbf{1}_2 \\ -\mathbf{1}_2 & \end{pmatrix} \begin{pmatrix} \mathbf{1}_2 & \\ z^{-1} & \mathbf{1}_2 \end{pmatrix}.$$

Since $w_s \mathbf{t}(A) w_s^{-1} = \mathbf{t}(A)$, we get

$$(3.6) \quad w_s \mathbf{n}(z) \in N_2 Z_2 \mathbf{d}(\det z) \mathbf{t}(z) w_s \begin{pmatrix} & \mathbf{1}_2 \\ -\mathbf{1}_2 & \end{pmatrix} \begin{pmatrix} \mathbf{1}_2 & \\ z^{-1} & \mathbf{1}_2 \end{pmatrix}.$$

Let $z = \begin{pmatrix} x & y \\ y & w \end{pmatrix}$. The inner integral converges as π_0 is unitary and generic.

Clearly, it depends only on $x + \mathfrak{p}^i$ and $w + \mathfrak{p}^i$. We may therefore assume that $x, w \notin \mathfrak{p}^{i+1}$. If $x \notin \mathfrak{p}^{-3i}$, then since $\text{ord } x < -3i$, $\text{ord } y \geq -i$ and $\text{ord } w \leq i$,

$$c_z = -z^{-1} = -(\det z)^{-1} \begin{pmatrix} w & -y \\ -y & x \end{pmatrix} \in \begin{pmatrix} \mathfrak{p}^{3i+1} & \mathfrak{p}^{2i+1} \\ \mathfrak{p}^{2i+1} & \mathfrak{p}^{-i} \end{pmatrix},$$

$$c'_z = \text{diag}[x^{-1}, w^{-1}]z = \begin{pmatrix} 1 & \frac{y}{x} \\ \frac{y}{w} & 1 \end{pmatrix} \in \begin{pmatrix} 1 & \mathfrak{p}^{2i+1} \\ \mathfrak{p}^{-2i} & 1 \end{pmatrix}$$

and hence

$$|\mathcal{W}(\phi(\mathbf{t}(\text{diag}[a, 1]) w_s \mathbf{n}(z)))| = |xw|_F^{\Re\sigma - 3/2} |\mathcal{W}(\phi(\mathbf{t}(\text{diag}[ax, w] c'_z) w_s \mathbf{n}(c_z) J_2))|$$

where there is a compact set K_i of $\text{GL}_2(F)$ such that $c'_z \in K_i$. Put

$$C_i = \sup_{c \in \text{Sym}_2(\mathfrak{p}^{-i}), c' \in K_i} \int_{F^\times} |\mathcal{W}(\phi(\mathbf{t}(\text{diag}[a, 1] c') w_s \mathbf{n}(c) J_2))| d^\times a.$$

We therefore conclude that

$$\int_{F \setminus \mathfrak{p}^{-3i}} \int_F \int_{F^\times} \left| \mathcal{W} \left(\phi \left(\mathbf{t}(\text{diag}[a, 1]) w_s \mathbf{n} \left(\begin{pmatrix} x & y \\ y & w \end{pmatrix} \right) \right) \right) \right| d^\times a dx dw$$

$$\leq C_i \sum_{n=3i}^{\infty} \sum_{m=-i}^{\infty} q^{(n+m)(2\Re\sigma-1)/2}.$$

The last summation clearly converges. \square

It is easy to see that $\mathbf{B}_{S',\Lambda}^{\mathcal{W},\sigma}$ is $R_{S'}$ -invariant and

$$(3.7) \quad \mathbf{B}_{S',\Lambda}^{\mathcal{W},\sigma}(\phi) = \lim_{i \rightarrow \infty} \int_{\text{Sym}_2^i} \mathbf{T}_{\Lambda}^{\mathcal{W}}(\phi(w_s \mathbf{n}(z))) \psi_{S'}(-z) dz.$$

3.3. Factorization of $J_{S,\Lambda}^{\mathcal{W}}$.

Proposition 3.4. *If K/F is split, then*

$$J_{S,\Lambda}^{\mathcal{W}}(\phi, \phi') = \Lambda_0(-1) \mathbf{B}_{S',\Lambda}^{\mathcal{W},\sigma}(\pi(\mathbf{m}(\varsigma))\phi) \mathbf{B}_{-S',\Lambda^{-1}}^{\mathcal{W},\sigma^{-1}}(\pi^\vee(\mathbf{m}(\varsigma)\mathbf{t}(\mathbf{J}))\phi')$$

for any $\phi \in \pi = I(\pi_0, \sigma)$ and $\phi' \in \pi^\vee = I(\pi_0, \sigma^{-1})$.

Proof. For our choice of the measure dg on $\mathcal{P}_2 \backslash \text{GSp}_4(F)$ we have

$$\begin{aligned} & \int_{\text{Sym}_2(F)}^{\text{st}} \mathcal{B}^\sharp(\pi(\mathbf{n}(z))\phi_1, \pi^\vee(\mathbf{t}(\mathbf{J}))\phi_2) \psi_S(-z) dz \\ &= \int_{\text{Sym}_2(F)}^{\text{st}} \int_{\text{Sym}_2(F)} \mathcal{B}(\phi_1(w_s \mathbf{n}(z') + z), \phi_2(w_s \mathbf{n}(z')\mathbf{t}(\mathbf{J}))) \psi_S(-z) dz' dz \end{aligned}$$

(see (3.1) and (3.2)). Set $\phi'_1 = \pi(\mathbf{m}(\varsigma))\phi_1$ and $\phi'_2 = \pi^\vee(\mathbf{m}(\varsigma)\mathbf{t}(\mathbf{J}))\phi_2$. Take sufficiently large i . Then the right hand side is equal to

$$\int_{\text{Sym}_2^i} \int_{\text{Sym}_2^i} \mathcal{B}(\phi'_1(w_s \mathbf{n}(z_1)), \phi'_2(w_s \mathbf{n}(-z_2))) \psi_{S'}(-z_1 - z_2) dz_1 dz_2$$

(see the proof of Lemma 3.3). The triple integral

$$\begin{aligned} & \int_{F^\times \backslash K^\times} \int_{\text{Sym}_2^i} \int_{\text{Sym}_2^i} |b_{\mathcal{W}}(\pi_0(\iota_\varsigma(t))\phi'_1(w_s \mathbf{n}(z_1)), \pi_0(\mathbf{J})\phi'_2(w_s \mathbf{n}(z_2)))| dz_1 dz_2 dt \\ &= \prod_{j=1}^2 \int_{\text{Sym}_2^i} \int_{F^\times} |\mathcal{W}(\pi_0(\text{diag}[a, 1])\phi''_j(w_s \mathbf{n}(z_j)))| da dz_j \end{aligned}$$

is convergent in view of Lemma 3.3, which justifies our formal manipulation.

Here we put $\phi''_1 = \phi'_1$ and $\phi''_2 = \pi_0(\mathbf{J})\phi'_2$.

Now we have the identity

$$\begin{aligned} & \int_{\text{Sym}_2(F)}^{\text{st}} b_{\mathcal{W}}^\sharp(\pi(\mathbf{t}(t)\mathbf{n}(z))\phi_1, \pi^\vee(\mathbf{t}(\mathbf{J}))\phi_2) \psi_S(-z) dz \\ &= \int_{\text{Sym}_2^i} \int_{\text{Sym}_2^i} b_{\mathcal{W}}(\pi_0(\iota_\varsigma(t))\phi'_1(w_s \mathbf{n}(z_1)), \phi'_2(w_s \mathbf{n}(z_2))) \psi_{S'}(-z_1 + z_2) dz_1 dz_2. \end{aligned}$$

Integrating over $t \in F^\times \backslash K^\times$ and changing the order of integration, we get

$$\begin{aligned} J_{S,\Lambda}^{\mathcal{W}}(\phi_1, \phi_2) &= \int_{\text{Sym}_2^i} \int_{\text{Sym}_2^i} dz_1 dz_2 \psi_{S'}(z_2 - z_1) \\ &\quad \times \int_{F^\times \backslash K^\times} b_{\mathcal{W}}(\pi_0(\iota_\varsigma(t))\phi'_1(w_s \mathbf{n}(z_1)), \phi'_2(w_s \mathbf{n}(z_2))) dt. \end{aligned}$$

Put $f'_1 = \phi'_1(w_s \mathbf{n}(z_1))$ and $f'_2 = \phi'_2(w_s \mathbf{n}(z_2))$. Then the inner integral equals

$$\int_{F^\times} \int_{F^\times} \mathcal{W}_{f'_1}(\mathbf{t}(ba)) \mathcal{W}_{f'_2}(\mathbf{t}(-b)) \Lambda_0(a)^{-1} d^\times b d^\times a = \Lambda_0(-1) \mathbf{T}_\Lambda^{\mathcal{W}}(f'_1) \mathbf{T}_{\Lambda^{-1}}^{\mathcal{W}}(f'_2).$$

We conclude that $\Lambda_0(-1) J_{S, \Lambda}^{\mathcal{W}}(\phi_1, \phi_2)$ is equal to

$$\int_{\text{Sym}_2^i} \int_{\text{Sym}_2^i} \mathbf{T}_\Lambda^{\mathcal{W}}(\phi'_1(w_s \mathbf{n}(z_1))) \mathbf{T}_{\Lambda^{-1}}^{\mathcal{W}}(\phi'_2(w_s \mathbf{n}(z_2))) \psi_{S'}(-z_1 + z_2) dz_1 dz_2,$$

which completes our proof by (3.7). \square

Remark 3.5. Let K/F be split. Then $T_\Lambda^{\mathcal{W}}(f) = \mathcal{Z}(\mathcal{W}_f \otimes \Lambda_0^{-1}, \frac{1}{2})$ is the zeta integral for $\pi_0 \otimes \Lambda_0^{-1}$. Since ${}_\varsigma \mathbf{J}_\varsigma^{-1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, for every $f \in \pi_0$

$$\mathbf{T}_{\Lambda^{-1}}^{\mathcal{W}}(\pi_0({}_\varsigma \mathbf{J}_\varsigma^{-1})f) = \Lambda_0(-1) \gamma \left(\frac{1}{2}, \pi_0 \otimes \Lambda_0^{-1}, \psi \right) \mathbf{T}_\Lambda^{\mathcal{W}}(\pi_0(\text{diag}[-1, 1])f)$$

by the functional equation in Hecke's theory and hence

$$\mathbf{B}_{S', \Lambda^{-1}}^{\mathcal{W}, \sigma}(\pi(\mathbf{m}({}_\varsigma \mathbf{J}_\varsigma^{-1}))\phi) = \frac{\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}(\pi(\mathbf{m}(\text{diag}[-1, 1]))\phi)}{\Lambda_0(-1) \gamma \left(\frac{1}{2}, \pi_0 \otimes \Lambda_0, \psi \right)}, \quad \phi \in \pi = I(\pi_0, \sigma).$$

4. EXPLICIT CALCULATIONS OF BESSEL INTEGRALS I: NEW VECTORS

Let π be an irreducible admissible representation of the form $I(\pi_0, \sigma)$, where π_0 is an irreducible unramified unitary principal series representation of $\text{PGL}_2(F)$ or the Steinberg representation twisted by an unramified quadratic character of F^\times and $\sigma \in \Omega(F^\times)^\circ$. Let $\phi_\sigma \in \pi$ be a new vector, i.e., ϕ_σ is a spherical vector in the former case and ϕ_σ is a paramodular vector in the latter case. Our task in Sections 4, 5 and 8 is to compute

$$\mathbb{B}_S^\Lambda(H) := \frac{B_S^\Lambda(H\phi_\sigma, H\phi_\sigma)}{b(\phi_\sigma, \phi_\sigma)} = \frac{B_{S, \Lambda}^{\mathcal{W}}(H\phi_\sigma, H\phi_\sigma)}{b_{\mathcal{W}}^\#(\phi_\sigma, M^*(\pi_0, \sigma)\phi_\sigma)}$$

for some Hecke operator H on $\text{GSp}_4(F)$, where B_S^Λ and $B_{S, \Lambda}^{\mathcal{W}}$ are the Bessel integrals defined in Definitions 2.2 and 3.1. Since there exists a constant $m(\pi_0, \sigma)$ such that

$$M^*(\pi_0, \sigma)\phi_\sigma = m(\pi_0, \sigma)\phi_{\sigma^{-1}},$$

it follows from Proposition 3.4 that

$$(4.1) \quad \mathbb{B}_S^\Lambda(\pi(\mathbf{m}(\varsigma^{-1}))) = \Lambda_0(-1) \frac{\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}(\phi_\sigma) \mathbf{B}_{-S', \Lambda^{-1}}^{\mathcal{W}, \sigma^{-1}}(\pi^\vee(\mathbf{t}(\varsigma \mathbf{J}_\varsigma^{-1}))\phi_{\sigma^{-1}})}{b_{\mathcal{W}}^\#(\phi_\sigma, \phi_{\sigma^{-1}})}.$$

Remark 4.1. Proposition 6.9 below gives

$$m(\pi_0, \sigma) = \gamma \left(\frac{1}{2}, \sigma_K^{-1} \Lambda, \psi_K \right) \frac{\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}(\phi_\sigma)}{\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma^{-1}}(\phi_{\sigma^{-1}})}.$$

4.1. **The unramified case.** The ratio $\frac{J_{S,\Lambda}^{\mathcal{W}}(\phi_\sigma, \phi_{\sigma^{-1}})}{b_{\mathcal{W}}^\sharp(\phi_\sigma, \phi_{\sigma^{-1}})}$ has been computed by Liu in Theorem 2.2 of [26] for split or unramified extensions K of F , and extended to ramified extensions in [10]. Since $\mathbb{B}_S^\Lambda(\text{Id}) = \frac{J_{S,\Lambda}^{\mathcal{W}}(\phi_\sigma, \phi_{\sigma^{-1}})}{b_{\mathcal{W}}^\sharp(\phi_\sigma, \phi_{\sigma^{-1}})}$, we obtain the following result:

Theorem 4.2 ([26, 10]). *Assume the following conditions*

- both π and Λ are unramified, unitary and generic;
- $a_0, b_0 \in \mathfrak{o}_F$; $c_0 \in \mathfrak{o}_F^\times$; $-4 \det S$ generates \mathfrak{d}_F ; ψ has order 0;
- When the residual characteristic of F is 2, we suppose that $F = \mathbb{Q}_2$;

If $\phi^0 \in V$ is $\text{GSp}_4(\mathfrak{o}_F)$ -invariant, then

$$\mathbb{B}_S^\Lambda(\text{Id}) = \frac{|\mathfrak{d}_K|_K^{1/2} \zeta(2) \zeta(4) L(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \Lambda)}{|\mathfrak{d}_F|^{1/2} L(1, \tau_{K/F}) L(1, \pi, \text{ad})}.$$

4.2. **The paramodular case.** The paramodular group $\text{K}(\mathfrak{p})$ is the subgroup of $k \in \text{GSp}_4(F)$ such that $\lambda(k)$ is in \mathfrak{o}_F^\times and

$$(4.2) \quad k \in \begin{pmatrix} \mathfrak{o}_F & \mathfrak{o}_F & \mathfrak{p}^{-1} & \mathfrak{o}_F \\ \mathfrak{p} & \mathfrak{o}_F & \mathfrak{o}_F & \mathfrak{o}_F \\ \mathfrak{p} & \mathfrak{p} & \mathfrak{o}_F & \mathfrak{p} \\ \mathfrak{p} & \mathfrak{o}_F & \mathfrak{o}_F & \mathfrak{o}_F \end{pmatrix}.$$

Proposition 5.1.2 of [31] gives the Iwasawa decomposition relative to $\text{K}(\mathfrak{p})$

$$\text{GSp}_4(F) = \mathcal{P}_2(F) \text{K}(\mathfrak{p}).$$

Define the Iwahori subgroup of $\text{GL}_2(\mathfrak{o}_F)$ by

$$\mathfrak{I} = \left\{ A \in \text{GL}_2(\mathfrak{o}_F) \mid A \equiv \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{\mathfrak{p}} \right\}.$$

Given $\mu \in \Omega(F^\times)$, we denote by $I_1(\mu) = \mu \times \mu^{-1}$ the principal series representation of $\text{PGL}_2(F)$. We write $\text{St} \subset I_1(\omega_F^{1/2})$ for the Steinberg representation. As is well-known, its subspace of \mathfrak{I} -invariant vectors is one-dimensional. Take $\sigma \in \Omega^1(F^\times)^\circ$ and an unramified quadratic character ε of F^\times . Let

$$\pi_0 \simeq \text{St} \otimes \varepsilon, \quad \pi \simeq I(\pi_0, \sigma).$$

Put $\mu = \varepsilon \omega_F^{1/2}$. Then

$$\pi_0 \subset I_1(\mu), \quad \pi \subset I(I_1(\mu), \sigma).$$

Remark 4.3. The representations $I(\text{St} \otimes \varepsilon, \sigma)$ are called type IIa in [31]. Their minimal paramodular level is \mathfrak{p} by Table A.12 of [31], namely, the $\text{K}(\mathfrak{p})$ -invariant subspace $\pi^{\text{K}(\mathfrak{p})}$ is one-dimensional. Representations of type IIa (with unramified ε and σ) are only the tempered representations of paramodular level \mathfrak{p} by Tables A.12, A.13 in loc. cit.

Let $f_{\pi_0} \in \pi_0^{\mathfrak{J}}$ be such that

$$\mathcal{W}_{f_{\pi_0}}(\mathbf{t}(a)) = \varepsilon(a)|a|_F \mathbf{1}_{\mathfrak{o}_F}(a).$$

The Iwasawa decomposition allows us to define $\phi_\sigma \in \pi^{\mathbf{K}(\mathfrak{p})}$ by

$$\phi_\sigma(\mathbf{n}(z)\mathbf{d}(\lambda)\mathbf{t}(A)k) = \sigma(\lambda)|\lambda|_F^{-3/2} f_{\pi_0}(A)$$

for $z \in \text{Sym}_2(F)$, $\lambda \in F^\times$, $A \in \text{GL}_2(F)$ and $k \in \mathbf{K}(\mathfrak{p})$.

Since π has no Bessel model relative to the trivial character of K^\times if K/F is the unramified quadratic extension, we assume that $K = Fe_1 \oplus Fe_2$ is split and let $S' = -\begin{pmatrix} 0 & 1 \\ \frac{1}{2} & 0 \end{pmatrix}$ throughout this subsection. Let $\Lambda = (\Lambda_0, \Lambda_0^{-1})$ be an unramified character of $K^\times = F^\times \times F^\times$. Put

$$\gamma = \sigma(\varpi), \quad \epsilon = -\varepsilon(\varpi), \quad \delta = \Lambda_0(\varpi).$$

Define a function $\mathbf{T}' : \text{PGL}_2(F) \rightarrow \mathbb{C}$ by

$$\mathbf{T}'(A) = \mathbf{T}_\Lambda(\pi_0(\mathbf{t}(A))f_{\pi_0}).$$

Observe that for $A \in \text{GL}_2(F)$, $a, b \in F^\times$ and $k \in \mathfrak{J}$

$$(4.3) \quad \mathbf{T}'(A\eta_0) = \epsilon\mathbf{T}'(A), \quad \mathbf{T}'(\text{diag}[a, b]Ak) = \Lambda_0(ab^{-1})\mathbf{T}'(A).$$

In particular, the value $\mathbf{T}'(\mathbf{n}(x))$ depends only on $\mathfrak{o}_F^\times x + \mathfrak{o}_F$. We will write

$$T'(m) = \mathbf{T}'(\mathbf{n}(\varpi^{-m}))$$

for non-negative integers m . For $x, y, w \in F$ we put

$$\mathbf{T}(x, y, w) = \mathbf{T}_\Lambda \left(\phi_\sigma \left(w_s \mathbf{n} \left(\begin{pmatrix} x & y \\ y & w \end{pmatrix} \right) \right) \right).$$

Recall the Bessel period

$$\mathbf{B}_{S', \Lambda}^\sigma(\phi) = \int_{\text{Sym}_2(F)} \mathbf{T}_\Lambda(\phi(w_s \mathbf{n}(z))) \psi_{S'}(-z) dz.$$

Conjugating ϕ_σ by $\mathbf{d}(\lambda)$ with $\lambda \in \mathfrak{o}_F^\times$, we get

$$\mathbf{B}_{S', \Lambda}^\sigma(\phi_\sigma) = \int_{F^3} \mathbf{T}(x, y, w) \psi(-\lambda y) dx dy dz$$

by a change of variables. Choosing $\lambda = 1 + u$ with $u \in \mathfrak{p}$ and integrating both sides of this equality over $u \in \mathfrak{p}$, we see that

$$\mathbf{B}_{S', \Lambda}^\sigma(\phi_\sigma) = \int_{\mathfrak{p}^{-1}} \int_{F^2} \mathbf{T}(x, y, w) \psi(-y) dx dw dy.$$

We define the function $\mathbf{v} : F \rightarrow \mathbb{N} \cup \{0\}$ via

$$q^{\mathbf{v}(x)} = [x\mathfrak{o}_F + \mathfrak{o}_F : \mathfrak{o}_F].$$

Lemma 4.4. *Let $z = \begin{pmatrix} x & y \\ y & w \end{pmatrix} \in \text{Sym}_2(F)$. Assume that the following conditions are satisfied:*

- $\mathbf{v}(y) \leq 1$.

- When $\mathbf{v}(y) = 0$, we suppose that $x \notin \mathfrak{o}_F$ and $w \notin \mathfrak{p}$.
- When $\mathbf{v}(y) = 1$ and $\mathbf{v}(w) = 0$, we suppose that if $\mathbf{v}(x) \geq 3$, then $w \in \mathfrak{o}_F^\times$, while if $\mathbf{v}(x) \leq 2$, then $w = 0$.
- When $\mathbf{v}(y) = 1$ and $\mathbf{v}(x) \leq 1$, we suppose that if $\mathbf{v}(w) \geq 2$, then $\mathbf{v}(x) = 1$, while if $\mathbf{v}(w) \leq 1$, then $x = 0$.

Then $z \in \mathrm{GL}_2(F)$ and

$$w_s \mathbf{n}(z) \in N_2 Z_2 \mathbf{d}(\det z) \mathbf{t}(z J_1) \mathbf{K}(\mathfrak{p}).$$

Remark 4.5. Let $a, b, c \in F$. Since

$$(4.4) \quad \mathbf{T}(a + \mathfrak{p}^{-1}, b + \mathfrak{o}_F, c + \mathfrak{o}_F) = \mathbf{T}(a, b, c),$$

if $\mathbf{v}(b) \leq 1$, then we can find a triplet $x, y, w \in F$ which satisfies the conditions above and such that $\mathbf{T}(a, b, c) = \mathbf{T}(x, y, w)$.

Proof. Note that

$$z^{-1} = (\det z)^{-1} \begin{pmatrix} w & -y \\ -y & x \end{pmatrix} \in \begin{pmatrix} \mathfrak{p} & \mathfrak{p} \\ \mathfrak{p} & \mathfrak{o}_F \end{pmatrix}$$

by assumption. Now the lemma follows from (3.6). \square

Lemma 4.6. *If $\mathbf{v}(y) \leq 1$, then the value $\mathbf{T}(x, y, w)$ depends only on $\mathbf{v}(x)$, $\mathbf{v}(y)$ and $\mathbf{v}(w)$. We may therefore write*

$$\mathbf{T}(x, y, w) = \mathbf{T}_{\mathbf{v}(y)}(\mathbf{v}(x), \mathbf{v}(w)).$$

(1) *If $i \geq 1$, then*

$$\mathbf{T}_0(i, j) = \epsilon \gamma^{-i-j} q^{-3(i+j)/2} \delta^{j-i+1} T'(0).$$

(2) *If $i \geq 2$ and $j \geq 1$, then*

$$\mathbf{T}_1(i, j) = \epsilon \gamma^{-i-j} q^{-3(i+j)/2} \delta^{j-i+1} T'(0).$$

(3) *If $i \geq 2$, then*

$$\mathbf{T}_1(i, 0) = \gamma^{-i} q^{-3i/2} \delta^{2-i} T'(1), \quad \mathbf{T}_1(1, 0) = \gamma^{-2} q^{-3} T'(0).$$

(4) *If $j \geq 1$, then*

$$\mathbf{T}_1(1, j) = \epsilon \gamma^{-j-1} q^{-3(j+1)/2} \delta^j T'(1).$$

Proof. In view of Remark 4.5 we may assume that x, y, w satisfy the assumptions of Lemma 4.4. Then

$$\begin{aligned} \mathbf{T}(x, y, w) &= \sigma(\det z) |\det z|^{-3/2} \mathbf{T}'(z J_1) \\ &= \epsilon \sigma(\det z) |\det z|^{-3/2} \mathbf{T}'(z \mathrm{diag}[\varpi, 1]) \end{aligned}$$

by (4.3). If $y = 0$ or if $\mathbf{v}(x) \geq 2$ and $\mathbf{v}(w) \geq 1$, then since

$$z \mathrm{diag}[\varpi, 1] = \begin{pmatrix} \varpi x & y \\ \varpi y & w \end{pmatrix} = \mathrm{diag}[\varpi x, w] \begin{pmatrix} 1 & \frac{y}{\varpi x} \\ \frac{\varpi y}{w} & 1 \end{pmatrix} \in \mathrm{diag}[\varpi x, w] \mathfrak{J},$$

we get

$$\mathbf{T}(x, y, w) = \epsilon \sigma(\det z) |\det z|^{-3/2} \Lambda_0(\varpi x w^{-1}) \mathbf{T}'(\mathbf{1}_2).$$

If $\mathbf{v}(x) \leq 1$ or $\mathbf{v}(w) = 0$, then we can use (4.4) to verify that $\mathbf{T}(x, y, w)$ depends only on $\mathbf{v}(x)$, $\mathbf{v}(y)$ and $\mathbf{v}(w)$ by conjugating ϕ_σ by $\mathbf{t}(\text{diag}[u, 1])$, $\mathbf{t}(\text{diag}[1, v])$ and $\mathbf{d}(\lambda)$ with $u, v, \lambda \in \mathfrak{o}_F^\times$. We have proved (1) and (2).

Next we shall prove (3). Let $j = 0$. If $i \geq 3$, then since

$$\begin{pmatrix} \varpi^{1-i} & \varpi^{-1} \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} \varpi^{-1} - \varpi^{1-i} & \varpi^{1-i} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

we get

$$\mathbf{T}(\varpi^{-i}, \varpi^{-1}, 1) = \frac{\epsilon \delta^{1-i}}{\gamma^i q^{3i/2}} \mathbf{T}' \left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) = \frac{\delta^{1-i}}{\gamma^i q^{3i/2}} \mathbf{T}' \left(\begin{pmatrix} \varpi & 1 \\ 0 & 1 \end{pmatrix} \right).$$

We can easily prove the case $i = 1$ as $\mathbf{T}(\varpi^{-1}, \varpi^{-1}, 0) = \mathbf{T}(0, \varpi^{-1}, 0)$. Since

$$\begin{pmatrix} \varpi^{-2} & \varpi^{-1} \\ \varpi^{-1} & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} \varpi^{-1} & \varpi^{-2} \\ 0 & \varpi^{-1} \end{pmatrix},$$

we get $\mathbf{T}(\varpi^{-2}, \varpi^{-1}, 0) = \gamma^{-2} q^{-3} \mathbf{T}'(\mathbf{n}(\varpi^{-1}))$.

If $j \geq 2$, then we can prove (4) by observing that

$$\begin{pmatrix} 1 & \varpi^{-1} \\ 1 & \varpi^{-j} \end{pmatrix} = \begin{pmatrix} 1 & \varpi^{-1} \\ 0 & \varpi^{-j} \end{pmatrix} \begin{pmatrix} 1 - \varpi^{j-1} & 0 \\ \varpi^j & 1 \end{pmatrix}.$$

From the computation

$$\begin{pmatrix} 0 & \varpi^{-1} \\ 1 & \varpi^{-1} \end{pmatrix} = \begin{pmatrix} 1 & \varpi^{-1} \\ 0 & \varpi^{-1} \end{pmatrix} \begin{pmatrix} -1 & 0 \\ \varpi & 1 \end{pmatrix}$$

we can deduce the remaining case $i = j = 1$. □

Now we are led to

$$\begin{aligned} \mathbf{B}_{S', \Lambda}^\sigma(\phi_\sigma) &= \int_{\mathfrak{p}^{-1}} \int_{F^2} \mathbf{T}_{\mathbf{v}(y)}(\mathbf{v}(x), \mathbf{v}(w)) \psi(y) \, dx dw dy \\ &= \sum_{i=1}^{\infty} \sum_{j=0}^{\infty} q^{i+j} (1 - q^{-1})^{\min\{1, i-1\} + \min\{1, j\}} (\mathbf{T}_0(i, j) - \mathbf{T}_1(i, j)). \end{aligned}$$

If $i \geq 2$ and $j \geq 1$, then $\mathbf{T}_0(i, j) = \mathbf{T}_1(i, j)$ by Lemma 4.6(1), (2). Hence

$$\begin{aligned} \mathbf{B}_{S', \Lambda}^\sigma(\phi_\sigma) &= q(\mathbf{T}_0(1, 0) - \mathbf{T}_1(1, 0)) \\ &\quad + (1 - q^{-1}) \sum_{i=2}^{\infty} q^i (\mathbf{T}_0(i, 0) - \mathbf{T}_1(i, 0)) \\ &\quad + (1 - q^{-1}) \sum_{j=1}^{\infty} q^{j+1} (\mathbf{T}_0(1, j) - \mathbf{T}_1(1, j)) \\ &= q(\mathbf{T}_0(1, 0) - \mathbf{T}_1(1, 0)) + (1 - q^{-1})(I_0 + J_0) + I_1 + J_1, \end{aligned}$$

where

$$I_0 = \sum_{i=2}^{\infty} q^i \mathbf{T}_0(i, 0), \quad I_1 = -(1 - q^{-1}) \sum_{i=2}^{\infty} q^i \mathbf{T}_1(i, 0),$$

$$J_0 = \sum_{j=1}^{\infty} q^{j+1} \mathbf{T}_0(1, j), \quad J_1 = -(1 - q^{-1}) \sum_{j=1}^{\infty} q^{j+1} \mathbf{T}_1(1, j).$$

Lemma 4.6(1) gives

$$I_0 = \sum_{i=2}^{\infty} q^i \epsilon \gamma^{-i} q^{-3i/2} \delta^{-i+1} T'(0) = \frac{\epsilon \gamma^{-2} q^{-1} \delta^{-1}}{1 - \gamma^{-1} \delta^{-1} q^{-1/2}} T'(0),$$

$$J_0 = \sum_{j=1}^{\infty} q^{j+1} \epsilon \gamma^{-1-j} q^{-3(1+j)/2} \delta^j T'(0) = \frac{\epsilon \gamma^{-2} q^{-1} \delta}{1 - \gamma^{-1} \delta q^{-1/2}} T'(0).$$

Lemma 4.6(3) gives

$$I_1 = -(1 - q^{-1}) \sum_{i=2}^{\infty} q^i \gamma^{-i} q^{-3i/2} \delta^{2-i} T'(1) = -\frac{\gamma^{-2} q^{-1} (1 - q^{-1})}{1 - \gamma^{-1} \delta^{-1} q^{-1/2}} T'(1).$$

Lemma 4.6(4) gives

$$J_1 = -(1 - q^{-1}) \sum_{j=1}^{\infty} q^{j+1} \epsilon \frac{\delta^j T'(1)}{\gamma^{j+1} q^{3(j+1)/2}} = -\frac{\epsilon \gamma^{-2} q^{-1} \delta (1 - q^{-1})}{1 - \gamma^{-1} \delta q^{-1/2}} T'(1).$$

Proposition 4.7. *If $m \geq 1$, then*

$$T'(m) = (1 + \epsilon \delta) \frac{(-\epsilon \delta^{-1})^m}{q^m (1 - q^{-1})} T'(0), \quad T'(0) = L(1, \epsilon \Lambda_0^{-1}).$$

Proof. Since π_0 has no $\mathrm{GL}_2(\mathfrak{o}_F)$ -invariant vector,

$$\sum_{x \in \mathfrak{o}_F/\mathfrak{p}} \mathbf{T}' \left(A \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \right) = -\mathbf{T}'(AJ_1) = -\epsilon \mathbf{T}'(\mathrm{Adiag}[\varpi, 1])$$

by (4.3). Observe that

$$\mathbf{n}(y) \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} = \begin{pmatrix} 1 + yx & y \\ x & 1 \end{pmatrix} = \begin{pmatrix} -x^{-1} & 1 + yx \\ 0 & x \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & x^{-1} \\ 0 & 1 \end{pmatrix}$$

for $x \in \mathfrak{o}_F^\times$. If $m \geq 2$, then we get

$$\mathbf{T}'(\mathbf{n}(\varpi^{1-m})) + \epsilon(q-1) \mathbf{T}'(\mathbf{n}(\varpi^{1-m}) \mathrm{diag}[\varpi, 1]) = -\epsilon \mathbf{T}'(\mathbf{n}(\varpi^{1-m}) \mathrm{diag}[\varpi, 1]),$$

letting $A = \mathbf{n}(y)$ and $y = \varpi^{1-m}$. It follows that

$$T'(m) = -q^{-1} \epsilon \delta^{-1} T'(m-1) = \dots = (-q^{-1} \epsilon \delta^{-1})^{m-1} T'(1).$$

Letting $y = 0$, we get

$$\mathbf{T}' \left(\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \right) = \epsilon \mathbf{T}'(\mathbf{n}(-x) \mathrm{diag}[\varpi, 1]) = \epsilon \delta \mathbf{T}'(\mathbf{n}(\varpi^{-1})) = \epsilon \delta T'(1)$$

for every $x \in \mathfrak{o}_F^\times$. We obtain

$$T'(1) = -(1 + \epsilon \delta^{-1})(q-1)^{-1} T'(0).$$

Since $\mathcal{W}_{f_{\pi_0}}(\mathrm{diag}[a, 1]) = \varepsilon(a) |a|_F \mathbf{1}_{\mathfrak{o}_F}(a)$, one can easily compute $T'(0)$. \square

Proposition 4.8.

$$\mathbf{B}_{S',\Lambda}^\sigma(\phi_\sigma) = \epsilon\gamma^{-1}q^{-1/2} \frac{L\left(\frac{1}{2}, \sigma_K^{-1}\Lambda\right)}{L\left(\frac{3}{2}, \epsilon\sigma\right) L(1, \sigma^{-2})} L(1, \epsilon\Lambda_0^{-1}).$$

Proof. Proposition 4.7 gives

$$\begin{aligned} I_1 &= -\frac{\gamma^{-2}q^{-1}(1-q^{-1})}{1-\gamma^{-1}\delta^{-1}q^{-1/2}}(1+\epsilon\delta)\frac{-\epsilon\delta^{-1}}{q(1-q^{-1})}T'(0) = \frac{\gamma^{-2}q^{-2}(1+\epsilon\delta^{-1})}{1-\gamma^{-1}\delta^{-1}q^{-1/2}}T'(0), \\ J_1 &= -\frac{\epsilon\gamma^{-2}q^{-1}\delta(1-q^{-1})}{1-\gamma^{-1}\delta q^{-1/2}}(1+\epsilon\delta)\frac{-\epsilon\delta^{-1}}{q(1-q^{-1})}T'(0) = \frac{\gamma^{-2}q^{-2}(1+\epsilon\delta)}{1-\gamma^{-1}\delta q^{-1/2}}T'(0). \end{aligned}$$

Now we have

$$\begin{aligned} I &:= \frac{(1-q^{-1})I_0 + I_1}{T'(0)} = (1-q^{-1})\frac{\epsilon\gamma^{-2}q^{-1}\delta^{-1}}{1-\gamma^{-1}\delta^{-1}q^{-1/2}} + \frac{\gamma^{-2}q^{-2}(1+\epsilon\delta^{-1})}{1-\gamma^{-1}\delta^{-1}q^{-1/2}} \\ &= \frac{\epsilon\gamma^{-2}q^{-1}\delta^{-1} + \gamma^{-2}q^{-2}}{1-\gamma^{-1}\delta^{-1}q^{-1/2}}, \\ J &:= \frac{(1-q^{-1})J_0 + J_1}{T'(0)} = \frac{\epsilon\gamma^{-2}q^{-1}\delta + \gamma^{-2}q^{-2}}{1-\gamma^{-1}\delta q^{-1/2}}. \end{aligned}$$

By Lemma 4.6(1), (3) we have

$$\begin{aligned} q\frac{\mathbf{T}_0(1,0) - \mathbf{T}_1(1,0)}{T'(0)} + J &= \epsilon\gamma^{-1}q^{-1/2} - \gamma^{-2}q^{-2} + \frac{\epsilon\gamma^{-2}q^{-1}\delta + \gamma^{-2}q^{-2}}{1-\gamma^{-1}\delta q^{-1/2}} \\ &= \frac{\epsilon\gamma^{-1}q^{-1/2} + \delta\gamma^{-3}q^{-5/2}}{1-\gamma^{-1}\delta q^{-1/2}}. \end{aligned}$$

We conclude that

$$\begin{aligned} \frac{\mathbf{B}_{S',\Lambda}^\sigma(\phi_\sigma)}{T'(0)} &= \frac{\epsilon\gamma^{-2}q^{-1}\delta^{-1} + \gamma^{-2}q^{-2}}{1-\gamma^{-1}\delta^{-1}q^{-1/2}} + \frac{\epsilon\gamma^{-1}q^{-1/2} + \delta\gamma^{-3}q^{-5/2}}{1-\gamma^{-1}\delta q^{-1/2}} \\ &= \epsilon\gamma^{-1}q^{-1/2} \frac{(1+\epsilon\gamma^{-1}q^{-3/2})(1-\gamma^{-2}q^{-1})}{(1-\gamma^{-1}\delta^{-1}q^{-1/2})(1-\gamma^{-1}\delta q^{-1/2})}. \end{aligned}$$

The proof is complete by Proposition 4.7. \square

We use the Iwasawa decomposition to define $\phi^{\text{pa}} \in I(\text{Id}, 1)$ by

$$\phi^{\text{pa}}(\mathbf{d}(\lambda)\mathbf{t}(A)\mathbf{n}(z)k) = |\lambda|^{-3/2},$$

where $\lambda \in F^\times$, $A \in \text{GL}_2(F)$, $z \in \text{Sym}_2(F)$ and $k \in \text{K}(\mathfrak{p})$. The elements $\phi_w \in I(I_1(\omega_F^{-1/2}), 1)$ are defined in Section 7 below. The pairing $\mathbf{q} : I(\text{Id}, 1) \times I(\text{Id}, 1) \rightarrow \mathbb{C}$ will be defined in (5.2).

Lemma 4.9. *We have*

$$\begin{aligned} \phi^{\text{pa}} &= \phi_{\mathbf{1}_4} + \phi_{s_1} + \phi_{s_2} + \phi_{s_1s_2} + q^{-3/2}(\phi_{s_2s_1} + \phi_{s_1s_2s_1} + \phi^\dagger + \phi_{s_2s_1s_2}), \\ \mathbf{q}(\phi^{\text{pa}}, \phi^{\text{pa}}) &= q^{-2}\zeta(1)^2\zeta(2)^{-2}. \end{aligned}$$

Proof. Since $s_2 \in K(\mathfrak{p})$, it is clear that

$$\begin{aligned}\phi^{\text{pa}}(\mathbf{1}_4) &= \phi^{\text{pa}}(s_1) = \phi^{\text{pa}}(s_2) = \phi^{\text{pa}}(s_1 s_2) = 1, \\ \phi^{\text{pa}}(s_2 s_1) &= \phi^{\text{pa}}(s_1 s_2 s_1) = \phi^{\text{pa}}(s_2 s_1 s_2) = \phi^{\text{pa}}(s_1 s_2 s_1 s_2).\end{aligned}$$

Since $\mathbf{m}(\text{diag}[\varpi^{-1}, 1])s_1 s_2 s_1 s_2 \in K(\mathfrak{p})$, we get $\phi^{\text{pa}}(s_1 s_2 s_1 s_2) = q^{-3/2}$. The Iwasawa decomposition relative to $\text{GSp}_4(\mathfrak{o}_F)$ gives

$$\mathbf{q}(\phi^{\text{pa}}, \phi^{\text{pa}}) = \int_{\text{GSp}_4(\mathfrak{o}_F)} \phi^{\text{pa}}(k)^2 dk = \sum_{w \in W} \phi^{\text{pa}}(w)^2 \text{vol}(\mathbb{I}w\mathbb{I}).$$

Clearly, $\text{vol}(\mathbb{I}w\mathbb{I}) = q_w \text{vol}(\mathbb{I})$ (see Definition 2.1). Since $\text{vol}(\mathbb{I}s_1 s_2 s_1 s_2 \mathbb{I}) + \text{vol}(\mathbb{I}s_2 s_1 s_2 \mathbb{I}) = 1$ for our choice of the measure, $\text{vol}(\mathbb{I}) = q^{-4}(1+q^{-1})^{-1}$ and

$$\begin{aligned}\mathbf{q}(\phi^{\text{pa}}, \phi^{\text{pa}}) &= (1+q^{-1})^{-1} \sum_{w \in W} \phi^{\text{pa}}(w)^2 q^{-4} q_w \\ &= (1+q^{-1})^{-1} \{q^{-4} + 2q^{-3} + q^{-2} + q^{-3}(q^{-2} + 2q^{-1} + 1)\},\end{aligned}$$

which proves the second identity. \square

Proposition 4.10. *Let $\sigma, \Lambda_0 \in \Omega^1(F^\times)^\circ$. Fix a quadratic unramified character $\varepsilon : F^\times \rightarrow \{\pm 1\}$. Put $\pi = I(\text{St} \otimes \varepsilon, \sigma)$. Then*

$$\mathbb{B}_S^\Lambda(\pi(\mathbf{m}(\varsigma^{-1}))) = -\varepsilon q(1+q^{-2})\Lambda_0(\varpi)^{-1} \frac{|\mathfrak{d}_K|_K^{1/2} \zeta(2)\zeta(4)L(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \Lambda)}{|\mathfrak{d}_F|^{1/2} L(1, \tau_{K/F})L(1, \pi, \text{ad})}.$$

Proof. We know that

$$(4.5) \quad \begin{aligned}L(s, \text{Spn}(\pi)_K \otimes \Lambda) &= L(s, \sigma_K^{-1} \Lambda) L(s, \sigma_K \Lambda) L(s, \omega_K^{1/2} \varepsilon_K \Lambda), \\ L(s, \pi, \text{ad}) &= \zeta(s)\zeta(s+1)L(s, \sigma^{-2})L(s, \sigma^2)L(s, \omega_F^{1/2} \varepsilon \sigma)L(s, \omega_F^{1/2} \varepsilon \sigma^{-1})\end{aligned}$$

(see [31, Table A.8] and [3, (13)]). We also remark that

$$\varepsilon(1/2, \text{St} \otimes \varepsilon \Lambda_0^{-1}) = \Lambda_0(\varpi)^{-1}(-\varepsilon),$$

(cf. [8, Lemma 3.1, Proposition 3.8]). By (4.1) and Remark 3.5

$$b_{\mathcal{W}}^\sharp(\phi_\sigma, \phi_{\sigma^{-1}}) \mathbb{B}_S^\Lambda(\pi(\mathbf{m}(\varsigma^{-1}))) = \gamma \left(\frac{1}{2}, \text{St} \otimes \varepsilon \Lambda_0^{-1}, \psi \right) \mathbf{B}_{S', \Lambda}^\sigma(\phi_\sigma) \mathbf{B}_{S', \Lambda}^{\sigma^{-1}}(\phi_{\sigma^{-1}}).$$

By Proposition 4.8 the right hand side is equal to

$$\begin{aligned}& \frac{L(1, \varepsilon \Lambda_0)}{L(1, \varepsilon \Lambda_0^{-1})} (-\varepsilon) \Lambda_0(\varpi)^{-1} \frac{L(\frac{1}{2}, \sigma_K^{-1} \Lambda) L(\frac{1}{2}, \sigma_K \Lambda) L(1, \varepsilon \Lambda_0^{-1})^2}{qL(\frac{3}{2}, \varepsilon \sigma) L(1, \sigma^{-2}) L(\frac{3}{2}, \varepsilon \sigma^{-1}) L(1, \sigma^2)} \\ &= (-\varepsilon) \Lambda_0(\varpi)^{-1} \zeta(1)\zeta(2) \frac{L(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \Lambda)}{qL(1, \pi, \text{ad})}.\end{aligned}$$

Since $\mathcal{W}_{\phi_\sigma}(\mathbf{t}(\text{diag}[a, 1])) = \varepsilon(a)|a|_F \mathbb{1}_{\mathfrak{o}_F}(a)$ for $a \in F^\times$, we have

$$b_{\mathcal{W}}(\pi(g)\phi_\sigma, \pi^\vee(g)\phi_{\sigma^{-1}}) = \zeta(2)\phi^{\text{pa}}(g)^2$$

and hence $b_{\mathcal{W}}^\sharp(\phi_\sigma, \phi_{\sigma^{-1}}) = \zeta(2)\mathbf{q}(\phi^{\text{pa}}, \phi^{\text{pa}}) = q^{-2}\zeta(1)^2\zeta(2)^{-1}$. \square

5. EXPLICIT CALCULATIONS OF BESSEL INTEGRALS II: DEGENERATE VECTORS

5.1. Degenerate principal series. Let D be a quaternion algebra over a local field F of characteristic zero. We retain the notation in Section 2. Fix a quadratic character ε of F^\times . For $\sigma \in \Omega(F^\times)$ we consider the normalized induced representation $\pi = I(\varepsilon \circ \mathbf{N}_F^D, \sigma)$, which is realized on the space of smooth functions $\phi : \mathrm{GU}_2^D(F) \rightarrow \mathbb{C}$ satisfying

$$\phi(\mathbf{d}(\lambda)\mathbf{t}(A)\mathbf{n}(z)g) = \sigma(\lambda)|\lambda|^{-3/2}\varepsilon(\mathbf{N}_F^D(A))\phi(g)$$

for $A \in D^\times$, $z \in D_-$, $\lambda \in F^\times$ and $g \in \mathrm{GU}_2^D(F)$.

In the \mathfrak{p} -adic case we fix a maximal compact subring \mathfrak{o}_D of D and set $\mathcal{K} = \mathrm{GU}_2^D(F) \cap \mathrm{GL}_2(\mathfrak{o}_D)$. In the archimedean case we fix a maximal compact subgroup \mathcal{K} of $\mathrm{GU}_2^D(F)$. For $g \in \mathrm{GU}_2^D(F)$ the quantity $|a(g)|$ is defined by setting $|a(g)| = |\lambda|_F^{-1}$, where we write $g = pk$ with $p = \mathbf{d}(\lambda)\mathbf{t}(A)\mathbf{n}(z) \in \mathcal{P}$ and $k \in \mathcal{K}$. For $\phi \in \pi$ and $s \in \mathbb{C}$ we define $\phi^{(s)} \in I(\varepsilon \circ \mathbf{N}_F^D, \sigma\omega_F^s)$ by $\phi^{(s)}(g) = \phi(g)|a(g)|^{-s}$. We define the intertwining operator

$$M(\varepsilon \circ \mathbf{N}_F^D, \sigma) : I(\varepsilon \circ \mathbf{N}_F^D, \sigma) \rightarrow I(\varepsilon \circ \mathbf{N}_F^D, \sigma^{-1})$$

by

$$[M(\varepsilon \circ \mathbf{N}_F^D, \sigma)\phi](g) = \lim_{s \rightarrow 0} \int_{D_-} \phi^{(s)} \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z)g \right) dz.$$

The integral converges for $\Re s + \Re \sigma < -\frac{3}{2}$ and admits meromorphic continuation to whole s -plane.

5.2. Degenerate Whittaker model. Fix $S \in D_-$. Put

$$d_0 = -\mathbf{N}_F^D(2S), \quad K = F(\sqrt{d_0}).$$

We identify K with the subalgebra $F + FS$ of D . Put

$$R_S = \{\mathbf{t}(t)\mathbf{n}(z) \mid z \in D_-, t \in K^\times\}.$$

We define an additive character ψ_S on D_- by $\psi_S(z) = \psi(\mathrm{Tr}_F^D(Sz))$ and associate a character Λ^S of R_S by $\Lambda^S(\mathbf{t}(t)\mathbf{n}(z)) = \Lambda(t)\psi_S(z)$ to a character Λ of $F^\times \backslash K^\times$. The integral

$$\mathbf{B}_S^{\sigma\omega_F^s}(\phi^{(s)}) = \int_{D_-} \phi^{(s)} \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z) \right) \overline{\psi_S(z)} dz$$

is absolutely convergent for $\Re s + \Re \sigma < -\frac{3}{2}$ and makes sense for all s by an entire analytic continuation. In the nonarchimedean case the integral stabilizes. The reader who are interested in the archimedean case can consult [37]. One can easily see that

$$(5.1) \quad \mathbf{B}_S^\sigma \circ \pi(\mathbf{t}(t)) = \varepsilon(\mathbf{N}_F^K(t))\mathbf{B}_S^\sigma$$

for $t \in K^\times$. Thus \mathbf{B}_S^σ is a Bessel functional on π relative to S and ε_K .

We introduce the $\mathrm{GU}_2^D(F)$ -invariant pairing

$$\mathbf{q} : I(\varepsilon \circ \mathbf{N}_F^D, \sigma) \times I(\varepsilon \circ \mathbf{N}_F^D, \sigma^{-1}) \rightarrow \mathbb{C}$$

by

$$(5.2) \quad \begin{aligned} \mathbf{q}(\phi_1, \phi_2) &= \int_{\mathcal{P} \backslash \mathrm{GU}_2^D(F)} \phi_1(g) \phi_2(g) \, dg \\ &= \int_{D_-} \phi_1 \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z) \right) \phi_2 \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z) \right) \, dz \end{aligned}$$

for $\phi_1 \in I(\varepsilon \circ \mathrm{N}_F^D, \sigma)$ and $\phi_2 \in I(\varepsilon \circ \mathrm{N}_F^D, \sigma^{-1})$. Define the Bessel integral by

$$\begin{aligned} J_S(\phi_1, \phi_2) &= \int_{D_-} \mathbf{q}(\pi(\mathbf{n}(z))\phi_1, \pi(\mathbf{d}(-1))\phi_2) \overline{\psi_S(z)} \, dz \\ &:= \lim_{s \rightarrow 0} \sigma(-1) \int_{D_-}^2 \phi_1^{(s)} \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z - z') \right) \phi_2^{(s)} \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z') \right) \, dz' \overline{\psi_S(z)} \, dz. \end{aligned}$$

The double integral absolutely converges for $\Re s \ll 0$ and can be continued as an entire function to the whole complex plane. We have the factorization

$$(5.3) \quad J_S(\phi, \phi') = \mathbf{B}_S^\sigma(\phi) \mathbf{B}_{-S}^{\sigma^{-1}}(\pi^\vee(\mathbf{d}(-1))\phi').$$

It follows that

$$J_S \in \mathrm{Hom}_{R_S \times R_S}(I(\varepsilon \circ \mathrm{N}_F^D, \sigma) \boxtimes I(\varepsilon \circ \mathrm{N}_F^D, \sigma^{-1}), \varepsilon_K^S \boxtimes \varepsilon_K^S).$$

If $I(\varepsilon \circ \mathrm{N}_F^D, \sigma)$ has a new vector ϕ_σ , then we have a functional equation

$$M(\varepsilon \circ \mathrm{N}_F^D, \sigma) \phi_\sigma = c(\varepsilon, \sigma) \phi_{\sigma^{-1}}$$

with factor $c(\varepsilon, \sigma)$ of proportionality. We set

$$\mathbf{B}_S(\phi, \phi') = J_S(\phi, M(\varepsilon \circ \mathrm{N}_F^D, \sigma) \phi').$$

Then

$$(5.4) \quad \frac{\mathbf{B}_S(\phi_\sigma, \phi_\sigma)}{\mathbf{q}(\phi_\sigma, M(\varepsilon \circ \mathrm{N}_F^D, \sigma) \phi_\sigma)} = \frac{\mathbf{B}_S^\sigma(\phi_\sigma) \mathbf{B}_{-S}^{\sigma^{-1}}(\pi^\vee(\mathbf{d}(-1))\phi_{\sigma^{-1}})}{\mathbf{q}(\phi_\sigma, \phi_{\sigma^{-1}})}.$$

5.3. The quaternion case. Let D be a quaternion division algebra over a \mathfrak{p} -adic field F of characteristic 0. We denote by \mathfrak{o}_D the maximal compact subring of D and by \mathfrak{P} the maximal proper two-sided ideal of \mathfrak{o}_D . Put $\mathfrak{o}_{\bar{D}} = \mathfrak{o}_D \cap D_-$. Define a maximal compact subgroup $\mathrm{K}(\mathfrak{P})$ of $\mathrm{GU}_2^D(F)$ by

$$(5.5) \quad \mathrm{K}(\mathfrak{P}) = \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{GU}_2^D(F) \mid \alpha, \delta \in \mathfrak{o}_D, \beta \in \mathfrak{P}^{-1}, \gamma \in \mathfrak{P} \right\}.$$

Then $\mathrm{GU}_2^D(F) = \mathcal{PK}(\mathfrak{P})$. Let $\sigma = \omega_F^s$ be an unramified character of F^\times and $\varepsilon = \omega_F^t$ an unramified quadratic character of F^\times . Put $\pi = I(\varepsilon \circ \mathrm{N}_F^D, \sigma)$. We define its L -factors as those of $I(\varepsilon \circ \mathrm{St}, \sigma)$.

Remark 5.1. The representations $I(\varepsilon \circ \mathrm{N}_F^D, \sigma)$ are called type IIa^G in [39]. The subgroup $\mathrm{K}(\mathfrak{P})$ is considered in [20, 29]. Appendix of [29] shows that representations of type IIa^G (with unramified ε and σ) are only the tempered representations having a non-zero $\mathrm{K}(\mathfrak{P})$ -fixed vector, which is up to scalar.

It follows from [3] that

$$\frac{L(\frac{1}{2}, \mathrm{Spn}(\pi)_K \otimes \varepsilon_K)}{L(1, \pi, \mathrm{ad})} = \frac{L(\frac{1}{2}, (\sigma^{-1}\varepsilon)_K)L(\frac{1}{2}, (\sigma\varepsilon)_K)L(1, \tau_{K/F})}{\zeta(2)L(1, \sigma^{-2})L(1, \sigma^2)L(\frac{3}{2}, \varepsilon\sigma^{-1})L(\frac{3}{2}, \varepsilon\sigma)}.$$

For $g \in \mathrm{GU}_2^D(F)$ the quantity $|\alpha(g)|$ is defined via

$$|\alpha(\mathbf{d}(\lambda)\mathbf{n}(A)\mathbf{n}(z)k)| = |\mathrm{N}_F^D(A)|_F,$$

where $\lambda \in F^\times$, $A \in D^\times$, $z \in D_-$ and $k \in \mathbf{K}(\mathfrak{P})$. We define $\phi_\sigma \in \pi$ by

$$\phi_\sigma(g) = |\lambda(g)|_F^{(2s-3)/2} |\alpha(g)|^{(-2s+2t+3)/2}.$$

Proposition 5.2 (Hirai [20]). *Let $S \in \mathfrak{o}_D^-$ and $K = F + FS$. Take the Haar measure dz on D_- which gives $D_- \cap \mathfrak{P}$ the volume 1.*

(1) *If $\varpi^{-1}S \notin \mathfrak{o}_D^-$, then*

$$\mathbf{B}_S^\sigma(\phi_\sigma) = \frac{q^{3/2}(\varepsilon\sigma)(\varpi)^{-1}L(\frac{1}{2}, (\varepsilon\sigma^{-1})_K)}{L(1, \sigma^{-2})L(\frac{3}{2}, \varepsilon\sigma^{-1})}.$$

(2) $\mathbf{q}(\phi_\sigma, \phi_{\sigma^{-1}}) = \zeta(2)\zeta(4)^{-1}$.

Proof. Put $\mathbf{K}(\mathfrak{P})' = \mathbf{d}(\varpi)^{-1}\mathbf{K}(\mathfrak{P})\mathbf{d}(\varpi)$. Observe that

$$\mathbf{K}(\mathfrak{P})' = \left\{ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{GU}_2^D(F) \mid \alpha, \delta \in \mathfrak{o}_D, \beta \in \mathfrak{P}, \gamma \in \mathfrak{P}^{-1} \right\}.$$

Define the function $A_{\mathfrak{p}} : \mathrm{GU}_2^D(F) \rightarrow \mathbb{C}$ by $A_{\mathfrak{p}}(g) = |\alpha(g\mathbf{d}(\varpi)^{-1})|$. Set

$$\alpha_{\mathfrak{p}}(\eta, x) = \int_{D_-} A_{\mathfrak{p}} \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z) \right)^{(2x+3)/2} \overline{\psi_\eta(z)} dz$$

for $\eta \in \varpi^{-1}\mathfrak{o}_D^-$ and $x \in \mathbb{C}$. Put

$$\phi'_\sigma = \pi(\mathbf{d}(\varpi)^{-1})\phi_\sigma, \quad S' = \varpi^{-1}S.$$

By a change of variables we have

$$\begin{aligned} \mathbf{B}_S^\sigma(\phi_\sigma) &= \int_{D_-} \phi'_\sigma \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \mathbf{n}(z)\mathbf{d}(\varpi) \right) \overline{\psi_S(z)} dz \\ &= q^{(2s-3)/2} \cdot q^3 \mathbf{B}_{S'}^\sigma(\phi'_\sigma) \\ &= q^{(2s+3)/2} \cdot q^{(2s-3)/2} \alpha_{\mathfrak{p}}(S', t-s). \end{aligned}$$

Theorem 2.3 of [20] explicitly computes $\alpha_{\mathfrak{p}}(\eta, s)$. By the assumption on S'

$$\alpha_{\mathfrak{p}}(S', s) = q^{(2s+3)/2} \frac{\zeta_K(s + \frac{1}{2})}{\zeta(2s+1)\zeta(s + \frac{3}{2})}.$$

We remind the reader that the measure in [20] gives $\mathfrak{P} \cap D_-$ volume 1. The second part follows from the obvious equality $\mathbf{q}(\phi_\sigma, \phi_{\sigma^{-1}}) = q^{-3}\alpha_{\mathfrak{p}}(0, \frac{3}{2})$. \square

Now we have the following conclusion by (5.4) and Proposition 5.2.

Corollary 5.3.

$$\frac{\mathbf{B}_S(\phi_\sigma, \phi_\sigma)}{\mathbf{q}(\phi_\sigma, M(\varepsilon \circ N_F^D, \sigma)\phi_\sigma)} = q^3(1 - q^{-2}) \frac{\zeta(4)L(\frac{1}{2}, \mathrm{Spn}(\pi)_K \otimes \varepsilon_K)}{\zeta(2)L(1, \tau_{K/F})L(1, \pi, \mathrm{ad})}.$$

5.4. The archimedean case. We discuss the case in which $F = \mathbb{R}$ and $K = \mathbb{C}$. Put $\zeta_F(s) = \pi^{-s/2}\Gamma(\frac{s}{2})$. We define the character of \mathbb{C} by $\mathbf{e}(z) = e^{2\pi\sqrt{-1}z}$ for $z \in \mathbb{C}$. Its restriction to F is denoted by ψ^F . The measure dx on \mathbb{R} is the Lebesgue measure and $d^\times x = \frac{dx}{|x|}$. Let dz be the standard measure on $\mathrm{Sym}_2(\mathbb{R})$ defined by viewing $\mathrm{Sym}_2(\mathbb{R})$ as \mathbb{R}^3 in an obvious fashion.

Denote by $\mathrm{Sym}_g(\mathbb{R})^+$ the set of positive definite symmetric matrices of rank g over \mathbb{R} . Let

$$\mathrm{GSp}_{2g}(\mathbb{R})^\circ = \{h \in \mathrm{GSp}_{2g} \mid \lambda(h) > 0\}$$

be the identity component of the real reductive group $\mathrm{GSp}_{2g}(\mathbb{R})$. We can define the action of the connected component $\mathrm{GSp}_{2g}(\mathbb{R})^\circ$ on the space

$$\mathfrak{H}_g = \{Z \in \mathrm{Sym}_g(\mathbb{C}) \mid \Im Z \in \mathrm{Sym}_g(\mathbb{R})^+\}$$

and the automorphy factor on $\mathrm{GSp}_{2g}(\mathbb{R})^\circ \times \mathfrak{H}_g$ by

$$hZ = (AZ + B)(CZ + D)^{-1}, \quad j(h, Z) = (\det h)^{-1/2} \det(CZ + D)$$

for $Z \in \mathfrak{H}_g$ and $h = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{GSp}_{2g}(\mathbb{R})^\circ$ with matrices A, B, C, D of size g . Put $\mathbf{i}_g = \sqrt{-1}\mathbf{1}_g$. Define the maximal compact subgroup of $\mathrm{GSp}_{2g}(\mathbb{R})^\circ$ by

$$U_n = \{h \in \mathrm{GSp}_{2g}(\mathbb{R})^\circ \mid h(\mathbf{i}_g) = \mathbf{i}_g\}.$$

Definition 5.4. For each positive integer κ we denote the lowest weight representation of $\mathrm{GSp}_{2g}(\mathbb{R})^\circ$ with lowest U_g -type $k \mapsto j(k, \mathbf{i}_g)^{-\kappa}$ by $\mathfrak{D}_\kappa^{(g)}$ and the highest weight representation with highest U_g -type $k \mapsto j(k, \mathbf{i}_g)^\kappa$ by $\mathfrak{D}_{-\kappa}^{(g)}$. The direct sum $D_\kappa^{(g)} = \mathfrak{D}_\kappa^{(g)} \oplus \mathfrak{D}_{-\kappa}^{(g)}$ extends to an irreducible representation of $\mathrm{GSp}_{2g}(\mathbb{R})$.

Remark 5.5. If $S \in \mathrm{Sym}_2(\mathbb{R})$ is positive or negative definite, then Theorem 3.10 of [30] says that $D_\kappa^{(2)}$ admits a Bessel model relative to (Λ, S) if and only if Λ is trivial. This fact is compatible with our observations in the previous subsection.

To simplify notation, we set

$$J(m) = I(\mathrm{sgn}^m \circ \det, \omega_{\mathbb{R}}^{(3-2m)/2}), \quad N(m) = M(\mathrm{sgn}^m \circ \det, \omega_{\mathbb{R}}^{(3-2m)/2})$$

for $m \in \mathbb{Z}$. As is well-known, $\mathfrak{D}_{\pm\kappa}^{(2)}$ are subrepresentations of the degenerate principal series $J(\kappa)$ which is here viewed as a representation of $\mathrm{GSp}_4(\mathbb{R})^\circ$. The representation $\mathfrak{D}_\kappa^{(2)}$ (resp. $\mathfrak{D}_{-\kappa}^{(2)}$) is generated by the function $\phi_\kappa(h) = j(h, \mathbf{i}_2)^{-\kappa}$ (resp. $\phi_{-\kappa}(h) = \overline{j(h, \mathbf{i}_2)^{-\kappa}}$). These functions $\phi_{\pm\kappa}$ are extended

uniquely to elements of $J(\kappa)$. Since $\phi_\kappa(h\mathbf{d}(-1)) = \phi_{-\kappa}(h)$, we can view $D_\kappa^{(2)}$ as a subrepresentation of $J(\kappa)$.

Put $\varphi_\kappa = \phi_\kappa^{(2\kappa-3)} \in J(3-\kappa)$ and $\mathbf{B}_S^s = \mathbf{B}_S^{\omega_{\mathbb{R}}^s}$. Observe that

$$\begin{aligned} \mathbf{B}_S^{(2s+3-2\kappa)/2}(\phi_\kappa^{(s)}) &= \int_{\mathrm{Sym}_2(\mathbb{R})} \phi_\kappa^{(s)}(J_2\mathbf{n}(z))\mathbf{e}(-\mathrm{tr}(Sz)) \, dz \\ &= \int_{\mathrm{Sym}_2(\mathbb{R})} |\det(z + \mathbf{i}_2)|^s \det(z + \mathbf{i}_2)^{-\kappa} \mathbf{e}(-\mathrm{tr}(Sz)) \, dz \\ &= \xi\left(\mathbf{1}_2, S; \kappa - \frac{s}{2}, -\frac{s}{2}\right). \end{aligned}$$

The confluent hypergeometric function $\xi(Y, S; \alpha, \beta)$ is extensively studied in [34]. By Lemma 3.1 of [40] the operator

$$\frac{M(\mathrm{sgn}^\kappa \circ \det, \omega_{\mathbb{R}}^{-s})}{L(-s - \frac{1}{2}, \mathrm{sgn}^\kappa)\Gamma(-s)}$$

is entire. In particular, $M(\mathrm{sgn}^\kappa \circ \det, \omega_{\mathbb{R}}^{-s})$ is holomorphic at $s = \kappa - \frac{3}{2}$. Letting $S = 0$, $s = 2\kappa - 3 + t$ and $t \rightarrow 0$, we get

$$\begin{aligned} N(3-\kappa)\varphi_\kappa &= \lim_{t \rightarrow 0} \xi\left(\mathbf{1}_2, 0; \frac{3-t}{2}, \frac{3-t}{2} - \kappa\right) \phi_\kappa^{(-t)} \\ &= \lim_{t \rightarrow 0} (-1)^\kappa 2^{-1} (2\pi)^3 \frac{\Gamma_2(\frac{3}{2} - t - \kappa)}{\Gamma_2(\frac{3-t}{2})\Gamma_2(\frac{3-t}{2} - \kappa)} 2^{2\kappa+2t-3} \phi_\kappa \\ &= (-1)^\kappa 4^{-1} (2\pi)^3 \Gamma_2(3/2)^{-1} 2^{2\kappa-3} \phi_\kappa = 2^{-1} (-4)^\kappa \pi^2 \phi_\kappa \end{aligned}$$

by (1.31) of [34], where $\Gamma_2(s) = \sqrt{\pi}\Gamma(s)\Gamma(s - \frac{1}{2})$. We write $\mathcal{D}_\kappa^{(2)}$ for the subrepresentation of $J(3-\kappa)$ generated by φ_κ . It has the module $D_\kappa^{(2)}$ as a quotient. The quotient map $\mathcal{D}_\kappa^{(2)} \rightarrow D_\kappa^{(2)}$ is realized by the operator $N(3-\kappa)$. Since $J(3-\kappa)$ is multiplicity free even as a representation of U_2 , any $\mathrm{GSp}_4(\mathbb{R})$ -invariant pairing $\mathcal{D}_\kappa^{(2)} \times D_\kappa^{(2)} \rightarrow \mathbb{C}$ factors through the quotient map. We construct a $\mathrm{GSp}_4(\mathbb{R})$ -invariant pairing $\mathbf{r} : D_\kappa^{(2)} \times D_\kappa^{(2)} \rightarrow \mathbb{C}$ in the following way: for $\phi, \phi' \in \mathcal{D}_\kappa^{(2)}$ we set

$$\mathbf{r}(\phi, \phi') = \mathbf{q}(\varphi, \phi'),$$

where we take $\varphi \in \mathcal{D}_\kappa^{(2)}$ so that $N(3-\kappa)\varphi = \phi$.

Definition 5.6 (Bessel integrals for $D_\kappa^{(2)}$). We define

$$\mathbf{A}_S^\kappa(\phi, \phi') = \int_{\mathrm{Sym}_2(\mathbb{R})} \mathbf{r}(D_\kappa^{(2)}(\mathbf{n}(z))\phi, D_\kappa^{(2)}(\mathbf{d}(-1))\phi') \overline{\mathbf{e}(\mathrm{tr}(Sz))} \, dz$$

for $\kappa \in \mathbb{N}$, $S \in \mathrm{Sym}_2(\mathbb{R})^+$ and $\phi, \phi' \in D_\kappa^{(2)}$.

Remark 5.7. We make the integral above meaningful by analytic continuation (see the previous subsection). If $\kappa \geq 2$, then $D_\kappa^{(2)}$ is a discrete series and this integral converges absolutely by Proposition 3.15 of [26].

Proposition 5.8. *For every positive integer κ and $S \in \text{Sym}_2(\mathbb{R})^+$ we have*

$$\frac{\mathbf{A}_S^\kappa(\phi_\kappa, \phi_\kappa)}{\mathbf{r}(\phi_\kappa, D_\kappa^{(2)}(\mathbf{d}(-1))\phi_\kappa)} = 2^{4\kappa-2}(2\pi)^{2\kappa-1} \frac{(\det S)^{(2\kappa-3)/2}}{\Gamma(2\kappa-1)} e^{-4\pi\text{tr}(S)}.$$

Proof. In view of (5.3) we arrive at

$$\begin{aligned} \mathbf{A}_S^\kappa(\phi, \phi') &= \int_{\text{Sym}_2(\mathbb{R})} \mathbf{q}(D_\kappa^{(2)}(\mathbf{n}(z))\varphi, D_\kappa^{(2)}(\mathbf{d}(-1))\phi') \overline{\mathbf{e}(\text{tr}(Sz))} dz \\ &= \mathbf{B}_S^{(2\kappa-3)/2}(\varphi) \mathbf{B}_{-S}^{(3-2\kappa)/2}(D_\kappa^{(2)}(\mathbf{d}(-1))\phi'). \end{aligned}$$

Let $\phi = \phi' = \phi_\kappa$ and $\varphi = 2(-4)^{-\kappa}\pi^{-2}\varphi_\kappa$. Then

$$\begin{aligned} \mathbf{r}(\phi_\kappa, D_\kappa^{(2)}(\mathbf{d}(-1))\phi_\kappa) &= 2(-4)^{-\kappa}\pi^{-2} \mathbf{q}(\varphi_\kappa, D_\kappa^{(2)}(\mathbf{d}(-1))\phi_\kappa) \\ &= 2(-4)^{-\kappa}\pi^{-2} \int_{\text{Sym}_2(\mathbb{R})} \varphi_\kappa(J_2\mathbf{n}(z))\phi_{-\kappa}(J_2\mathbf{n}(z)) dz \\ &= 2(-4)^{-\kappa}\pi^{-2} \xi\left(\mathbf{1}_2, 0; \frac{3}{2}, \frac{3}{2}\right) = 2(-4)^{-\kappa}. \end{aligned}$$

From (4.34.K) and (4.35.K) of [34]

$$\begin{aligned} \mathbf{B}_S^{(3-2\kappa)/2}(\phi_\kappa) &= \xi(\mathbf{i}_2, S; \kappa, 0) = (-1)^\kappa 4\pi^{(4\kappa-1)/2} \frac{(4 \det S)^{(2\kappa-3)/2}}{\Gamma(\kappa)\Gamma(\kappa - \frac{1}{2})} e^{-2\pi\text{tr}(S)}, \\ \mathbf{B}_S^{(2\kappa-3)/2}(\varphi_\kappa) &= \xi\left(\mathbf{i}_2, S; \frac{3}{2}, \frac{3}{2} - \kappa\right) = (-4)^\kappa \pi^2 e^{-2\pi\text{tr}(S)}. \end{aligned}$$

We get $\mathbf{B}_S^{(2\kappa-3)/2}(\varphi) = 2e^{-2\pi\text{tr}(S)}$ and conclude that

$$\begin{aligned} \frac{\mathbf{A}_S^\kappa(\phi_\kappa, \phi_\kappa)}{\mathbf{r}(\phi_\kappa, D_\kappa^{(2)}(\mathbf{d}(-1))\phi_\kappa)} &= 2^{-1}(-4)^\kappa \mathbf{B}_S^{(2\kappa-3)/2}(\varphi) \mathbf{B}_{-S}^{(3-2\kappa)/2}(D_\kappa^{(2)}(\mathbf{d}(-1))\phi_\kappa) \\ &= 2^{-1}(-4)^\kappa 2e^{-2\pi\text{tr}(S)} \overline{\mathbf{B}_S^{(3-2\kappa)/2}(\phi_\kappa)}, \end{aligned}$$

which completes our proof. \square

6. BESSEL PERIODS ON PRINCIPAL SERIES REPRESENTATIONS

6.1. Tate's local zeta integral. Let us recall Tate's theory for local factors of quasi-characters of the multiplicative group of a local field L . Denote by $\mathcal{S}(L)$ the space of Bruhat-Schwartz functions on L . Fix a non-trivial additive character ψ_L of L . Tate's local zeta integral is defined by

$$\mathcal{Z}(\Phi, \sigma) = \int_{L^\times} \Phi(x)\sigma(x) d^\times x$$

for $\sigma \in \Omega(L^\times)$ and $\Phi \in \mathcal{S}(L)$. The gamma factor

$$\gamma(s, \sigma, \psi_L) = \varepsilon(s, \sigma, \psi_L) \frac{L(1-s, \sigma^{-1})}{L(s, \sigma)}$$

is defined as the proportionality constant of the functional equation

$$\mathcal{Z}(\widehat{\Phi}, \sigma^{-1}\omega_L^{1-s}) = \gamma(s, \sigma, \psi_L)\mathcal{Z}(\Phi, \sigma\omega_L^s),$$

where

$$\widehat{\Phi}(y) = \int_L \Phi(x)\psi_L(yx) dx$$

is the Fourier transform with respect to ψ_L . We repeatedly use the equation

$$(6.1) \quad \gamma(s, \sigma, \psi_L)\gamma(1-s, \sigma^{-1}, \psi_L) = \sigma(-1),$$

Define the additive character ψ_K on K by $\psi_K(x) = \psi(\text{Tr}_F^K(\mathfrak{T}^{-1}x))$, where

$$\mathfrak{T} = \begin{cases} e_1 - e_2 & \text{if } K = Fe_1 \oplus Fe_2 \text{ is split,} \\ \theta - \bar{\theta} & \text{if } K \text{ is not split.} \end{cases}$$

6.2. Principal series representations. Let $\chi_1, \chi_2, \sigma \in \Omega(F^\times)$ be such that $\chi_1\chi_2 = \sigma^{-2}$. We consider the principal series representation

$$\pi = I_2(\chi) = \chi_1 \times \chi_2 \rtimes \sigma := \text{Ind}_{\mathcal{B}_2}^{\text{GSp}_4(F)} \chi,$$

where the character χ of \mathcal{B}_2 is defined by

$$\chi(\mathbf{m}(\text{diag}[a, d], \lambda)u) = \chi_1(a)\chi_2(d)\sigma(\lambda) \quad (a, d, \lambda \in F^\times, u \in \mathcal{U}_2(F)).$$

The induction is always normalized, i.e., the space V of π consists of \mathbb{C} -valued functions on $\text{GSp}_4(F)$ with the transformation property

$$\phi(\mathbf{m}(\text{diag}[a, d], \lambda)ug) = \chi_1(a)\chi_2(d)\sigma(\lambda)|a|^2|d||\lambda|^{-3/2}\phi(g).$$

If χ_1 and χ_2 are unitary, then π is irreducible by Lemma 3.2 of [32].

Then π is equivalent to the induced representation $I(\pi_0, \sigma)$, where we put $\pi_0 = I_1(\chi_1\sigma)$. A ψ -Whittaker functional \mathbf{W} on $I_1(\chi_1\sigma)$ is constructed by the Jacquet integral

$$\mathbf{W}_f(g) = \mathbf{W}(\pi(g)f) := \int_F^{\text{st}} f(J_1\mathbf{n}(x)g)\psi(-x) dx.$$

We define the $\text{GL}_2(F)$ -invariant pairing $b_{\mathbf{W}} : \pi_0 \times \pi_0^\vee \rightarrow \mathbb{C}$ by

$$b_{\mathbf{W}}(f, f') = \int_{F^\times} \mathbf{W}_f(\text{diag}[a, 1])\mathbf{W}_{f'}(\text{diag}[-a, 1]) da$$

and identify π_0^\vee with $I(\pi_0^\vee, \sigma^{-1}) \simeq \chi_1^{-1} \times \chi_2^{-1} \rtimes \sigma^{-1}$ via the pairing

$$b_{\mathbf{W}}^\sharp(\phi, \phi') = \int_{\text{Sym}_2(F)} b_{\mathbf{W}}(\phi(w_s\mathbf{n}(z)), \phi'(w_s\mathbf{n}(z))) dz.$$

For a Weyl element w of $\text{GSp}_4(F)$ we define $\chi^w \in \Omega(\mathcal{T}_2)$ by $\chi^w(t) = \chi(w^{-1}tw)$ and define the intertwining operator $M_w(\chi) : I_2(\chi) \rightarrow I_2(\chi^w)$ by the integral

$$[M_w(\chi)\phi](g) = \int_{\mathcal{U}_2 \cap w\mathcal{U}_2w^{-1} \setminus \mathcal{U}_2} \phi(w^{-1}ug) du.$$

This integral is absolutely convergent if χ lies in some open set, and can be meromorphically continued to all χ . Let Σ_+ be the set of positive roots of

GSp_4 . For each $\alpha \in \Sigma$, let G_α be the derived group of the centralizer in GSp_4 of the kernel of α . Then G_α has relative semi-simple rank one. Letting $\iota_\alpha : \mathrm{SL}_2 \rightarrow G_\alpha$ be the relevant homomorphism, we define $\chi_\alpha \in \Omega(F^\times)$ by $\chi_\alpha(a) = \chi(\iota_\alpha(\mathrm{diag}[a, a^{-1}]))$ for $a \in F^\times$. Now we define the normalized intertwining operator

$$M_w^*(\chi) = \prod_{\alpha \in \Sigma_+, \alpha^w \notin \Sigma_+} \gamma(0, \chi_\alpha, \psi) \cdot M_w(\chi).$$

For example,

$$\begin{aligned} M_{w^\dagger}^*(\chi) &= \gamma(0, \chi_1 \chi_2^{-1}, \psi) \gamma(0, \chi_1 \chi_2, \psi) \gamma(0, \chi_1, \psi) \gamma(0, \chi_2, \psi) M_{w^\dagger}(\chi), \\ M_{w_s}^*(\chi) &= \gamma(0, \chi_1 \chi_2, \psi) \gamma(0, \chi_1, \psi) \gamma(0, \chi_2, \psi) M_{w_s}(\chi). \end{aligned}$$

6.3. Toric periods on principal series representations. Let $\pi_0 = I_1(\mu)$. We define the toric period of $f \in I_1(\mu)$ in the split case by

$$\mathbf{T}_\Lambda^\mu(f) = \int_{F^\times} f(\mathfrak{J} \mathrm{diag}[a, 1]) \Lambda_0(a)^{-1} d^\times a, \quad \mathfrak{J} = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix},$$

where we have written $\Lambda = (\Lambda_0, \Lambda_0^{-1})$, and in the non-split case by

$$\mathbf{T}_\Lambda^\mu(f) = \int_{F^\times \setminus K^\times} f(\iota(t)) \Lambda(t)^{-1} dt$$

otherwise. The former integral is convergent if $\Re \mu > -\frac{1}{2}$.

Lemma 6.1. *If K/F is split, then for $f \in I_1(\mu)$*

$$\mathbf{T}_\Lambda^\mu = \gamma\left(\frac{1}{2}, \mu^{-1} \Lambda_0^{-1}, \psi\right) \mathbf{T}_\Lambda^{\mathbf{W}}(f).$$

Proof. For each $\Phi = \Phi_1 \otimes \Phi_2 \in \mathcal{S}(F \oplus F)$ we define the Godement section $f = f_\mu^\Phi$ as in (6.2). The left hand side equals

$$\begin{aligned} \int_{F^\times} \mu(a) |a|_F^{1/2} \int_{F^\times} \Phi\left((0, b) \begin{pmatrix} 0 & -1 \\ a & 1 \end{pmatrix}\right) \mu(b)^2 |b|_F \Lambda_0(a)^{-1} d^\times b d^\times a \\ = \mathcal{Z}(\Phi_1, \mu \Lambda_0^{-1} \omega_F^{1/2}) \mathcal{Z}(\Phi_2, \mu \Lambda_0 \omega_F^{1/2}). \end{aligned}$$

The right hand side equals the product of $\gamma\left(\frac{1}{2}, \mu^{-1} \Lambda_0^{-1}, \psi\right)$ and

$$\begin{aligned} \int_{F^\times} \int_F^{\mathrm{st}} \mu(a) |a|_F^{1/2} \int_{F^\times} \Phi\left((0, b) \begin{pmatrix} 0 & -1 \\ a & x \end{pmatrix}\right) \mu(b)^2 |b|_F \frac{\psi(-x)}{\Lambda_0(a)} d^\times b dx d^\times a \\ = \mathcal{Z}(\Phi_1, \mu \Lambda_0^{-1} \omega_F^{1/2}) \mathcal{Z}(\widehat{\Phi}_2, \mu^{-1} \Lambda_0^{-1} \omega_F^{1/2}) (\mu \Lambda_0)(-1). \end{aligned}$$

The lemma follows from the functional equation for Tate's local integral. \square

We associate to $\Lambda \in \Omega^1(F^\times \setminus K^\times)$ the toric integral

$$P_\Lambda \in \mathrm{Hom}_{K^\times \times K^\times}(I_1(\mu) \boxtimes I_1(\mu^{-1}), \Lambda \boxtimes \Lambda)$$

by the convergent integral

$$P_\Lambda(f, f') = L(1, \tau_{K/F}) \int_{F^\times \backslash K^\times} b_{\mathbf{W}}(\pi_0(t)f, \pi_0(\mathbf{J})f') \Lambda^{-1}(t) dt.$$

The normalized intertwining operator $\mathcal{M}(\mu_1, \mu_2) : I(\mu_1, \mu_2) \rightarrow I(\mu_2, \mu_1)$ is defined by the integral

$$[\mathcal{M}(\mu_1, \mu_2)f](g) := \gamma(0, \mu_1 \mu_2^{-1}, \psi) \int_F f(J_1 \mathbf{n}(x)g) dx$$

if $\Re(\mu_1 \mu_2^{-1}) > 1$, and by meromorphic continuation otherwise. To simplify notation, we will write $\mathcal{M}(\sigma) = \mathcal{M}(\sigma, \sigma^{-1})$.

Lemma 6.2.

$$\mathbf{T}_\Lambda^{\sigma^{-1}} \circ \mathcal{M}(\sigma) = \gamma\left(\frac{1}{2}, \sigma_K \Lambda, \psi_K\right) \mathbf{T}_\Lambda^\sigma.$$

Proof. Since $\mathbf{W}(\mathcal{M}(\sigma)f) = \mathbf{W}(f)$ for the choice of a normalization of the intertwining operator, if K/F is split, then Lemma 6.1 gives

$$\mathbf{T}_\Lambda^{\sigma^{-1}} \circ \mathcal{M}(\sigma) = \frac{\gamma\left(\frac{1}{2}, \sigma \Lambda_0^{-1}, \psi\right)}{\gamma\left(\frac{1}{2}, \sigma^{-1} \Lambda_0^{-1}, \psi\right)} \mathbf{T}_\Lambda^\sigma = \gamma\left(\frac{1}{2}, \sigma_K \Lambda, \psi_K\right) \mathbf{T}_\Lambda^\sigma.$$

Let K be a field. To each $\Phi \in \mathcal{S}(F \oplus F)$ we associate the Godement section

$$(6.2) \quad f_\sigma^\Phi(g) = \sigma(\det g) |\det g|_F^{1/2} \int_{F^\times} \Phi((0, b)g) \sigma(b)^2 |b|_F d^\times b \in I_1(\sigma).$$

We shall identify Φ with a Bruhat-Schwartz function on K in such a way that $\Phi(a\theta + b) = \Phi(a, b)$. Define the Fourier transforms of Φ by

$$\widehat{\Phi}(z) = \int_K \Phi(x) \psi_K(xz) dx, \quad \widetilde{\Phi}(z) = \int_K \Phi(x) \psi_K(x\bar{z}) dx.$$

The proof of Lemma 14.7.1 of [22] tells us that

$$(6.3) \quad \mathcal{M}(\sigma) f_\sigma^\Phi = f_{\sigma^{-1}}^{\widetilde{\Phi}}.$$

Notice that $\psi_K((a\theta + b)(x\bar{\theta} + y)) = \psi(ay - bx)$.

Observe that

$$\mathbf{T}_\Lambda^\sigma(f_\sigma^\Phi) = \int_{F^\times \backslash K^\times} \int_{F^\times} \Phi(bt) \frac{\sigma(b^2 t \bar{t}) |b^2 t \bar{t}|_F^{1/2}}{\Lambda(t)} d^\times b d^\times t = \mathcal{Z}(\Phi, \sigma_K \Lambda^{-1} \omega_K^{1/2}).$$

We define $\Phi^\tau \in \mathcal{S}(K)$ by $\Phi^\tau(x) = \Phi(\bar{x})$. Since $\widetilde{\Phi}(z) = \widehat{\Phi^\tau}(-z)$ and $\Lambda(-\bar{t}) = \Lambda(t)^{-1}$ for $t \in K^\times$, the lemma follows again from (6.3) and the functional equation for $\mathrm{GL}_1(K)$. \square

6.4. Factorizations. Given $\phi \in \pi = I_2(\chi)$ and $\phi' \in I_2(\chi^{-1})$, we define

$$J_S^\Lambda(\phi, \phi') = \int_{F^\times \backslash K^\times} \int_{\text{Sym}_2(F)}^{\text{st}} b_{\mathbf{W}}^\sharp(\pi(\mathbf{n}(z)\mathbf{t}(t))\phi, \phi') \overline{\Lambda^S(\mathbf{n}(z)\mathbf{t}(t))} dt dz.$$

Put $\chi_1 = \mu\sigma^{-1}$ and $\chi_2 = \mu^{-1}\sigma^{-1}$. Then $I(\pi_0, \sigma)$ is equivalent to $I_2(\chi)$. Let π'_0 be a generic irreducible subrepresentation of $I_1(\mu)$. Since \mathbf{T}_Λ^μ is necessarily proportional to $\mathbf{T}_\Lambda^{\mathcal{W}}$ on π'_0 by uniqueness, Lemma 3.3 allows us to define the Bessel period $\mathbf{B}_{S', \Lambda}^\chi \in \text{Hom}_{R_{S'}}(I(\pi'_0, \sigma), \Lambda_{S'})$ by

$$\mathbf{B}_{S', \Lambda}^\chi(\phi) = \lim_{i \rightarrow \infty} \int_{\text{Sym}_2^i} \mathbf{T}_\Lambda^\mu(\phi(w_s \mathbf{n}(z))) \psi_{S'}(-z) dz.$$

If K is a field, then the pairing

$$I_1(\mu) \otimes I_1(\mu^{-1}) \ni f \otimes f' \mapsto \int_{F^\times \backslash K^\times} f(c) f'(c) dc$$

is also $\text{GL}_2(F)$ -invariant as $\text{GL}_2(F) = \mathcal{B}_1 K^\times$.

Lemma 6.3. *If K/F is not split, then for $f \in I_1(\mu)$ and $f' \in I_1(\mu^{-1})$*

$$b_{\mathbf{W}}(f, f') = \mu(-1) \frac{|\mathfrak{d}_F|^{1/2} \zeta(1)}{|\mathfrak{d}_K|^{1/2} L(1, \tau_{K/F})} \int_{F^\times \backslash K^\times} f(c) f'(c) dc.$$

Proof. Define $f_\mu^\dagger \in I_1(\mu)$ by $f_\mu^\dagger(\mathcal{B}_1) = 0$ and $f_\mu^\dagger(J_1 \mathbf{n}(z)) = \mathbb{1}_{\mathfrak{o}_F}(z)$. Then

$$\mathbf{W}_{f_\mu^\dagger}(\text{diag}[a, 1]) = \mu(a)^{-1} |a|^{1/2} \mathbb{1}_{\mathfrak{o}_F}(a)$$

and hence $b_{\mathbf{W}}(h_\mu, f_{\mu^{-1}}^\dagger) = \mu(-1) \zeta(1)$. Since

$$(6.4) \quad \iota(x) = \begin{pmatrix} x + \text{Tr}(\theta) & -\text{N}(\theta) \\ 1 & x \end{pmatrix} = \begin{pmatrix} \text{N}(x + \theta) & x + \text{Tr}(\theta) \\ 0 & 1 \end{pmatrix} J_1 \mathbf{n}(x),$$

we have

$$f_\mu^\dagger(\iota(x + \theta)) = \mu_K(x + \theta) |x + \theta|_K^{1/2} \mathbb{1}_{\mathfrak{o}_F}(x)$$

for $x \in F$. Since $f_\mu^\dagger(F^\times) = 0$,

$$\int_{F^\times \backslash K^\times} f_\mu^\dagger(c) f_{\mu^{-1}}^\dagger(c) dc = \frac{|\mathfrak{d}_K|^{1/2}}{|\mathfrak{d}_F|^{1/2}} \int_{F^\times} (f_\mu^\dagger f_{\mu^{-1}}^\dagger)(\iota(x + \theta)) \frac{L(1, \tau_{K/F})}{|x + \theta|_K} dx$$

The identity therefore holds if $f = f_\mu^\dagger$ and $f' = f_{\mu^{-1}}^\dagger$. Since the $\text{GL}_2(F)$ -invariant pairings must be proportional, it holds in general. \square

Proposition 6.4. *For $f \in I_1(\mu)$ and $f' \in I_1(\mu^{-1})$*

$$P_\Lambda(f, f') = \mu(-1) \zeta(1) |\mathfrak{d}_F|^{1/2} |\mathfrak{d}_K|^{-1/2} \mathbf{T}_\Lambda^\mu(\pi_0(\varsigma) f) \mathbf{T}_{\Lambda^{-1}}^{\mu^{-1}}(\pi_0(\varsigma \mathbf{J}) f').$$

Proof. Set $h = \pi_0(\varsigma) f$ and $h' = \pi_0(\varsigma \mathbf{J}) f'$. In the split case we have

$$\zeta(1)^{-1} P_\Lambda(f, f') = \int_{F^\times \backslash K^\times} b_{\mathbf{W}}(\pi_0(\iota_\varsigma(t)) h, h') \Lambda(t)^{-1} d^\times t$$

$$= \int_{F^\times} \int_{F^\times} \mathbf{W}_h(\mathbf{t}(ba)) \mathbf{W}_{h'}(\mathbf{t}(-b)) \Lambda_0(a)^{-1} d^\times b d^\times a.$$

Lemma 6.1 now proves the declared identity.

Next we shall prove the non-split case. Lemma 6.3 gives

$$P_\Lambda(f, f') = \mu(-1) \zeta(1) \frac{|\mathfrak{d}_F|^{1/2}}{|\mathfrak{d}_K|^{1/2}} \int_{F^\times \setminus K^\times} \int_{F^\times \setminus K^\times} f(ct) f'(c\mathbf{J}) \Lambda(t)^{-1} dc dt.$$

The double integral above is clearly equal to $\mathbf{T}_\Lambda^\mu(f) \mathbf{T}_{\Lambda^{-1}}^{\mu^{-1}}(\pi_0(\mathbf{J})f')$. \square

Proposition 6.5. *For $\phi \in I_2(\chi)$ and $\phi' \in I_2(\chi^{-1})$*

$$J_S^\Lambda(\phi, \phi') = \frac{\mu(-1) \zeta(1) |\mathfrak{d}_F|^{1/2}}{L(1, \tau_{K/F}) |\mathfrak{d}_K|^{1/2}} \mathbf{B}_{S', \Lambda}^\chi(\pi(\mathbf{m}(\varsigma))\phi) \mathbf{B}_{-S', \Lambda^{-1}}^{\chi^{-1}}(\pi(\mathbf{m}(\varsigma)\mathbf{t}(\mathbf{J}))\phi').$$

Proof. One can prove Proposition 6.5 in the same way as in the proof of Proposition 3.4, using Proposition 6.4. \square

6.5. Functional equations for $\mathbf{B}_{S, \Lambda}^\chi$. Our goal is to prove the following functional equation:

Proposition 6.6.

$$\mathbf{B}_{S', \Lambda}^{\chi^{-1}} \circ M_{w_\dagger}^*(\chi) = \gamma \left(\frac{1}{2}, \mu_K \Lambda, \psi_K \right) \gamma \left(\frac{1}{2}, \sigma_K^{-1} \Lambda, \psi_K \right) \mathbf{B}_{S', \Lambda}^\chi.$$

By uniqueness we arrive at a functional equation

$$\mathbf{B}_{S', \Lambda}^{\chi^w} \circ M_w^*(\chi) = c(w, \chi, \Lambda, \psi) \mathbf{B}_{S', \Lambda}^\chi.$$

When both χ and Λ are unramified, the factor $c(w, \chi, \Lambda, \psi)$ and Lemma 6.2 were calculated in [7]. We will generalize these results to ramified characters.

Proposition 6.7.

$$c(s_1, \chi, \Lambda, \psi) = \gamma \left(\frac{1}{2}, (\chi_1 \sigma)_K \Lambda, \psi_K \right), \quad c(s_2, \chi, \Lambda, \psi) = 1.$$

Proof. Let $\phi \in I_2(\chi)$. From the expression

$$\mathbf{B}_{S', \Lambda}^{\chi^{s_1}}(M_{s_1}^*(\chi)\phi) = \lim_{i \rightarrow \infty} \int_{\text{Sym}_i^2} \mathbf{T}_\Lambda^{(\chi_1 \sigma)^{-1}}(\mathcal{M}(\chi_1 \sigma)\phi(w_s \mathbf{n}(z))) \psi_{S'}(-z) dz.$$

We deduce the assertion for s_1 from Lemma 6.2.

To prove the assertion for s_2 , we consider the following embedding

$$\iota : \text{GL}_2 \rightarrow \text{GSp}_4, \quad \iota \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = \begin{pmatrix} ad - bc & & & \\ & a & & b \\ & & 1 & \\ & c & & d \end{pmatrix}$$

and the subgroup U_2'' of N_2 given by

$$U_2'' = \left\{ \mathbf{n} \left(\begin{pmatrix} 0 & y \\ y & w \end{pmatrix} \right) \mid y, w \in F \right\}$$

Then $\iota(J_1) = s_2$, and we can write a unique expression

$$w_s n \mathbf{t}(t) = \iota(J_1 \mathbf{n}(x)) g', \quad g' = s_2 w_s u'' \mathbf{t}(t), \quad u'' \in U_2''.$$

Put $\mathbf{B}_{S,\Lambda}^\chi = \mathbf{B}_{S',\Lambda}^\chi \circ \pi(\mathbf{m}(\varsigma))$ with $\pi = I_2(\chi)$. Recall that the upper left entry of S is 1. If $\Re\sigma > -\frac{1}{2}$, then $\mathbf{B}_{S,\Lambda}^{\chi^{s_2}}(M_{s_2}^*(\chi)\phi)$ is equal to

$$\int_{F^\times \backslash K^\times} \int_{U_2''} \int_F [M_{s_2}^*(\chi)\phi](\iota(J_1 \mathbf{n}(x))g') \psi(-x) dx \overline{\psi_S(u'')\Lambda^{-1}(t)} du'' dt.$$

Note that $\chi^{s_2} = (\chi_1, \chi_2^{-1}, \chi_2\sigma)$. We define the function $f : \mathrm{GL}_2(F) \rightarrow \mathbb{C}$ via $f(A) = \phi(\iota(A)g')$. Clearly,

$$f \in I(\sigma^{-1}, \chi_1\sigma), \quad [M_{s_2}^*(\chi)\phi](\iota(A)g') = [\mathcal{M}(\sigma^{-1}, \chi_1\sigma)f](A).$$

Since $\mathbf{W}(\mathcal{M}(\sigma^{-1}, \chi_1\sigma)h) = \mathbf{W}(h)$ for any $h \in I(\sigma^{-1}, \chi_1\sigma)$, we find that

$$\begin{aligned} & \mathbf{B}_{S,\Lambda}^{\chi^{s_2}}(M_{s_2}^*(\chi)\phi) \\ &= \int_{F^\times \backslash K^\times} \int_{U_2''} \int_F \phi(\iota(J_1 \mathbf{n}(x))g') \psi(-x) dx \overline{\psi_S(u'')\Lambda^{-1}(t)} du'' dt = \mathbf{B}_{S,\Lambda}^\sigma(\phi), \end{aligned}$$

which proves the assertion for s_2 . \square

Now we will prove Proposition 6.6. Observe that

$$\chi^{s_2} = (\chi_1, \chi_2^{-1}, \chi_2\sigma), \quad \chi^{s_1 s_2} = (\chi_2^{-1}, \chi_1, \chi_2\sigma), \quad \chi^{s_2 s_1 s_2} = (\chi_2^{-1}, \chi_1^{-1}, \sigma^{-1})$$

and $\chi^{s_1 s_2 s_1 s_2} = \chi^{-1}$. Proposition 6.7 gives

$$\begin{aligned} & c(w_\dagger, \chi, \Lambda, \psi) \\ &= c(s_1, \chi^{s_2 s_1 s_2}, \Lambda, \psi) c(s_2, \chi^{s_1 s_2}, \Lambda, \psi) c(s_1, \chi^{s_2}, \Lambda, \psi) c(s_2, \chi, \Lambda, \psi) \\ &= \gamma \left(\frac{1}{2}, (\chi_2\sigma)_K^{-1} \Lambda, \psi_K \right) \cdot 1 \cdot \gamma \left(\frac{1}{2}, (\chi_1 \chi_2 \sigma)_K \Lambda, \psi_K \right) \cdot 1. \end{aligned}$$

6.6. Local coefficients. The factor $c(w, \chi, \Lambda, \psi)$ is an analogue of the local coefficients for Bessel models instead of Whittaker models and have been studied in [11] in more general situations. We will discuss a functional equation for the Bessel periods $\mathbf{B}_{S',\Lambda}^{\mathcal{W},\sigma}$ introduced in Definition 3.2, which is of interest in its own right. This result is not used in our later discussion and the reader can skip the rest of this section and continue reading from the next section onwards.

Conjecture 6.8. Let π_0 be an irreducible admissible unitary generic representation of $\mathrm{PGL}_2(F)$ and $\sigma \in \Omega(F^\times)$. Then

$$\mathbf{B}_{S',\Lambda}^{\mathcal{W},\sigma^{-1}} \circ M^*(\pi_0, \sigma) = \gamma \left(\frac{1}{2}, \sigma_K^{-1} \Lambda, \psi_K \right) \mathbf{B}_{S',\Lambda}^{\mathcal{W},\sigma}.$$

We will prove this conjecture, provided that π_0 is not supercuspidal. One will be able to prove the supercuspidal case by the global method.

Proposition 6.9. *Conjecture 6.8 is true if π_0 is not supercuspidal and $-\frac{1}{2} < \Re \sigma < \frac{1}{2}$.*

Proof. By uniqueness the Bessel period admits a functional equation

$$\mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma^{-1}} \circ M^*(\pi_0, \sigma) = c(\pi_0, \sigma, \Lambda, \psi) \mathbf{B}_{S', \Lambda}^{\mathcal{W}, \sigma}.$$

Take $\mu \in \Omega(F^\times)$ with $\Re \mu > -\frac{1}{2}$ such that π_0 is equivalent to the (unique) irreducible subrepresentation of the principal series representation $I_1(\mu)$ of $\mathrm{GL}_2(F)$. Since $\chi^{w_s} = (\chi_2^{-1}, \chi_1^{-1}, \sigma^{-1})$, we have $I_2(\chi^{w_s}) = I(\pi_0, \sigma^{-1})$. The restriction of $M_{w_s}^*(\chi)$ to $I(\pi_0, \sigma)$ agrees with the normalized intertwining operator $M^*(\pi_0, \sigma)$, and consequently

$$c(\pi_0, \sigma, \Lambda, \psi) = c(w_s, \chi, \Lambda, \psi).$$

Since $c(\pi_0, \sigma \omega_F^s, \psi)$ is a meromorphic function in s , it suffices to prove the equality for σ in a general position. We may therefore suppose that $\gamma(s, \sigma \Lambda_0^{-1}, \psi)$ and $\gamma(s, \sigma^{-1} \chi_1^{-1} \Lambda_0^{-1}, \psi)$ have no pole or zero at $s = \frac{1}{2}$. Then

$$c(w_s, \chi, \Lambda, \psi) = 1 \cdot c(s_1, \chi^{s_2}, \Lambda, \psi) \cdot 1 = \gamma\left(\frac{1}{2}, \sigma_K^{-1} \Lambda, \psi_K\right)$$

by Proposition 6.7. □

7. THE IWAHORI HECKE ALGEBRAS AND THE ORDINARY PROJECTOR e_{ord}^0

We introduce the ordinary projector on principal series representations of $\mathrm{GSp}_4(F)$. Define the Iwahori subgroup of $\mathrm{GSp}_4(\mathfrak{o}_F)$ by

$$\mathbb{I} = \left\{ g \in \mathrm{GSp}_4(\mathfrak{o}_F) \mid g \equiv \begin{pmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & 0 \\ 0 & 0 & * & * \end{pmatrix} \pmod{\mathfrak{p}} \right\}.$$

We define elements of $\mathrm{GSp}_4(F)$ by

$$\begin{aligned} \delta_1 &= \mathrm{diag}[\varpi, 1, \varpi^{-1}, 1], & \delta_2 &= \mathrm{diag}[-\varpi, -\varpi, 1, 1], \\ w_\dagger &= \begin{pmatrix} 0 & \mathbf{1}_2 \\ -\mathbf{1}_2 & 0 \end{pmatrix}, & s_1 &= \mathbf{m} \left(\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right), \\ s_2 &= \begin{pmatrix} 1 & & & \\ & & 1 & \\ & & & 1 \\ -1 & & & \end{pmatrix}, & \eta &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & \varpi & 0 & 0 \\ \varpi & 0 & 0 & 0 \end{pmatrix}, & s_0 &= \begin{pmatrix} & & & -\frac{1}{\varpi} \\ & 1 & & \\ \varpi & & & \\ & & & 1 \end{pmatrix}. \end{aligned}$$

Observe that

$$w_\dagger = s_1 s_2 s_1 s_2, \quad \eta = s_2 s_1 s_2 \delta_2, \quad s_0 = \eta s_2 \eta^{-1}, \quad \eta s_1 \eta^{-1} = s_1.$$

Let (π, V) be an admissible representation of $\mathrm{GSp}_4(F)$. For an open compact subgroup \mathcal{K} of $\mathrm{GSp}_4(F)$ the subspace

$$V^{\mathcal{K}} = \{\phi \in V \mid \pi(k)\phi = \phi \text{ for } k \in \mathcal{K}\}$$

consists of \mathcal{K} -invariant vectors.

Definition 7.1 (Hecke operators). Given $g \in \mathrm{GSp}_4(F)$, we write $\mathbb{I}g\mathbb{I} = \bigsqcup_{u \in I_g} ug\mathbb{I}$ and define the operator $[\mathbb{I}g\mathbb{I}]$ on $V^\mathbb{I}$ by

$$[\mathbb{I}g\mathbb{I}]v = \sum_{u \in I_g} \pi(ug)v.$$

Put $q_g = [\mathbb{I} : \mathbb{I} \cap g\mathbb{I}g^{-1}] = \sharp I_g$. Define the Hecke operators by

$$U^{\mathcal{Q}} = [\mathbb{I}\delta_1\mathbb{I}], \quad U^{\mathcal{P}} = [\mathbb{I}\delta_2\mathbb{I}].$$

Let \mathcal{T}'_2 be the diagonal torus of Sp_4 and \tilde{N} the normalizer of $\mathcal{T}'_2(F)$ in $\mathrm{Sp}_4(F)$. The Weyl group $W = \tilde{N}/\mathcal{T}'_2(F)$ has 8 elements and is generated by the images of s_1, s_2 . We may view W as a subgroup of $S_{p_4(\mathfrak{o}_F)}$ and will not distinguish in notation between the matrices s_1, s_2 and their images in W . The affine Weyl group $\tilde{W} = \tilde{N}/\mathcal{T}'_2(\mathfrak{o}_F)$ is generated by the images of s_0, s_1 and s_2 . The length $\ell(w)$ of $w \in \tilde{W}$ is defined as the minimum number of uses of s_0, s_1 and s_2 required to express w . If $\ell(ww') = \ell(w)\ell(w')$, then

$$(7.1) \quad q_{ww'} = q_w q_{w'}, \quad [\mathbb{I}ww'\mathbb{I}] = [\mathbb{I}w\mathbb{I}][\mathbb{I}w'\mathbb{I}].$$

Let $\chi_1, \chi_2, \sigma \in \Omega^1(F^\times)^\circ$ be such that $\chi_1\chi_2 = \sigma^{-2}$. We consider the unramified principal series representation

$$\pi = \chi_1 \times \chi_2 \rtimes \sigma = \mathrm{Ind}_{\mathcal{B}_2}^{\mathrm{GSp}_4(F)} \chi,$$

which is irreducible by Lemma 3.2 of [32]. Put

$$\alpha = \chi_1(\varpi), \quad \beta = \chi_2(\varpi), \quad \gamma = \sigma(\varpi), \quad \alpha_0 = \alpha\gamma.$$

The space $V^\mathbb{I}$ has the basis $\{\phi_w\}_{w \in W}$, where ϕ_w is the unique \mathbb{I} -invariant vector of V such that $\phi_w(w) = 1$ and $\phi_w(w') = 0$ for $w \neq w' \in W$. We will primarily be interested in $\phi^\dagger = \phi_{w^\dagger} = \phi_{s_1 s_2 s_1 s_2}$. It is convenient to order the basis as follows:

$$\phi_{\mathbf{1}_4}, \quad \phi_{s_1}, \quad \phi_{s_2}, \quad \phi_{s_2 s_1}, \quad \phi_{s_1 s_2 s_1}, \quad \phi_{s_1 s_2}, \quad \phi_{s_1 s_2 s_1 s_2}, \quad \phi_{s_2 s_1 s_2}.$$

With respect to this basis the actions of $[\mathbb{I}s_i\mathbb{I}]$ and $[\mathbb{I}\eta\mathbb{I}]$ on $V^\mathbb{I}$ are given by

$$[\mathbb{I}s_1\mathbb{I}] = \begin{pmatrix} 0 & q & & & & & & & \\ 1 & q-1 & & & & & & & \\ & & 0 & q & & & & & \\ & & 1 & q-1 & & & & & \\ & & & & q-1 & 1 & & & \\ & & & & q & 0 & & & \\ & & & & & & q-1 & 1 & \\ & & & & & & q & 0 & \end{pmatrix},$$

$$[\mathbb{I}s_2\mathbb{I}] = \begin{pmatrix} 0 & 0 & q & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & q & 0 & 0 & 0 \\ 1 & 0 & q-1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & q & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & q & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & q-1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & q-1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & q-1 & 0 \end{pmatrix}.$$

$$[\mathbb{I}\eta\mathbb{I}] = \begin{pmatrix} & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ & & & & & & & & & \gamma q^{3/2} \\ \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} & \frac{\alpha\beta\gamma}{q^{3/2}} \end{pmatrix}.$$

thanks to Lemma 2.1.1² of [33]. We have $[\mathbb{I}s_0\mathbb{I}] = [\mathbb{I}\eta\mathbb{I}][\mathbb{I}s_2\mathbb{I}][\mathbb{I}\eta\mathbb{I}]^{-1}$. Let

$$\phi_\chi^0 = \phi_{\mathbf{1}_4} + \phi_{s_1} + \phi_{s_2} + \phi_{s_2 s_1} + \phi_{s_1 s_2 s_1} + \phi_{s_1 s_2} + \phi^\dagger + \phi_{s_2 s_1 s_2}$$

be the unique element of π that takes the value 1 on $\mathrm{GSp}_4(\mathfrak{o}_F)$.

Definition 7.2 ((α^{-1}, γ) -stabilizations). Introduce the ordinary projector

$$e_{\mathrm{ord}}^0 := \frac{\alpha}{\gamma^3 q^{13/2}} (U^\mathcal{Q} - q^2/\beta)(U^\mathcal{P} - q^{3/2}/\gamma)(U^\mathcal{P} - q^{3/2}\gamma\alpha)(U^\mathcal{P} - q^{3/2}\gamma\beta)$$

(cf. [28]). Define stabilizations of ϕ_χ^0 by

$$\phi^\ddagger = e_{\mathrm{ord}}^0 \phi_\chi^0,$$

$$\phi^\flat = q^{-15/2} \gamma^{-1} \alpha^3 (U^\mathcal{P} - q^{3/2}\gamma\beta)(U^\mathcal{Q} - q^2\alpha)(U^\mathcal{Q} - q^2\beta)(U^\mathcal{Q} - q^2\beta^{-1}) \phi_\chi^0.$$

Remark 7.3. The operators $U^\mathcal{Q}$ and $U^\mathcal{P}$ are commutative.

Proposition 7.4. (1) The support of ϕ^\ddagger is contained in $\mathcal{B}_2 w_\dagger \mathcal{U}_2(\mathfrak{o}_F)$.

(2) ϕ^\ddagger is an eigenform for both $U^\mathcal{Q}$ and $U^\mathcal{P}$, i.e.,

$$U^\mathcal{Q} \phi^\ddagger = q^2 \alpha^{-1} \phi^\ddagger, \quad U^\mathcal{P} \phi^\ddagger = q^{3/2} \gamma \phi^\ddagger.$$

(3) ϕ^\ddagger and ϕ^\flat are eigenforms for both $U^\mathcal{Q}$ and $U^\mathcal{P}$. Moreover,

$$\phi^\ddagger = (1 - \alpha q^{-1})(1 - \beta q^{-1})(1 - \gamma^2 \alpha^2 q^{-1})(1 - \gamma^{-2} q^{-1}) \phi^\ddagger, \quad \phi^\flat = (\alpha + 1) \phi^\ddagger.$$

Remark 7.5. One can partially deduce (2) from (1), namely,

$$[\mathbb{I}\delta_i\mathbb{I}] \phi^\ddagger = ([\mathbb{I}\delta_i\mathbb{I}] \phi^\ddagger)(w_\dagger) \phi^\ddagger.$$

Let $g \in \mathrm{GSp}_4(F)$ be such that $([\mathbb{I}\delta_i\mathbb{I}] \phi^\ddagger)(g) \neq 0$. There exists $u \in \mathbb{I}$ such that $\phi^\ddagger(g u \delta_i) \neq 0$. We have $g u \delta_i \in \mathcal{B}_2 w_\dagger \mathcal{U}_2(\mathfrak{o}_F)$ in view of (1). Since $\delta_i \mathcal{U}_2(\mathfrak{o}_F) \delta_i^{-1} \subset \mathcal{U}_2(\mathfrak{o}_F)$, we get $g \in \mathcal{B}_2 w_\dagger \mathbb{I} = \mathcal{B}_2 w_\dagger \mathcal{U}_2(\mathfrak{o}_F)$.

Proof. Put

$$\bar{\mathcal{U}}_2 = w_\dagger \mathcal{U}_2 w_\dagger^{-1}, \quad \bar{\mathcal{U}}_2(\mathfrak{p}) = \{u \in \bar{\mathcal{U}}_2(\mathfrak{o}_F) \mid u \equiv \mathbf{1}_4 \pmod{\mathfrak{p}}\}.$$

Thanks to the Iwahori factorization $\mathbb{I} = \bar{\mathcal{U}}_2(\mathfrak{p}) \mathcal{B}_2(\mathfrak{o}_F) = \mathcal{B}_2(\mathfrak{o}_F) \bar{\mathcal{U}}_2(\mathfrak{p})$, we get

$$\mathcal{B}_2 \mathbb{I} w_\dagger \mathbb{I} = \mathcal{B}_2 \bar{\mathcal{U}}_2(\mathfrak{p}) w_\dagger \mathbb{I} = \mathcal{B}_2 w_\dagger \mathbb{I} = \mathcal{B}_2 w_\dagger \bar{\mathcal{U}}_2(\mathfrak{p}) \mathcal{B}_2(\mathfrak{o}_F) = \mathcal{B}_2 w_\dagger \mathcal{U}_2(\mathfrak{o}_F).$$

To prove (2), one can show that

$$U^\mathcal{Q} = [\mathbb{I}s_1\mathbb{I}][\mathbb{I}s_2\mathbb{I}][\mathbb{I}s_1\mathbb{I}][\mathbb{I}s_0\mathbb{I}], \quad U^\mathcal{P} = [\mathbb{I}s_2\mathbb{I}][\mathbb{I}s_1\mathbb{I}][\mathbb{I}s_2\mathbb{I}][\mathbb{I}\eta\mathbb{I}],$$

using (7.1). By direct computations the matrix representation of $U^\mathcal{Q}$ is

²The matrix for $[\mathbb{I}s_1\mathbb{I}]$ in Lemma 2.1.1 of [33] contains a typo.

$$\begin{bmatrix} q^2\alpha & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \alpha q(q-1) & q^2\beta & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & q^2\alpha & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & q(q-1)(\beta+1) & \alpha q(q-1) & \frac{q^2}{\beta} & 0 & 0 & 0 & 0 & 0 \\ (\alpha+1)(q-1) & (q-1)^2 & \alpha(q-1)^2 & \frac{q^2-q}{\beta} & \frac{q^2}{\alpha} & \beta q(q-1) & 0 & 0 & 0 \\ 0 & 0 & \alpha q(q-1) & 0 & 0 & q^2\beta & 0 & 0 & 0 \\ \frac{\alpha}{q}(q-1)^2 & \beta(q-1) & (q-1)\{\alpha(q-1+\frac{1}{q})+1\} & 0 & 0 & (q-1)^2(\beta+1) & \frac{q^2}{\alpha} & \frac{q^2-q}{\beta} & q \\ \alpha(q-1) & 0 & \alpha(q-1)^2 & 0 & 0 & q(q-1)(\beta+1) & 0 & 0 & \frac{q^2}{\beta} \end{bmatrix}$$

and the matrix representation of $U^{\mathcal{P}}$ is given by

$$\begin{bmatrix} \frac{q^{3/2}}{\gamma} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{q^{3/2}}{\gamma} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{q^{1/2}(q-1)}{\gamma} & 0 & \gamma\alpha q^{3/2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{q^{1/2}(q-1)}{\gamma} & 0 & \gamma\alpha q^{3/2} & 0 & 0 & 0 & 0 & 0 \\ \frac{q-1}{\gamma q^{1/2}} & \frac{(q-1)^2}{\gamma q^{1/2}} & 0 & \gamma\alpha q^{1/2}(q-1) & \gamma\beta q^{3/2} & 0 & 0 & 0 & 0 \\ 0 & \frac{q^{1/2}(q-1)}{\gamma} & \frac{q^{1/2}(q-1)}{\gamma\beta} & 0 & 0 & \gamma\beta q^{3/2} & 0 & 0 & 0 \\ \frac{(q-1)^2}{\gamma q^{3/2}} & \frac{(q^3-2q^2+2q-1)}{\gamma q^{3/2}} & \frac{\gamma\alpha(q-1)}{q^{1/2}} & \frac{\gamma\alpha(q-1)^2}{q^{1/2}} & \gamma\beta q^{1/2}(q-1) & 0 & \gamma q^{3/2} & 0 & 0 \\ \frac{q-1}{\gamma q^{1/2}} & \frac{(q-1)^2}{\gamma q^{1/2}} & 0 & \gamma\alpha q^{1/2}(q-1) & 0 & \frac{q^{1/2}(q-1)}{\gamma\alpha} & 0 & \gamma q^{3/2} & 0 \end{bmatrix}$$

From these we can prove (2) and observe that both ϕ^{\ddagger} and ϕ^{\flat} are multiples of ϕ^{\dagger} . By a brute force calculation one can show that

$$\begin{aligned} & \alpha\gamma^{-3}q^{-13/2}(U^{\mathcal{Q}} - q^2\beta^{-1})(U^{\mathcal{P}} - q^{3/2}\gamma^{-1})(U^{\mathcal{P}} - q^{3/2}\gamma\alpha)(U^{\mathcal{P}} - q^{3/2}\gamma\beta)e_0 \\ & = (1 - \alpha q^{-1})(1 - \beta q^{-1})(1 - \gamma^2\alpha^2q^{-1})(1 - \gamma^{-2}q^{-1})e_7, \end{aligned}$$

where $e_0 = {}^t(1, 1, 1, 1, 1, 1, 1, 1)$ and $e_7 = {}^t(0, 0, 0, 0, 0, 0, 1, 0)$. \square

8. EXPLICIT CALCULATIONS OF BESSEL INTEGRALS III: ORDINARY VECTORS

Let $\mu, \sigma \in \Omega(F^\times)^\circ$. Put $\pi_0 = I_1(\mu)$ and $\chi = (\mu\sigma^{-1}, \mu^{-1}\sigma^{-1}, \sigma)$. Let $\pi = I(\pi_0, \sigma) = I_2(\chi)$ be an irreducible unramified unitary principal series representation of $\mathrm{PGSp}_4(F)$. Recall that $\phi^{\ddagger} = e_{\mathrm{ord}}^0\phi_\chi^0$ is the \mathfrak{p} -stabilization of the spherical section ϕ_χ^0 in $I_2(\chi)$ obtained by the ordinary projector e_{ord}^0 in Definition 7.2. Let $\Lambda \in \Omega(F^\times \setminus K^\times)$.

Definition 8.1. When Λ is trivial on \mathfrak{r}^\times , set $c(\Lambda) = 0$. Otherwise we put

$$c(\Lambda) = \max\{s \in \mathbb{N} \mid \Lambda \text{ is trivial on } 1 + \mathfrak{p}^s\mathfrak{r}\}.$$

For any given positive integer n we put

$$\xi^{(n)} := \mathbf{m}(\zeta_{\mathfrak{p}}^{(n)}), \quad \zeta_{\mathfrak{p}}^{(n)} := \begin{cases} \begin{pmatrix} \theta & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \varpi^n & 0 \\ 0 & 1 \end{pmatrix} & \text{if } K = F \oplus F, \\ \begin{pmatrix} 0 & 1 \\ \varpi^n & 0 \end{pmatrix} & \text{otherwise.} \end{cases}$$

Set $n = \max\{1, c(\Lambda)\}$. Our task in this section is to compute

$$\mathbb{B}_S^\Lambda(\pi(\xi^{(n)})e_{\text{ord}}^0) = \frac{J_S^\Lambda(\pi(\xi^{(n)})e_{\text{ord}}^0\phi_\chi^0, M_{w_\dagger}^*(\chi)\pi(\xi^{(n)})e_{\text{ord}}^0\phi_\chi^0)}{b_{\mathbf{W}}^\sharp(\phi_\chi^0, M_{w_\dagger}^*(\chi)\phi_\chi^0)},$$

where we use the pairing $b_{\mathbf{W}}^\sharp$ to define J_S^Λ . Recall the unique section $\phi^\dagger \in I_2(\chi)$ supported in $\mathcal{B}_2 w_\dagger N_2(\mathfrak{o}_F)$ with $\phi^\dagger(w_\dagger) = 1$. Since $(\varsigma^{(n)})^{-1}\mathbf{J}\varsigma^{(n)} \in \mathfrak{I}$, the following result readily follows upon combining Propositions 6.5 and 6.6:

$$(8.1) \quad L(1, \tau_{K/F}) \frac{J_S^\Lambda(\pi(\xi^{(n)})\phi^\dagger, M_{w_\dagger}^*(\chi)\pi(\xi^{(n)})\phi^\dagger)}{\zeta(1)\gamma\left(\frac{1}{2}, \mu_K\Lambda, \psi_K\right)\gamma\left(\frac{1}{2}, \sigma_K^{-1}\Lambda, \psi_K\right)} \\ = |\mathfrak{d}_F|^{1/2} |\mathfrak{d}_K|^{-1/2} \mathbf{B}_{S', \Lambda}^\chi(\pi(\mathbf{m}(\varsigma\varsigma^{(n)}))\phi^\dagger) \mathbf{B}_{-S', \Lambda^{-1}}^\chi(\pi(\mathbf{m}(\varsigma\varsigma^{(n)}))\phi^\dagger).$$

We retain the notation in the previous section. Put

$$\alpha = \mu(\varpi), \quad \gamma = \sigma(\varpi).$$

In the split case we write $\Lambda = (\Lambda_0, \Lambda_0^{-1})$. Set

$$\pi_0 \simeq I_1(\mu), \quad \chi = (\mu\sigma^{-1}, \mu^{-1}\sigma^{-1}, \sigma), \quad \pi = I_2(\chi) \simeq I(\pi_0, \sigma).$$

Definition 8.2. Define the modified \mathfrak{p} -Euler factor

$$e(\pi, \Lambda) = \frac{\alpha^{c(\Lambda)}}{L\left(\frac{1}{2}, (\chi_1\sigma)_K\Lambda\right) L\left(\frac{1}{2}, \sigma_K^{-1}\Lambda\right)}.$$

Recall the element $f_\mu^\dagger \in I_1(\mu)$ defined in the proof of Lemma 6.3.

Lemma 8.3. *We have*

$$\mathbf{T}_\Lambda^\mu(\pi_0(\varsigma\varsigma_{\mathfrak{p}}^{(n)})f_\mu^\dagger) = \frac{|\mathfrak{d}_K|_K^{1/2} L(1, \tau_{K/F})}{|\mathfrak{d}_F|^{1/2} (\alpha\gamma)^n q^{n/2}} \times \begin{cases} \Lambda_0(-1) & \text{if } K = F \oplus F, \\ 1 & \text{otherwise.} \end{cases}$$

Proof. Note that

$$\varsigma\varsigma_{\mathfrak{p}}^{(n)} = \begin{cases} \begin{pmatrix} \mathbb{1}\varpi^n & -1 \\ 0 & 1 \end{pmatrix} & \text{if } K/F \text{ is split,} \\ \begin{pmatrix} 0 & 1 \\ \varpi^n & 0 \end{pmatrix} & \text{if } K/F \text{ is non-split.} \end{cases}$$

Since $\mathcal{W}_{f_\mu^\dagger}(\text{diag}[a, 1]) = \mu(a)^{-1}|a|^{1/2}\mathbb{1}_{\mathfrak{o}_F}(a)$, if K/F is split, then

$$\frac{\mathbf{T}_\Lambda(\pi_0(\varsigma\varsigma_{\mathfrak{p}}^{(n)})f_\mu^\dagger)}{\gamma\left(\frac{1}{2}, \mu^{-1}\Lambda_0^{-1}, \psi\right)} = \int_{F^\times} \mathcal{W}_{\mu^{-1}}^\dagger \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \mathbb{1}\varpi^n & -1 \\ 0 & 1 \end{pmatrix} \right) \Lambda_0(a)^{-1} d^\times a \\ = \int_{F^\times} \psi(-a)|a\varpi^n|^{1/2} \mu(a\varpi^n)^{-1} \mathbb{1}_{\mathfrak{o}_F}(a\varpi^n) \Lambda_0(a)^{-1} d^\times a \\ = \mathcal{Z}(\Phi, \Lambda_0^{-1}\mu^{-1}\omega_F^{1/2}) \Lambda_0(\varpi)^n,$$

where $\Phi(x) = \psi\left(-\frac{x}{\varpi^n}\right) \mathbb{1}_{\mathfrak{o}_F}(x)$. Since $\widehat{\Phi}(x) = \mathbb{1}_{\varpi^{-n} + \mathfrak{o}_F}$, we obtain

$$\mathcal{Z}(\Phi, \Lambda_0^{-1} \mu^{-1} \omega_F^{1/2}) = \frac{\mathcal{Z}(\widehat{\Phi}, \Lambda_0 \mu \omega_F^{1/2})}{\gamma\left(\frac{1}{2}, \Lambda_0^{-1} \mu^{-1}, \psi\right)} = \frac{(\alpha\gamma)^{-n} \Lambda_0(\varpi)^{-n} q^{n/2}}{\gamma\left(\frac{1}{2}, \Lambda_0^{-1} \mu^{-1}, \psi\right)} \text{vol}(\varpi^{-n} + \mathfrak{o}_F).$$

Observe that $\text{vol}(\varpi^{-n} + \mathfrak{o}_F) = \text{vol}(1 + \mathfrak{p}^n) = q^{-n} \zeta(1)$.

Next we assume K to be a field. Since

$$K^\times = F^\times (1 + \mathfrak{o}_F \theta) \bigsqcup F^\times (\mathfrak{p} + \theta),$$

we use the formula

$$\int_{F^\times \setminus K^\times} f(t) dt = \int_{\mathfrak{o}_F} f(\iota(1 + y\theta)) d'y + \int_{\mathfrak{p}} f(\iota(y + \theta)) \frac{d'y}{|y + \theta|_K},$$

where $d'y$ is the Haar measure on F giving \mathfrak{o}_F the volume $L(1, \tau_{K/F}) \frac{|\mathfrak{d}_K|_K^{1/2}}{|\mathfrak{d}_F|^{1/2}}$.

Since $f_\mu^\dagger(\iota(y + \theta) \varsigma_{\mathfrak{p}}^{(n)}) = 0$ by (6.4),

$$\begin{aligned} \mathbf{T}_\Lambda^\mu(\pi_0(\varsigma_{\mathfrak{p}}^{(n)} f_\mu^\dagger)) &= \int_{\mathfrak{o}_F} f_\mu^\dagger \left(\begin{pmatrix} 1 + y\text{Tr}(\theta) & -y\mathbf{N}(\theta) \\ y & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ \varpi^n & 0 \end{pmatrix} \right) \Lambda(1 + y\theta) d'y \\ &= \int_{\mathfrak{o}_F} f_\mu^\dagger \left(\begin{pmatrix} \mathbf{N}(1 + y\theta) & * \\ 0 & \varpi^n \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & \varpi^{-n}y \end{pmatrix} \right) \Lambda(1 + y\theta) d'y \\ &= \mu(\varpi)^{-n} q^{n/2} \int_{\mathfrak{o}_F} \mathbb{1}_{\mathfrak{o}_F}(\varpi^{-n}y) d'y. \end{aligned}$$

This finishes the proof. \square

Lemma 8.4. *We have*

$$\mathbf{B}_{S, \Lambda}^\chi(\pi(\mathbf{m}(\varsigma)\xi^{(n)})\phi^\dagger) = \gamma^n q^{-3n/2} \mathbf{T}_\Lambda^\mu(\pi_0(\varsigma \varsigma_{\mathfrak{p}}^{(n)}) f_\mu^\dagger).$$

Proof. For any $\phi \in \pi$ we have

$$[\pi(\mathbf{m}(\varsigma)\xi^{(n)})\phi](w_s \mathbf{n}(z)) = \gamma^n q^{3n/2} \pi_0(\varsigma \varsigma_{\mathfrak{p}}^{(n)}) \phi(w_s \mathbf{n}(\varsigma_{\mathfrak{p}}^{(n)-1} \varsigma^{-1} z \varsigma^{-1} t_{\varsigma_{\mathfrak{p}}^{(n)-1}})).$$

Since $S[\varsigma_{\mathfrak{p}}^{(n)}] \in \text{Sym}_2(\mathfrak{o}_F)$ and since $\phi^\dagger(\mathbf{t}(A)w_s \mathbf{n}(z)) = f_\mu^\dagger(A) \mathbb{1}_{\text{Sym}_2(\mathfrak{o}_F)}(z)$ by definition, we find the first identity by (3.7). Remark 3.5 and Lemma 8.3 give the second identity. \square

The main result of this section is the following explicit formula for the Bessel integral of ordinary vectors.

Proposition 8.5. *Let $n = \max\{1, c(\Lambda)\}$ and $\alpha = \chi_1(\varpi)$. Then*

$$\frac{\mathbb{B}_S^\Lambda(\pi(\mathbf{m}(\xi^{(n)})e_{\text{ord}}^0)}{L(1, \tau_{K/F})\zeta(2)\zeta(4)} = \frac{|\mathfrak{d}_K|_K^{1/2} L(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \Lambda)}{|\mathfrak{d}_F|^{1/2} L(1, \text{ad}, \pi)} e(\pi, \Lambda)^2 \alpha^{-2n} q^{-4n}.$$

Proof. It is proved in Proposition 7.4 (2) that

$$e_{\text{ord}}^0 \phi_\chi^0 = \mathbf{d}(\chi)^{-1} \phi^\dagger, \quad \mathbf{d}(\chi) = L(1, \chi_1) L(1, \chi_2) L(1, (\chi_1 \sigma)^2) L(1, \sigma^{-2}).$$

Put $\phi' = \pi(\xi^{(n)})\phi^\dagger$. By Lemmas 8.3, 8.4 and (8.1)

$$\frac{L(1, \tau_{K/F}) J_S^\Lambda(\phi', M_{w_\dagger}^*(\chi)\phi')}{\zeta(1)\gamma\left(\frac{1}{2}, \mu_K\Lambda, \psi_K\right)\gamma\left(\frac{1}{2}, \sigma_K^{-1}\Lambda, \psi_K\right)} = \gamma^{2n} q^{-3n} \frac{|\mathfrak{d}_K|_K^{1/2} L(1, \tau_{K/F})^2}{|\mathfrak{d}_F|^{1/2} (\alpha\gamma)^{2n} q^n}.$$

Since

$$\begin{aligned} \gamma\left(\frac{1}{2}, \sigma_K^{-1}\Lambda, \psi_K\right) &= \gamma^{-2c(\Lambda)} \frac{L\left(\frac{1}{2}, \sigma_K\Lambda^{-1}\right)}{L\left(\frac{1}{2}, \sigma_K^{-1}\Lambda\right)}, \\ \gamma\left(\frac{1}{2}, \mu_K\Lambda, \psi_K\right) &= (\alpha\gamma)^{2c(\Lambda)} \frac{L\left(\frac{1}{2}, \mu_K^{-1}\Lambda^{-1}\right)}{L\left(\frac{1}{2}, \mu_K\Lambda\right)} \end{aligned}$$

by the definition of the gamma factors, we get

$$\begin{aligned} \frac{J_S^\Lambda(\phi', M_{w_\dagger}^*(\chi)\phi')}{\zeta(1)L(1, \tau_{K/F})} &= \frac{|\mathfrak{d}_K|_K^{1/2}}{|\mathfrak{d}_F|^{1/2}} \alpha^{2c(\Lambda)-2n} q^{-4n} \frac{L\left(\frac{1}{2}, \sigma_K\Lambda^{-1}\right) L\left(\frac{1}{2}, \mu_K^{-1}\Lambda^{-1}\right)}{L\left(\frac{1}{2}, \sigma_K^{-1}\Lambda\right) L\left(\frac{1}{2}, \mu_K\Lambda\right)} \\ &= |\mathfrak{d}_F|^{-1/2} |\mathfrak{d}_K|_K^{1/2} L(1/2, \mathrm{Spn}(\pi)_K \otimes \Lambda) e(\pi, \Lambda)^2 \alpha^{-2n} q^{-4n}. \end{aligned}$$

In view of (9) of [2] we have

$$M_{w_\dagger}^*(\chi)\phi_\chi^0 = \frac{L(1, \pi, \mathrm{ad})}{\zeta(1)^2 \mathbf{d}(\chi)^2} \phi_{\chi^{-1}}^0.$$

Let f_μ^0 be a unique section of $I_1(\mu)$ such that $f_\mu^0(\mathrm{GL}_2(\mathfrak{o}_F)) = 1$. Then

$$\mathbf{W}(f_\mu^0) = L(1, \mu^2)^{-1}, \quad b_{\mathbf{W}}(f_\mu^0, f_{\mu^{-1}}^0) = \frac{\zeta(1)L(1, \mathrm{ad}, \pi_0)}{\zeta(2)L(1, \mu^2)L(1, \mu^{-2})} = \frac{\zeta(1)^2}{\zeta(2)}.$$

Since $\phi_\chi^0(\mathfrak{t}(A)w_s) = f_\mu^0(A)$, we have

$$b_{\mathbf{W}}^\#(\phi_\chi^0, \phi_{\chi^{-1}}^0) = \frac{\zeta(1)^2}{\zeta(2)} \int_{\mathrm{Sym}_2(F)} |a(w_s \mathbf{n}(z))|^{-3} dz = \frac{\zeta(1)^3}{\zeta(2)\zeta(4)}.$$

We conclude that

$$\mathbf{d}(\chi)^2 b_{\mathbf{W}}(\phi_\chi^0, M_{w_\dagger}^*(\chi)\phi_\chi^0) = L(1, \pi, \mathrm{ad})\zeta(1)\zeta(2)^{-1}\zeta(4)^{-1}.$$

From these our proof is complete. \square

9. GLOBAL BESSEL PERIODS FOR GU_2^D

9.1. Notation. If L is a number field, then \mathfrak{o}_L is the ring of integers of L , \mathbb{A}_L is the adèle ring of L and $L_\infty = L \otimes_{\mathbb{Q}} \mathbb{R}$ is the infinite part of \mathbb{A}_L . When $L = \mathbb{Q}$, we suppress the subscript L . Let $\bar{\mathbb{Z}}$ be the ring of algebraic integers of $\bar{\mathbb{Q}}$, $\bar{\mathbb{Z}}_\ell$ the ℓ -adic completion of $\bar{\mathbb{Z}}$ in $\mathbb{C}_\ell = \widehat{\bar{\mathbb{Q}}}_\ell$ and $\widehat{\mathbb{Z}} = \prod_\ell \mathbb{Z}_\ell$ the finite completion of \mathbb{Z} . Given an abelian group M , we put $\widehat{M} = M \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}$. In particular, $\mathbb{A}_L = L_\infty \oplus \widehat{L}$. Let $\mathbf{e} = \prod_v \mathbf{e}_v$ denote the standard additive character of \mathbb{A}/\mathbb{Q} such that $\mathbf{e}_\infty(x) = e^{2\pi\sqrt{-1}x}$ for $x \in \mathbb{R}$. Set $\psi^L = \mathbf{e} \circ \mathrm{Tr}_{\mathbb{Q}}^L$. When G is a reductive algebraic group over L , we denote by $\mathcal{A}_{\mathrm{cusp}}(G)$ the space of cusp forms on $G(\mathbb{A}_L)$. For an adèle point $g \in G(\mathbb{A}_L)$ we denote its projections to $G(\widehat{L})$, $G(L_\infty)$ and $G(L_v)$ by $g_{\mathbf{f}}$, g_∞ and g_v , respectively. We fix

once and for all an embedding $\iota_\infty : \bar{\mathbb{Q}} \hookrightarrow \mathbb{C}$ and an isomorphism $j_\ell : \mathbb{C} \simeq \mathbb{C}_\ell$ for each rational prime ℓ . Let $\iota_\ell = j_\ell \circ \iota_\infty$ be their composition. We regard L as a subfield of \mathbb{C} (resp. \mathbb{C}_ℓ) via ι_∞ (resp. ι_ℓ) and $\text{Hom}(L, \bar{\mathbb{Q}}) = \text{Hom}(L, \mathbb{C}_\ell)$.

Let F be a totally real number field of degree d and K a totally imaginary quadratic extension of F . We denote by Δ_F (resp. Δ_K) the discriminant of F (resp. K), by $\mathfrak{d}_F = \prod_{\mathfrak{l}} \mathfrak{d}_{F_{\mathfrak{l}}}$ the different of F , by \mathfrak{D}_F^K the relative different of K/F and by $\tau_{K/F}$ the quadratic Hecke character of \mathbb{A}^\times corresponding to K/F . Fix a square free ideal $\mathfrak{N} = \mathfrak{N}^+ \mathfrak{N}^-$ of \mathfrak{o}_F such that every prime factor of \mathfrak{N}^+ (resp. \mathfrak{N}^-) is split (resp. not split) in \mathfrak{o}_K . Fix a decomposition

$$\mathfrak{N}^+ \mathfrak{o}_K = \mathfrak{N}_0^+ \overline{\mathfrak{N}_0^+}.$$

Suppose that the number of prime factors of \mathfrak{N}^- is even. Then there exists a totally indefinite quaternion algebra D over F of discriminant \mathfrak{N}^- , i.e., D is a central simple algebra of dimension 4 over F such that $D_v := D \otimes_v F_v$ is a division algebra if and only if v divides \mathfrak{N}^- . Put

$$N_{\mathbb{Q}}^D = N_F^D \circ N_F^D, \quad \text{Tr}_{\mathbb{Q}}^D = \text{Tr}_F^D \circ \text{Tr}_F^D, \quad N = \#(\mathfrak{o}_F/\mathfrak{N}), \quad N^- = \#(\mathfrak{o}_F/\mathfrak{N}^-).$$

Once and for all we fix a prime \mathfrak{p} of F , which does not divide \mathfrak{N} , CM type Σ of K and a finite idèle $d_F = (d_{F_{\mathfrak{l}}}) \in \widehat{F}^\times$ such that $d_{F_{\mathfrak{l}}}$ is a generator of the local different $\mathfrak{d}_{F_{\mathfrak{l}}}$ for each finite prime \mathfrak{l} . We identify Σ with the set of real places of F . Fix a maximal order \mathfrak{o}_D of D . For any finite prime \mathfrak{p} we set $\mathfrak{o}_{D_{\mathfrak{p}}} = \mathfrak{o}_D \otimes_{\mathfrak{o}_F} \mathfrak{o}_{F_{\mathfrak{p}}}$. If \mathfrak{p} divides \mathfrak{N}^- , then we write $\mathfrak{P}_{\mathfrak{p}}$ for the maximal ideal of $\mathfrak{o}_{D_{\mathfrak{p}}}$. We choose an element $\theta \in K$ such that

- $\Im \tau(\theta) > 0$ for every $\tau \in \Sigma$;
- $\{1, \theta\}$ is an $\mathfrak{o}_{F_{\mathfrak{l}}}$ -basis of $\mathfrak{o}_{K_{\mathfrak{l}}}$ for every prime \mathfrak{l} dividing $\mathfrak{p} \mathfrak{D}_F^K \mathfrak{N}$;
- θ is a uniformizer of $\mathfrak{o}_{K_{\mathfrak{l}}}$ for every prime \mathfrak{l} ramified in K .

We regard K as a subalgebra of D . Put $S = S_\theta := \frac{1}{2}(\theta - \bar{\theta}) \in D_-(F)$.

Recall that $J_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Put $J_\star = \text{diag}[\mathbf{1}_2, J_1]$. For $v \nmid \mathfrak{N}^-$ we fix an isomorphism $i_v : M_2(F_v) \simeq D_v$ by which we identify $M_{2m}(F_v)$ with $M_m(D_v)$. Since $i_v^{-1}(\bar{x}) = J_1^{-1} i_v^{-1}(x) J_1$ for $x \in D_v$, we arrive at

$$J_\star \text{GU}_2^D(F_v) J_\star^{-1} = \text{GSp}_4(F_v), \quad J_1 D_-(F_v) = \text{Sym}_2(F_v).$$

We identify $\begin{pmatrix} \text{Tr}_F^K(\theta) & -N_F^K(\theta) \\ 1 & 0 \end{pmatrix}$ with θ . Then

$$J_1 S_\theta = \begin{pmatrix} 1 & -\frac{\text{Tr}_F^K(\theta)}{2} \\ -\frac{\text{Tr}_F^K(\theta)}{2} & N_F^K(\theta) \end{pmatrix}.$$

We always take the adèlic measure dg on $\text{PGU}_2^D(\mathbb{A}_F)$ to be the Tamagawa measure. We define the bilinear pairing by

$$\langle \phi, \phi' \rangle = \int_{\text{PGU}_2^D(F) \backslash \text{PGU}_2^D(\mathbb{A}_F)} \phi(g) \phi'(g) dg.$$

Let $\pi \simeq \otimes'_v \pi_v$ be an irreducible admissible tempered representation of $\mathrm{PGU}_2^D(\mathbb{A}_F)$ which is realized on a subspace V of $\mathcal{A}_{\mathrm{cusp}}(\mathrm{PGU}_2^D)$. The space $\mathcal{A}_{\mathrm{cusp}}(\mathrm{PGSp}_4)$ satisfies multiplicity one thanks to the work of Arthur. It is conjectured in general that $\mathcal{A}_{\mathrm{cusp}}(\mathrm{PGU}_2^D)$ satisfies multiplicity one, which we assume. Then since $\pi_v \simeq \pi_v^\vee$ for every v , we have $V = \bar{V} := \{\bar{\phi} \mid \phi \in V\}$. Thus the restriction of the pairing $\langle \cdot, \cdot \rangle$ to $V \times V$ is nondegenerate.

Let dz denote the Tamagawa measure on $D_-(\mathbb{A}_F)$. When \mathfrak{l} and \mathfrak{N}^- are coprime, we take the Haar measure $dz_{\mathfrak{l}}$ on $D_-(F_{\mathfrak{l}})$ so that the measure of $D_-(F_{\mathfrak{l}}) \cap \mathfrak{o}_{D_{\mathfrak{l}}}$ is 1. For each prime factor \mathfrak{q} of \mathfrak{N}^- we take the Haar measure $dz_{\mathfrak{q}}$ on $D_-(F_{\mathfrak{q}})$ so that the measure of $D_-(F_{\mathfrak{q}}) \cap \mathfrak{P}_{\mathfrak{q}}$ is 1. For $v \in \Sigma$ we define the Haar measure dz_v on $D_-(F_v)$ by identifying $D_-(F_v) \simeq \mathrm{Sym}_2(F_v)$ with F_v^3 with respect to the standard basis. Then by Lemma 2.1 of [41]

$$(9.1) \quad dz = \Delta_F^{-3/2} (N^-)^{-2} \prod_v dz_v.$$

Fix a Hecke character $\Lambda \in \Omega^1(K^\times \mathbb{A}_F^\times \backslash \mathbb{A}_K^\times)$. Let dt be the invariant measure on $K^\times \mathbb{A}_F^\times \backslash \mathbb{A}_K^\times$ normalized to have total volume $2A(1, \tau_{K/F})$, where

$$\Lambda(s, \tau_{K/F}) = \pi^{-d(s+1)/2} \Gamma((s+1)/2)^d L(s, \tau_{K/F})$$

is the complete Hecke L -function of $\tau_{K/F}$.

We define the S th Fourier coefficient and the Bessel period relative to S and Λ of a cusp form $\phi \in \mathcal{A}_{\mathrm{cusp}}(\mathrm{PGU}_2^D)$ by

$$\begin{aligned} \mathbf{W}_S(\phi, g) &= \int_{D_-(F) \backslash D_-(\mathbb{A}_F)} \phi(\mathbf{n}(z)g) \overline{\psi^F(\tau(Sz))} dz, \\ B_S^\Lambda(\phi, g) &= \int_{K^\times \mathbb{A}_F^\times \backslash \mathbb{A}_K^\times} \mathbf{W}_S(\phi, \mathbf{t}(t)g) \Lambda(t)^{-1} dt. \end{aligned}$$

Here e is the identity element in $\mathrm{GU}_2^D(\mathbb{A}_F)$. We will write $B_S^\Lambda(\phi) = B_S^\Lambda(\phi, e)$.

9.2. The refined Gross-Prasad conjecture for the Bessel periods.

For each place v we normalize the local Bessel integrals by

$$\mathcal{B}_S^{\Lambda_v} = c(\pi_v, \Lambda_v)^{-1} B_S^{\Lambda_v}, \quad c(\pi_v, \Lambda_v) = \zeta_{F_v}(2) \zeta_{F_v}(4) \frac{L(\frac{1}{2}, \mathrm{Spn}(\pi_v)_{K_v} \otimes \Lambda_v)}{L(1, \tau_{K_v/F_v}) L(1, \pi_v, \mathrm{ad})}.$$

We denote the complete Dedekind zeta function of F by $\xi_F(s)$, the complete adjoint L -function of π by $\Lambda(s, \pi, \mathrm{ad})$ and the complete Godement-Jacquet L -function of an automorphic representation Π of a general linear group by $\Lambda(s, \Pi)$.

We rewrite the theorem of Furusawa and Morimoto in a form which is suitable for our later use (cf. Theorem 1.2).

Theorem 9.1 (Furusawa-Morimoto). *Let $\phi_i = \otimes_v \phi_{i,v}$ for $i = 1, 2, 3, 4$. If $\langle \phi_3, \phi_4 \rangle \neq 0$, then*

$$\frac{B_S^\Lambda(\phi_1) B_S^\Lambda(\phi_2)}{\langle \phi_3, \phi_4 \rangle} = \xi_F(2) \xi_F(4) \frac{\Lambda(\frac{1}{2}, \mathrm{Spn}(\pi)_K \otimes \Lambda)}{(N^-)^2 \Delta_F^{3/2} 2^{\ell(\pi)} \Lambda(1, \pi, \mathrm{ad})} \prod_v \frac{\mathcal{B}_S^{\Lambda_v}(\phi_{1,v}, \phi_{2,v})}{\langle \phi_{3,v}, \phi_{4,v} \rangle_v}.$$

Proof. The Petersson inner product (\cdot, \cdot) is defined by $(\phi, \varphi) = \langle \phi, \bar{\varphi} \rangle$. Fix a factorization $(\cdot, \cdot) = \prod_v (\cdot, \cdot)_v$, where $(\cdot, \cdot)_v$ is an invariant Hermitian perfect pairing on π_v . Furusawa and Morimoto [15] use $(\cdot, \cdot)_v$ to define a local Bessel integral

$$\begin{aligned} \alpha_{S, \Lambda_v}^{\natural}(\phi_v, \varphi_v) &= c(\pi_v, \Lambda_v)^{-1} \\ &\times \int_{F_v^\times \backslash K_v^\times} \int_{D_-(F_v)}^{\text{st}} (\pi_v(\mathbf{n}(z_v)\mathbf{t}(t_v))\phi_v, \varphi_v)_v \overline{\psi_v(\text{Tr}_F^D(Sz_v))\Lambda_v(t_v)} dz_v dt_v \end{aligned}$$

When v is archimedean, the integral is defined via regularization (see [26]).

We fixed a local invariant \mathbb{C} -bilinear pairing $\langle \cdot, \cdot \rangle_v : \pi_v \times \pi_v^\vee \rightarrow \mathbb{C}$, which is equivalent to fixing a *sesquilinear* equivariant isomorphism $^c : \pi_v^\vee \simeq \pi_v$ such that $\langle \phi_v, \varphi_v \rangle_v = (\phi_v, \varphi_v^c)_v$. Fix a factorization $0 \neq \phi = \otimes_v \phi_v \in \pi \simeq \otimes_v \pi_v$ and define a factorization $\pi^\vee \simeq \otimes_v \pi_v^\vee$ by $\bar{\phi} = \otimes_v \phi_v^c$. Then it is formal to see that $\bar{\varphi} = \otimes_v \varphi_v^c$ for all pure tensors $\varphi = \otimes_v \varphi_v \in \pi$.

Theorem 1.2 of [15] gives

$$\frac{B_S^\Lambda(\phi_1)\overline{B_S^\Lambda(\phi)}}{(\phi_3, \bar{\phi}_4)} = \xi_F(2)\xi_F(4) \frac{\Lambda(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \Lambda)}{(N^-)^2 \Delta_F^{3/2} 2^{\ell(\pi)} \Lambda(1, \pi, \text{ad})} \prod_v \frac{\alpha_{S, \Lambda_v}^{\natural}(\phi_{1,v}, \phi_v)}{(\phi_{3,v}, \varphi_{4,v}^c)_v}$$

by the uniqueness of Bessel period and the identity $\frac{(\phi, \phi)}{(\phi_3, \phi_4)} = \prod_v \frac{(\phi_v, \phi_v)_v}{(\phi_{3,v}, \phi_{4,v}^c)_v}$.

We defined the local Bessel integral

$$\mathcal{B}_S^{\Lambda_v}(\phi_v, \varphi_v) = \alpha_{S, \Lambda_v}^{\natural}(\phi_v, \pi_v(\mathbf{m}(\mathbf{J}, -1))\varphi_v^c)$$

in Definition 2.2. Let $\phi = \pi(\mathbf{m}(\mathbf{J}, -1))\bar{\phi}_2$. Since $B_S^\Lambda(\phi) = B_{-S}^{\Lambda^{-1}}(\bar{\phi}_2) = \overline{B_S^\Lambda(\phi_2)}$, we get the stated formula. \square

Remark 9.2. (1) The theorem of Furusawa and Morimoto is a special case of the refined Gross-Prasad conjecture formulated by Liu [26] for $\text{SO}(m) \times \text{SO}(l)$. It is easily seen that $dt = \prod_v dt_v$. Thus $C_{G_0} = 1$ and $\mathcal{B}_{\pi_0}(\varphi_0, \bar{\varphi}_0) = 2\Lambda(1, \tau_{K/F})$ in the notation of [26].

(2) Furusawa and Morimoto [13, 14] proved the Liu's conjecture for irreducible cuspidal tempered representations of $\text{SO}(2n+1)$ and the trivial character of $\text{SO}(2)$. In the course of the proof they verified that π has the weak lift $\text{Spn}(\pi)$ to $\text{GL}_{2n}(\mathbb{A}_F)$ and obtain $L(s, \pi, \text{ad})$ to be the symmetric square L -function of $\text{Spn}(\pi)$, which is holomorphic and nonzero at $s = 1$, for the exterior square L -function of $\text{Spn}(\pi)$ has a pole at $s = 1$.

9.3. A central value formula. Let $\kappa \in \mathbb{N}^\Sigma$ be a tuple of d natural numbers indexed by Σ . We define the action of $\text{GU}_2^D(F_\infty)^\circ$ on the space

$$\mathfrak{H}_2^* := \{Z \in \text{M}_2(F \otimes_{\mathbb{Q}} \mathbb{C}) \mid {}^t(ZJ_1^{-1}) = ZJ_1^{-1}, \Im(ZJ_1^{-1}) > 0\}$$

and the automorphy factor $J_\kappa : \text{GU}_2^D(F_\infty)^\circ \times \mathfrak{H}_2^* \rightarrow \mathbb{C}^\times$ by

$$hZ = (h_v Z_v)_{v \in \Sigma}, \quad h_v Z_v = (a_v Z_v + b_v)(c_v Z_v + d_v)^{-1},$$

$$J_\kappa(h, Z) = \prod_{v \in \Sigma} j(h_v, Z_v)^{\kappa_v}, \quad j(h_v, Z_v) = N_{F_v}^{D_v}(c_v Z_v + d_v) / N_{F_v}^{D_v}(h_v)^{1/2},$$

where we write $h_v = \begin{pmatrix} a_v & b_v \\ c_v & d_v \end{pmatrix}$. Let $\mathbf{i} = \sqrt{-1}J_1 \in \mathfrak{H}_2^*$. Put

$$U_2^\Sigma = \{g \in U_2^D(F_\infty) \mid g(\mathbf{i}) = \mathbf{i}\}.$$

The open compact subgroup $K(\mathfrak{l})$ (resp. $K(\mathfrak{P}_\mathfrak{l})$) of $\mathrm{GU}_2^D(F_\mathfrak{l})$ is defined in (4.2) (resp. (5.5)). The paramodular subgroup of level \mathfrak{N} is defined by

$$\mathcal{K}_D(\mathfrak{N}) = \prod_{\mathfrak{l} \mid \mathfrak{N}^+} J_\star^{-1} K(\mathfrak{l}) J_\star \times \prod_{\mathfrak{l} \mid \mathfrak{N}^-} K(\mathfrak{P}_\mathfrak{l}) \times \prod_{\mathfrak{l} \mid \mathfrak{N}} J_\star^{-1} \mathrm{GSp}_4(\mathfrak{o}_{F_\mathfrak{l}}) J_\star.$$

From now on let π be an irreducible cuspidal automorphic representation of $\mathrm{PGU}_2^D(\mathbb{A}_F)$ whose archimedean component is $\otimes_{v \in \Sigma} D_{\kappa_v}^{(2)}$ and such that $\pi_\mathfrak{l}$ is generic for each finite prime \mathfrak{l} . Let $V_\kappa(\pi, \mathfrak{N})$ denote the subspace of V on which the group $U_2^\Sigma \times \mathcal{K}_D(\mathfrak{N})$ acts by the character $k \mapsto J_\kappa(k_\infty, \mathbf{i})^{-1}$.

Definition 9.3. For $\phi, \phi' \in V_\kappa(\pi, \mathfrak{N})$ we normalize the pairing by

$$\langle \phi, \phi' \rangle_{\mathcal{K}_D(\mathfrak{N})} = \langle \phi, \phi' \rangle \prod_{\mathfrak{l} \mid \mathfrak{N}^+} (q_\mathfrak{l}^2 + 1) \prod_{\mathfrak{l} \mid \mathfrak{N}^-} (q_\mathfrak{l}^2 - 1).$$

Suppose that $\dim V_\kappa(\pi, \mathfrak{N}) = 1$. The representations considered in the next section satisfy and are characterized by this condition (see Remarks 4.3 and 5.1). Fix $0 \neq \phi_\pi \in V_\kappa(\pi, \mathfrak{N})$. Then $\phi_\pi = \otimes_v \phi_v^0$ is a pure tensor of new vectors. Put

$$\epsilon_\mathfrak{l}(\pi) = \varepsilon \left(\frac{1}{2}, \mathrm{Spn}(\pi_\mathfrak{l}) \right), \quad \epsilon_{\mathfrak{N}^+}(\pi) = \prod_{\mathfrak{l} \mid \mathfrak{N}^+} \epsilon_\mathfrak{l}(\pi).$$

Take $\chi_1, \chi_2, \sigma \in \Omega(F_\mathfrak{p}^\times)^\circ$ so that $\pi_\mathfrak{p} \simeq \chi_1 \times \chi_2 \rtimes \sigma$. Put $\alpha_\mathfrak{p} = \chi_1(\varpi_\mathfrak{p})$ and $\gamma_\mathfrak{p} = \sigma(\varpi_\mathfrak{p})$. Define the $(\alpha_\mathfrak{p}^{-1}, \gamma_\mathfrak{p})$ -stabilization of ϕ_π by

$$e_{\mathrm{ord}, \mathfrak{p}}^0 \phi_\pi = (\otimes_{v \neq \mathfrak{p}} \phi_v^0) \otimes e_{\mathrm{ord}, \mathfrak{p}}^0 \phi_\mathfrak{p}^0,$$

where the ordinary projector $e_{\mathrm{ord}, \mathfrak{p}}^0$ is defined in Definition 7.2 with respect to $(\alpha_\mathfrak{p}^{-1}, \gamma_\mathfrak{p})$. For $a \in F_\infty^\times$ put $a^\kappa = \prod_{v \in \Sigma} |a|_{F_v}^{\kappa_v}$. For $\mathfrak{l} \nmid \mathfrak{N}$ we take $\theta_\mathfrak{l} \in K_\mathfrak{l}$ and $A_\mathfrak{l} \in \mathrm{GL}_2(F_\mathfrak{l})$ so that $\mathfrak{o}_{K_\mathfrak{l}} = \mathfrak{o}_{F_\mathfrak{l}} + \mathfrak{o}_{F_\mathfrak{l}} \theta_\mathfrak{l}$ and $J_1 S_{\theta_\mathfrak{l}} = (J_1 S_\theta)[A_\mathfrak{l}]$. Recall

$$\varsigma = \begin{pmatrix} 1 & -\theta \\ -1 & \theta \end{pmatrix}.$$

Definition 9.4. For each positive integer n we define $\varsigma_\mathfrak{p}^{(n)} \in \mathrm{GL}_2(F_\mathfrak{p})$ by

$$\varsigma_\mathfrak{p}^{(n)} = \begin{pmatrix} \vartheta & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \varpi_\mathfrak{p}^n & 0 \\ 0 & 1 \end{pmatrix},$$

where $\vartheta = \theta$ if \mathfrak{p} splits in K and $\vartheta = 0$ otherwise, and define

$$\zeta^{(n)} = \mathbf{m} \left(i_\mathfrak{p} \left(\varsigma_\mathfrak{p}^{(n)} \right) \right) \prod_{\mathfrak{l} \mid \mathfrak{N}^+} \mathbf{m}(\varsigma^{-1}) \prod_{\mathfrak{l} \mid \mathfrak{N}} \mathbf{m}(i_\mathfrak{l}(A_\mathfrak{l})) \prod_{\mathfrak{l}} \mathbf{d}(d_{F_\mathfrak{l}}) \in \mathrm{GU}_2^D(\widehat{F}).$$

Put $\tau_\infty = \prod_{v \in \Sigma} \tau_v$, where $\tau_v = \mathbf{d}(-1) \in \mathrm{GU}_2^D(F_v)$.

Theorem 9.5. *We suppose that Λ_l is unramified for every prime l distinct from \mathfrak{p} . Put $n = \max\{1, c(\Lambda_{\mathfrak{p}})\}$. Assume that $\dim V_\kappa(\pi, \mathfrak{N}) = 1$. Assume that $F_l = \mathbb{Q}_2$ if $l \neq \mathfrak{p}$ and 2 is divisible by l . Then*

$$\begin{aligned} & \frac{B_S^\Lambda(e_{\mathrm{ord}, \mathfrak{p}}^0 \phi_\pi, \zeta^{(n)})^2}{\langle \phi_\pi, \pi(\tau_\infty) \phi_\pi \rangle_{\mathcal{K}_D(\mathfrak{N})} e^{4\pi\sqrt{-1}\mathrm{Tr}_{\mathbb{Q}}^D(\mathrm{Si})}} = \frac{\Delta_F^2 \xi_F(2) \xi_F(4) N_F^D(4S)^\kappa [\mathfrak{o}_K : \mathfrak{o}_F + \theta \mathfrak{o}_F]^{-3}}{\epsilon_{\mathfrak{N}^+}(\pi) \Lambda(\mathfrak{N}_0^+) 2^{2d+\ell(\pi)} \Delta_K^{1/2} N_{\mathbb{Q}}^D(S_\theta)^{3/2}} \\ & \times \frac{e(\pi_{\mathfrak{p}}, \Lambda_{\mathfrak{p}})^2}{\alpha_{\mathfrak{p}}^{2n} q_{\mathfrak{p}}^{4n}} L(1, \tau_{K_{\mathfrak{p}}/F_{\mathfrak{p}}})^2 \frac{\Lambda(\frac{1}{2}, \mathrm{Spn}(\pi)_K \otimes \Lambda)}{N\Lambda(1, \pi, \mathrm{ad})} \prod_{\mathfrak{l} | \mathfrak{N}^- \cap \mathfrak{O}_F^K, \mathfrak{l} = \mathfrak{l}_K^2} (1 - \epsilon_{\mathfrak{l}}(\pi) \Lambda(\mathfrak{l}_K)), \end{aligned}$$

where $e(\pi_{\mathfrak{p}}, \Lambda_{\mathfrak{p}})$ is the \mathfrak{p} -adic multiplier

$$e(\pi_{\mathfrak{p}}, \Lambda_{\mathfrak{p}}) = \alpha_{\mathfrak{p}}^{c(\Lambda_{\mathfrak{p}})} \cdot L\left(\frac{1}{2}, (\chi_1 \sigma)_{K_{\mathfrak{p}}} \Lambda_{\mathfrak{p}}\right)^{-1} L\left(\frac{1}{2}, \sigma_{K_{\mathfrak{p}}}^{-1} \Lambda_{\mathfrak{p}}\right)^{-1}.$$

Proof. Put $\mathcal{B}_S^{\Lambda_l}(H) = \frac{\mathbb{B}_S^{\Lambda_l}(H)}{c(\pi_l, \Lambda_l)}$, where $\mathbb{B}_S^{\Lambda_l}$ is defined with respect to an additive character of order 0. It should be remarked that when F_l is of residual characteristic ℓ , we have defined $\mathcal{B}_S^{\Lambda_l}$ with respect to the additive character $\mathbf{e}_\ell \circ \mathrm{Tr}_{\mathbb{Q}_\ell}^{F_l}$ on F_v . Taking Remark 2.4 into account, we have

$$\mathcal{B}_{S_\theta}^{\Lambda_l}(\pi_l(\mathbf{m}(A_l, d_{F_l}))H\phi_l, \pi_l(\mathbf{m}(A_l, d_{F_l}))H\phi_l) = |\mathfrak{d}_{F_l}|^{-3} |\det A_l|^3 \mathcal{B}_{S_{\theta_l}}^{\Lambda_l}(H).$$

Since $\prod_l |\det A_l|_{F_l}^{-1} = [\mathfrak{o}_K : \mathfrak{o}_F + \theta \mathfrak{o}_F]$, it follows from Theorem 9.1 that

$$\begin{aligned} & \frac{B_S^\Lambda(\phi_\pi^\dagger, \xi^{(n)})^2}{\langle \phi_\pi, \pi(\tau_\infty) \phi_\pi \rangle} = \xi_F(2) \xi_F(4) \frac{D_F^{3/2} \Lambda(\frac{1}{2}, \mathrm{Spn}(\pi)_K \otimes \Lambda)}{[\mathfrak{o}_K : \mathfrak{o}_F + \theta \mathfrak{o}_F]^3 (N^-)^2 2^{\ell(\pi)} \Lambda(1, \pi, \mathrm{ad})} \mathcal{B}_{S_{\theta_{\mathfrak{p}}}}^{\Lambda_{\mathfrak{p}}}(e_{\mathrm{ord}}^0) \\ & \times \prod_{v \in \Sigma} \frac{\mathcal{B}_S^{\Lambda_v}(\phi_v^0, \phi_v^0)}{\mathbf{r}(\phi_v^0, \pi_v(\mathbf{d}(-1))\phi_v^0)} \prod_{\mathfrak{l} | \mathfrak{N}^+} \mathcal{B}_S^{\Lambda_l}(\pi_l(\mathbf{m}(\varsigma^{-1}))) \prod_{\mathfrak{l} | \mathfrak{N}^+} \mathcal{B}_{S_{\theta_l}}^{\Lambda_l}(\mathrm{Id}). \end{aligned}$$

Taking Remark 2.4 into account, we deduce from Theorem 4.2 that

$$\mathcal{B}_{S_{\theta_l}}^{\Lambda_l}(\mathrm{Id}) = |\mathfrak{d}_{K_l}|_{K_l}^{1/2} |\mathfrak{d}_{F_l}|^{-1/2}$$

if l and $\mathfrak{p}\mathfrak{N}$ are coprime. If l divides \mathfrak{N}^- , then by Corollary 5.3 and (5.1)

$$\mathcal{B}_S^{\Lambda_l}(\mathrm{Id}) = \frac{|\mathfrak{d}_{K_l}|_{K_l}^{1/2}}{|\mathfrak{d}_{F_l}|^{1/2}} q_l^3 (1 - q_l^{-2}) \times \begin{cases} 1 & \text{if } K_l/F_l \text{ is unramified,} \\ 1 - \epsilon_l(\pi) \Lambda(\mathfrak{l}_K) & \text{if } \mathfrak{l} = \mathfrak{l}_K^2 \text{ is ramified.} \end{cases}$$

If \mathfrak{N}^+ is divisible by l , then Proposition 4.10 gives

$$\mathcal{B}_S^{\Lambda_l}(\mathbf{m}(\varsigma^{-1})) = \epsilon_l(\pi) \Lambda(\varpi_l)^{-1} |\mathfrak{d}_{K_l}|_{K_l}^{1/2} |\mathfrak{d}_{F_l}|^{-1/2} q_l (1 + q_l^{-2}).$$

Since the measure dt_v gives $F_v^\times \backslash K_v^\times$ the volume 2, Proposition 5.8 gives

$$\frac{\mathcal{B}_{S_\theta}^{\Lambda_v}(\phi_v^0, \phi_v^0)}{\mathbf{r}(\phi_v^0, \pi_v(\mathbf{d}(-1))\phi_v^0)} = 2^{4\kappa_v - 2} N_{F_v}^{D_v}(S_\theta)^{(2\kappa_v - 3)/2} e^{4\pi\sqrt{-1}\mathrm{Tr}_{\mathbb{Q}}^D(\mathrm{Si})}$$

for $v \in \Sigma$ in view of $c(\pi_v, \Lambda_v) = \frac{(2\pi)^{2\kappa_v}}{\Gamma(2\kappa_v-1)\pi}$. Proposition 8.5 gives

$$\mathcal{B}_{S_{\theta_p}^{\Lambda_p}}(\pi_{\mathfrak{p}}(\mathbf{m}(\xi^{(n)})e_{\text{ord},\mathfrak{p}}^0) = |\mathfrak{d}_K|_K^{1/2} |\mathfrak{d}_F|^{-1/2} L(1, \tau_{K_{\mathfrak{p}}/F_{\mathfrak{p}}})^2 e(\pi_{\mathfrak{p}}, \Lambda_{\mathfrak{p}})^2 \alpha_{\mathfrak{p}}^{-2n} q^{-4n}.$$

Upon combining these calculations we obtain Theorem 9.5. \square

10. THETA ELEMENTS AND p -ADIC L -FUNCTIONS

10.1. **Quaternionic modular forms.** Let

$$D_-^+ := \{S \in D_-(F) \mid J_1 S > 0 \text{ for every } v \in \Sigma\}.$$

Given $B \in D_-^+$, we define a function $W_B^{(\kappa)} : \text{GU}_2^D(F_\infty)^\circ \rightarrow \mathbb{C}$ by

$$(10.1) \quad W_B^{(\kappa)}(h) = e^{2\pi\sqrt{-1}\text{Tr}_{\mathbb{Q}}^D(Bh(\mathbf{i}))} J_\kappa(h, \mathbf{i})^{-1}.$$

Definition 10.1 (adèlic quaternionic cusp forms). Let \mathcal{K} be an open compact subgroup of $\text{GU}_2^D(\widehat{F})$. A quaternionic cusp form of weight κ and level \mathcal{K} is a \mathbb{C} -valued function ϕ on $\text{GU}_2^D(F) \backslash \text{GU}_2^D(\mathbb{A}_F) / \mathcal{K}$ which satisfies

$$\phi(zhk) = \phi(h) J_\kappa(k, \mathbf{i})^{-1}$$

for every $k \in \text{U}_2^\Sigma$ and $z \in \mathbb{A}^\times$ and admits a Fourier expansion of the form

$$\phi(h) = \sum_{B \in D_-^+} \mathbf{W}_B(\phi, h) = \sum_{B \in D_-^+} \mathbf{w}_B(\phi, h_{\mathbf{f}}) W_B^{(\kappa)}(h_\infty)$$

for $h \in \text{GU}_2^D(F_\infty)^\circ \text{GU}_2^D(\widehat{F})$, where $\mathbf{w}_B(\phi, -) : h_{\mathbf{f}} \mapsto \mathbf{w}_B(\phi, h_{\mathbf{f}})$ is a locally constant \mathbb{C} -valued function on $\text{GU}_2^D(\widehat{F})$.

We denote the space of adèlic quaternionic cusp forms of weight κ and level \mathcal{K} by $\mathcal{A}_\kappa^0(\mathcal{K})$. The space $\mathcal{A}_\kappa^0(\mathcal{K})$ is contained in the subspace of $\mathcal{A}_{\text{cusp}}(\text{PGU}_2^D)$ which consists of right \mathcal{K} -invariant cuspidal automorphic forms with scalar K -type $k \mapsto J_\kappa(k, \mathbf{i})^{-1}$ (cf. [2]). The finite adèle group $\text{PGU}_2^D(\widehat{F})$ acts on the space $\mathcal{A}_\kappa^0 = \bigcup_{\mathcal{K}} \mathcal{A}_\kappa^0(\mathcal{K})$ by right translation. If an irreducible cuspidal automorphic representation of $\text{PGU}_2^D(\mathbb{A}_F)$ has the lowest weight representation with minimal K -type $k \mapsto J_\kappa(k, \mathbf{i})^{\pm 1}$ as its archimedean component, then its non-archimedean component appears in the decomposition of \mathcal{A}_κ^0 .

10.2. **Theta elements.** Let (π, V) be an irreducible cuspidal automorphic representation of $\text{PGU}_2^D(\mathbb{A}_F)$ such that $\pi_v \simeq D_{\kappa_v}^{(2)}$ for $v \in \Sigma$, such that π_l is generic for every finite prime l and such that $\dim V_\kappa(\pi, \mathfrak{N}) = 1$. Fix a basis vector $\phi_\pi = \otimes_v \phi_v^0 \in V_\kappa(\pi, \mathfrak{N})$. Let $\mathcal{O}_{\mathfrak{p}^n} = \mathfrak{o}_F + \mathfrak{p}^n \mathfrak{o}_K$ be the order of \mathfrak{o}_K of conductor \mathfrak{p}^n and $\mathcal{G}_n = K^\times \backslash \widehat{K}^\times / \widehat{\mathcal{O}}_{\mathfrak{p}^n}^\times$ its Picard group. We identify \mathcal{G}_n with the Galois group of the ring class field $K_{\mathfrak{p}^n}$ of conductor \mathfrak{p}^n over K via geometrically normalized reciprocity law. Denote by $[\cdot]_n : \widehat{K}^\times \rightarrow \mathcal{G}_n$ the natural projection map. Define

$$x_n : \widehat{K}^\times \rightarrow \text{GU}_2^D(\widehat{F}), \quad x_n(t) = \mathbf{t}(t) \zeta^{(n)}.$$

Definition 10.2. Let $\alpha_{\mathcal{Q}} = q_{\mathfrak{p}}^{\kappa-1} \alpha_{\mathfrak{p}}^{-1}$. Define the n th theta element by

$$\Theta_n^S(\phi_{\pi}) = \alpha_{\mathcal{Q}}^{-n} \sum_{[a]_n \in \mathcal{G}_n} q_{\mathfrak{p}}^{\kappa n} \mathbf{w}_S(e_{\text{ord}, \mathfrak{p}}^0 \phi_{\pi}, x_n(a)) [a]_n \in \mathbb{C}[\mathcal{G}_n].$$

The sequence $\{\Theta_n^S(\phi_{\pi})\}_n$ satisfies the following compatibility condition:

Lemma 10.3. Let $\Pi_n^{n+1} : \mathcal{G}_{n+1} \rightarrow \mathcal{G}_n$ be the natural quotient map. Then

$$\Pi_n^{n+1}(\Theta_{n+1}^S(\phi_{\pi})) = \Theta_n^S(\phi_{\pi}).$$

Proof. For $n' > n$, let $K_n^{n'}$ be the kernel of the quotient map $\mathcal{G}_{n'} \rightarrow \mathcal{G}_n$. Recall that

$$(10.2) \quad U_{\mathfrak{p}}^{\mathcal{Q}} \phi = \sum_{x \in \mathfrak{o}_F/\mathfrak{p}} \sum_{y \in \mathfrak{o}_F/\mathfrak{p}} \sum_{z \in \mathfrak{o}_F/\mathfrak{p}^2} \pi_{\mathfrak{p}} \left(\mathbf{n} \left(\begin{pmatrix} z & y \\ y & 0 \end{pmatrix} \right) \mathbf{m} \left(\begin{pmatrix} \varpi_{\mathfrak{p}} & x \\ 0 & 1 \end{pmatrix} \right) \right) \phi.$$

Since $U_{\mathfrak{p}}^{\mathcal{Q}} \phi_{\pi}^{\dagger} = q_{\mathfrak{p}}^2 \alpha_{\mathfrak{p}}^{-1} \phi_{\pi}^{\dagger}$, we have

$$\sum_{x \in \mathfrak{o}_F/\mathfrak{p}} \mathbf{w}_S \left(\phi_{\pi}^{\dagger}, \mathbf{t}(a) \zeta^{(n)} \mathbf{m} \left(\begin{pmatrix} \varpi_{\mathfrak{p}} & x \\ 0 & 1 \end{pmatrix} \right) \right) = q_{\mathfrak{p}}^{-1} \alpha_{\mathfrak{p}}^{-1} \mathbf{w}_S \left(\phi_{\pi}^{\dagger}, \mathbf{t}(a) \zeta^{(n)} \right).$$

Observing that

$$(\zeta_{\mathfrak{p}}^{(n)})^{-1} (1 + \varpi_{\mathfrak{p}}^n x \theta) \zeta_{\mathfrak{p}}^{(n)} = \begin{cases} \begin{pmatrix} 1 + \varpi_{\mathfrak{p}}^n x \theta & -x \\ 0 & 1 + \varpi_{\mathfrak{p}}^n x \bar{\theta} \end{pmatrix} & \text{if } \mathfrak{p} \text{ splits in } K, \\ \begin{pmatrix} 1 & -x \\ \varpi_{\mathfrak{p}}^n x \mathbf{N}(\theta) & 1 + \varpi_{\mathfrak{p}}^n x \mathbf{Tr}(\theta) \end{pmatrix} & \text{otherwise,} \end{cases}$$

we get

$$\sum_{x \in \mathfrak{o}_F/\mathfrak{p}} \mathbf{w}_S \left(\phi_{\pi}^{\dagger}, \mathbf{t}(a(1 + \varpi_{\mathfrak{p}}^n x \theta)) \zeta^{(n+1)} \right) = q_{\mathfrak{p}}^{-1} \alpha_{\mathfrak{p}}^{-1} \mathbf{w}_S \left(\phi_{\pi}^{\dagger}, \mathbf{t}(a) \zeta^{(n)} \right).$$

The left hand side is $\sum_{u \in K_n^{n+1}} \mathbf{w}_S \left(\phi_{\pi}^{\dagger}, x_{n+1}(au) \right)$ in view of the description

$$K_n^{n'} = [\hat{\mathcal{O}}_{\mathfrak{p}^n}^{\times}]_{n'} = \{[1 + \varpi_{\mathfrak{p}}^n x \theta]_{n'} \mid x \in \mathfrak{o}_F/\mathfrak{p}^{n'-n}\}.$$

The proof is complete by Definition 10.2. \square

Put $\mathcal{G}_{\infty} = \varprojlim_n \mathcal{G}_n$. Lemma 10.3 enables us to define

$$\Theta^S(\phi_{\pi}) := \{\Theta_n^S(\phi_{\pi})\}_n \in \varprojlim_n \mathbb{C}[\mathcal{G}_n].$$

Assuming that $c(\Lambda_{\mathfrak{l}}) = 0$ for $\mathfrak{l} \neq \mathfrak{p}$, we will write $c(\Lambda) = c(\Lambda_{\mathfrak{p}})$. When $n \geq c(\Lambda)$, we can view Λ as a character of \mathcal{G}_n and extend it linearly to a function $\Lambda : \mathbb{C}[\mathcal{G}_n] \rightarrow \mathbb{C}$. Let W_K be the group of roots of unity in K and w_K its order. Put $Q_K = [\mathfrak{o}_K^{\times} : W_K \mathfrak{o}_F^{\times}] \in \{1, 2\}$.

Proposition 10.4. *Assume that $F_l = \mathbb{Q}_2$ if $l \neq \mathfrak{p}$ and 2 is divisible by l . If $n \geq 1$ and $n \geq c(\Lambda)$, then*

$$\begin{aligned} \frac{\Lambda(\Theta_n^S(\phi_\pi))^2}{\langle \phi_\pi, \pi(\tau_\infty)\phi_\pi \rangle_{\mathcal{K}_D(\mathfrak{N})}} &= Q_K^2 w_K^2 \frac{\Delta_F \Delta_K^{1/2} N_F^D(4S)^\kappa}{2^{4d+2+\ell(\pi)} N_{\mathbb{Q}}^D(S)^{3/2}} \xi_F(2) \xi_F(4) \frac{\Lambda(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \Lambda)}{N\Lambda(1, \pi, \text{ad})} \\ &\times \frac{e(\pi_{\mathfrak{p}}, \Lambda_{\mathfrak{p}})^2 \epsilon_{\mathfrak{N}^+}(\pi)}{[\mathfrak{o}_K : \mathfrak{o}_F + \theta \mathfrak{o}_F]^3 \Lambda(\mathfrak{N}_0^+)} \prod_{\substack{\mathfrak{l} | \mathfrak{N}^- \cap \mathfrak{D}_F^K, \\ \mathfrak{l} = \mathfrak{l}_K^2}} (1 - \epsilon_{\mathfrak{l}}(\pi) \Lambda(\mathfrak{l}_K)). \end{aligned}$$

Proof. We may assume that $n = \max\{1, c(\Lambda)\}$ by Lemma 10.3. Denote by $\text{vol}(\hat{\mathcal{O}}_{\mathfrak{p}^n}^\times)$ the volume of the image of $K_\infty^\times \hat{\mathcal{O}}_{\mathfrak{p}^n}^\times$ in $K^\times \mathbb{A}_F^\times \backslash \mathbb{A}_K^\times$ with respect to the measure dt . Remark 9.2(1) together with the class number formula gives

$$\text{vol}(\hat{\mathcal{O}}_{\mathfrak{p}^n}^\times) = \text{vol}(\hat{\mathfrak{o}}_K^\times) L(1, \tau_{K_{\mathfrak{p}}/F_{\mathfrak{p}}}) q_{\mathfrak{p}}^{-n} = 2^{d+1} Q_K^{-1} w_K^{-1} \sqrt{\Delta_F \Delta_K^{-1}} L(1, \tau_{K_{\mathfrak{p}}/F_{\mathfrak{p}}}) q_{\mathfrak{p}}^{-n}.$$

Since $W_S^{(\kappa)}(\mathfrak{t}(t)g) = W_S^{(\kappa)}(g)$ for $t \in K_\infty^\times$ by (10.1),

$$\begin{aligned} B_S^\Lambda(\phi_\pi^\dagger, \zeta^{(n)}) &= \frac{(\det S)^{\kappa/2}}{e^{2\pi\sqrt{-1}\text{Tr}_{\mathbb{Q}}^D(Si)}} \int_{K^\times \hat{F}^\times \backslash \hat{K}^\times} \mathbf{w}_S(\phi_\pi^\dagger, \mathfrak{t}(t)\zeta^{(n)}) \Lambda(t)^{-1} dt \\ &= e^{-2\pi\sqrt{-1}\text{Tr}_{\mathbb{Q}}^D(Si)} \text{vol}(\hat{\mathcal{O}}_{\mathfrak{p}^n}^\times) q_{\mathfrak{p}}^{-n} \alpha_{\mathfrak{p}}^{-n} \Lambda(\Theta_n^S(\phi_\pi)). \end{aligned}$$

Theorem 9.5 gives the declared formula. \square

10.3. Classical quaternionic cusp forms. Hereafter let $F = \mathbb{Q}$. Thus

$$\mathfrak{N} = N = N^+ N^-, \quad K = \mathbb{Q}(\sqrt{-\Delta_K}), \quad \mathcal{K}_D(N) = \mathcal{K}_D(N) \cap \text{U}_2^D(\mathbb{Q}).$$

It is important to note that

$$(10.3) \quad D^\times(\mathbb{A}) = D^\times \cdot D_\infty^{\times \circ} \hat{R}^\times, \quad \text{GU}_2^D(\mathbb{A}) = \text{GU}_2^D(\mathbb{Q}) \text{GU}_2^D(\mathbb{R})^\circ \mathcal{K}_D(N).$$

We associate to $h_\infty \in \text{U}_2^D(\mathbb{R})$ and a function $f : \mathfrak{H}_2^* \rightarrow \mathbb{C}$ another function

$$f|_{\kappa} h_\infty : \mathfrak{H}_2^* \rightarrow \mathbb{C}, \quad f|_{\kappa} h_\infty(Z) = f(h_\infty Z) J_\kappa(h_\infty, Z)^{-1}.$$

Symbolically, we will abbreviate $q^B = e^{2\pi\sqrt{-1}\text{Tr}_{\mathbb{C}}^{D_\infty \otimes \mathbb{C}}(BZ)}$ for $B \in D_-^+$.

Definition 10.5 (classical quaternionic cusp forms). A quaternionic cusp form of weight κ with respect to a discontinuous subgroup $\mathbf{K} \subset \text{U}_2^D(\mathbb{Q})$ is a holomorphic function f on \mathfrak{H}_2^* which satisfies $f|_{\kappa}\gamma = f$ for every $\gamma \in \mathbf{K}$ and admits for every $\beta \in \text{U}_2^D(\mathbb{Q})$ a Fourier expansion of the form

$$f|_{\kappa}\beta(Z) = \sum_{B \in D_-^+} \mathbf{c}_B(f|_{\kappa}\beta) q^B.$$

Let $S_\kappa(\mathbf{K}, \mathbb{C})$ stand for the space of such cusp forms.

Let \mathcal{K} be an open compact subgroup of $\text{GU}_2^D(\hat{\mathbb{Q}})$. Set $\mathbf{K} = \text{U}_2^D(\mathbb{Q}) \cap \mathcal{K}$. If

$$\text{GU}_2^D(\mathbb{A}) = \text{GU}_2^D(\mathbb{Q}) \text{GU}_2^D(\mathbb{R}) \mathcal{K},$$

then we can associate to each $f \in S_\kappa(\mathbb{K}, \mathbb{C})$ a unique $\phi_f \in \mathcal{A}_\kappa^0(\mathcal{K})$ such that

$$f(Z) = \phi_f(h_\infty) J_\kappa(h_\infty, \mathbf{i}) \quad (h_\infty \in \mathrm{GU}_2^D(\mathbb{R})^\circ, h_\infty(\mathbf{i}) = Z).$$

We shall call ϕ_f the adèlic lift of f . By definition $\mathbf{w}_B(\phi_f, e) = \mathbf{c}_B(f)$. Let \mathbb{I}_p be the standard Iwahori subgroup in $\mathrm{GSp}_4(\mathbb{Z}_p)$ in Section 7. Put

$$\mathcal{K}_D(N, p) = \{g \in \mathcal{K}_D(N) \mid g_p \in \mathbb{I}_p\}, \quad \mathbf{K}_D(N, p) = \mathcal{K}_D(N, p) \cap \mathrm{U}_2^D(\mathbb{Q}).$$

Recall that \mathfrak{J}_ℓ is the Iwahori subgroup of $\mathrm{GL}_2(\mathbb{Z}_\ell)$. Let R be an Eichler order of level N^+ in \mathfrak{o}_D . We identify $R_\ell = R \otimes_{\mathbb{Z}} \mathbb{Z}_\ell$ with $\mathrm{M}_2(\mathbb{Z}_\ell)$ or \mathfrak{J}_ℓ via i_ℓ according to whether $\ell \nmid N$ or $\ell \mid N^+$. Put

$$\begin{aligned} R^\perp &= \{x \in D \mid \mathrm{Tr}_{\mathbb{Q}}^D(xy) \in \mathbb{Z} \text{ for all } y \in R\}, \\ \check{R}_- &= \{x \in D_- \mid \mathrm{Tr}_{\mathbb{Q}}^D(xy) \in \mathbb{Z} \text{ for all } y \in R^\perp \cap D_-\}. \end{aligned}$$

Observe that if N^+ is divisible by ℓ , then

$$J_\star^{-1} \mathbf{K}(\ell) J_\star = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, d \in R_\ell, b \in R_\ell^\perp, c \in \ell R_\ell^\perp \right\}.$$

It follows that

$$\mathbf{K}_D(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, d \in R, b \in R^\perp, c \in NR^\perp \right\}.$$

Thus the Fourier coefficients of cusp forms in the spaces $S_\kappa(\mathbf{K}_D(N), \mathbb{C})$ and $S_\kappa(\mathbf{K}_D(N, p), \mathbb{C})$ are indexed by $\check{R}_-^+ = \check{R}_- \cap D_-^+$.

The operators $U_p^{\mathcal{P}}$ and $U_p^{\mathcal{Q}}$ on the space $\mathcal{A}_\kappa^0(\mathcal{K}_D(N, p))$ are defined in Definition 7.1. We define the operators $\mathbf{U}_p^{\mathcal{P}}$ and $\mathbf{U}_p^{\mathcal{Q}}$ on $S_\kappa(\mathbf{K}_D(N, p), \mathbb{C})$ by

$$\begin{aligned} [\mathbf{U}_p^{\mathcal{P}} f](Z) &= p^{\kappa-3} \cdot [U_p^{\mathcal{P}} \phi_f](h_\infty) \cdot J_\kappa(h_\infty, \mathbf{i}), \\ [\mathbf{U}_p^{\mathcal{Q}} f](Z) &= p^{\kappa-3} \cdot [U_p^{\mathcal{Q}} \phi_f](h_\infty) \cdot J_\kappa(h_\infty, \mathbf{i}), \end{aligned}$$

where $f \in S_\kappa(\mathbf{K}_D(N, p), \mathbb{C})$ and $h_\infty \in \mathrm{GU}_2^D(\mathbb{R})^\circ$ with $h_\infty(\mathbf{i}) = Z$.

Proposition 10.6. *Let $f \in S_\kappa(\mathbf{K}_D(N, p), \mathbb{C})$. Then*

$$[\mathbf{U}_p^{\mathcal{P}} f](Z) = \sum_{B \in \check{R}_-^+} \mathbf{c}_{pB}(f) q^{pB}, \quad [\mathbf{U}_p^{\mathcal{Q}} f](Z) = \sum_{x=1}^p \sum_{B \in \check{R}_-^+} \mathbf{c}_{\tilde{\gamma}_x B \gamma_x}(f) q^B,$$

where $\gamma_x \in D^\times$ is such that $\gamma_x \in i_p \left(\begin{pmatrix} p & x \\ 0 & 1 \end{pmatrix} \right) D_\infty^{\times \circ} \widehat{R}^\times$.

Proof. The first formula is easy to prove. We see by (10.2) and (10.1) that

$$\begin{aligned} [U_p^{\mathcal{Q}} \phi](h_\infty) &= \sum_{x=1}^p \sum_{X \in R_- / \gamma_x R_- \tilde{\gamma}_x} \phi \left(\begin{pmatrix} \gamma_x^{-1} & 0 \\ 0 & \tilde{\gamma}_x \end{pmatrix}_\infty \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix}_\infty h_\infty \right) \\ &= p^{-\kappa} \sum_{x=1}^p \sum_{B \in \check{R}_-^+} \sum_{X \in R_- / \gamma_x R_- \tilde{\gamma}_x} \mathbf{c}_B(f) e^{2\pi \sqrt{-1} \mathrm{Tr}_{\mathbb{Q}}^D(B \gamma_x^{-1} (Z+X) \tilde{\gamma}_x^{-1})} \end{aligned}$$

$$= p^{-\kappa} \sum_{x=1}^p \sum_{B \in \check{R}_-^+} \mathbf{c}_B(f) q^{\check{\gamma}_x^{-1} B \gamma_x^{-1}} \sum_{X \in R_- / \gamma_x R_- \check{\gamma}_x} e^{2\pi\sqrt{-1}\mathrm{Tr}_{\mathbb{Q}}^D(\check{\gamma}_x^{-1} B \gamma_x^{-1} X)}$$

where $R^- = R \cap D_-$. Note that

$$\sum_{X \in R_- / \gamma_x R_- \check{\gamma}_x} e^{2\pi\sqrt{-1}\mathrm{Tr}_{\mathbb{Q}}^D(\check{\gamma}_x^{-1} B \gamma_x^{-1} X)} = \begin{cases} \#(R_- / \gamma_x R_- \check{\gamma}_x) & \text{if } B \in \check{\gamma}_x \check{R}_- \gamma_x, \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand,

$$R_- / \gamma_x R_- \check{\gamma}_x \simeq \mathrm{Sym}_2(\mathbb{Z}_p) / \begin{pmatrix} p & 0 \\ x & 1 \end{pmatrix} \mathrm{Sym}_2(\mathbb{Z}_p) \begin{pmatrix} p & x \\ 0 & 1 \end{pmatrix}.$$

We find that $\#(R / \gamma_x R \check{\gamma}_x) = p^3$. \square

Definition 10.7. For each subring $A \subset \mathbb{C}$ the space $S_\kappa(\mathbb{K}, A)$ consists of cusp forms $f \in S_\kappa(\mathbb{K}, \mathbb{C})$ such that $\mathbf{c}_B(f) \in A$ for every $B \in D_-^+$.

The following result follows from Proposition 10.6 immediately.

Corollary 10.8. $\mathbf{U}_p^{\mathcal{Q}}$ and $\mathbf{U}_p^{\mathcal{P}}$ stabilize $S_\kappa(\mathbb{K}_D(N, p), A)$ for any A .

Lemma 10.9. *If $f \in S_\kappa(\mathbb{K}_D(N, p), A)$, then for every $B \in D_-^+$ and $t \in \widehat{K}^\times$*

$$p^{n\kappa} \mathbf{w}_B(\phi_f, x_n(t)) \in A.$$

Proof. Let $\mathcal{R} \subset R$ be the Eichler order of level pN^+ . Given $t \in \widehat{K}^\times$, we use (10.3) to write $t\varsigma_p^{(n)} = \gamma \mathbf{f} u$ with $\gamma \in D^\times(\mathbb{Q})$, $N_{\mathbb{Q}}^D(\gamma) > 0$ and $u \in \widehat{\mathcal{R}}^\times$. Then $\mathbf{t}(\gamma \mathbf{f})^{-1} x_n(t) \mathbf{d}(p^n) \in \mathcal{K}_D(N, p)$. Let $h_\infty \in \mathrm{GU}_2^D(\mathbb{R})^\circ$. Put $Z = h_\infty(\mathbf{i})$. Then

$$\begin{aligned} \phi_f(h_\infty x_n(t)) &= \phi_f \left(\begin{pmatrix} \gamma^{-1} & \\ & p^n \gamma^{-1} \end{pmatrix}_\infty \cdot h_\infty \right) \\ &= \frac{f(p^{-n} \gamma^{-1} Z \gamma)}{J_\kappa \left(\begin{pmatrix} \gamma^{-1} & 0 \\ 0 & p^n \gamma^{-1} \end{pmatrix}, Z \right)} = p^{-n\kappa} f(p^{-n} \gamma^{-1} Z \gamma). \end{aligned}$$

Thus $p^{n\kappa} \mathbf{w}_B(\phi_f, x_n(t)) = \mathbf{w}_{p^n \gamma^{-1} B \gamma}(\phi_f, e) = \mathbf{c}_{p^n \gamma^{-1} B \gamma}(f) \in A$. \square

10.4. Anticyclotomic p -adic L -functions. Let $f \in S_\kappa(\mathbb{K}_D(N), \mathbb{C})$ be a Hecke eigenform and π an irreducible cuspidal automorphic representation of $\mathrm{PGU}_2^D(\mathbb{A})$ generated by the associated adèlic lift $\phi_\pi := \phi_f \in \mathcal{A}_\kappa^0(\mathcal{K}_D(N))$. Denote the ring of integers of the Hecke field F_π of π by \mathfrak{o}_π . We may assume that f belongs to $S_\kappa(\mathbb{K}_D(N), \mathfrak{o}_\pi)$ (cf. [12, Proposition 1.8 on p. 146] or [24]). The Hecke field F_π is a totally real number field for π is self-dual.

Define an automorphic of $\mathrm{GU}_2(D)$ by $g^c = \mathbf{d}(-1)g\mathbf{d}(-1)$. Since

$$g_\infty^c(\mathbf{i}) = -\overline{g_\infty(\mathbf{i})}, \quad J_\kappa(g_\infty^c, \mathbf{i}) = \overline{J_\kappa(g_\infty, \mathbf{i})}$$

for $g_\infty \in \mathrm{GU}_2(\mathbb{R})^\circ$, the cusp form $\overline{\pi(\tau_\infty)\phi_\pi}$ corresponds to $\sum_{B \in D_-^+} \mathbf{c}_B(f) q^B$, and hence $\overline{\phi_\pi} = \pi(\tau_\infty)\phi_\pi$. We may further assume that not all of the Fourier coefficients of f are zero modulo the maximal ideal of \mathfrak{o}_π with respect to ι_p .

Let $\{\alpha_p, \alpha_p^{-1}\gamma_p^{-2}, \gamma_p\}$ be the Satake parameters of π_p . Put

$$\begin{aligned}\alpha_{\mathcal{P}} &= p^{\kappa-3/2}\gamma_p, & \beta_{\mathcal{P}} &= p^{\kappa-3/2}\gamma_p^{-1}\alpha_p^{-1}, \\ \alpha_{\mathcal{Q}} &= p^{2-\kappa}\alpha_{\mathcal{P}}\beta_{\mathcal{P}} = p^{\kappa-1}\alpha_p^{-1}, & \beta_{\mathcal{Q}} &= p^{\kappa-1}\alpha_{\mathcal{P}}\beta_{\mathcal{P}}^{-1} = p^{\kappa-1}\alpha_p\gamma_p^2.\end{aligned}$$

Definition 10.10. Let

$$f^\ddagger := \alpha_{\mathcal{P}}^{-3}\alpha_{\mathcal{Q}}^{-1} \cdot (\mathbf{U}_p^{\mathcal{Q}} - \beta_{\mathcal{Q}})(\mathbf{U}_p^{\mathcal{P}} - p^{2\kappa-3}\alpha_{\mathcal{P}}^{-1})(\mathbf{U}_p^{\mathcal{P}} - p^{2\kappa-3}\beta_{\mathcal{P}}^{-1})(\mathbf{U}_p^{\mathcal{P}} - \beta_{\mathcal{P}})f.$$

Let $\text{ord}_p : \overline{\mathbb{Q}}_p^\times \rightarrow \mathbb{Q}$ denote the normalized additive valuation. From now on we assume one of the parameters of π_p to satisfy

$$(\mathcal{Q}\text{-ord}) \quad \text{ord}_p \iota_p(\alpha_{\mathcal{Q}}) = 0.$$

It is convenient to suppose that another parameter satisfies

$$(\mathcal{P}\text{-ord}) \quad \text{ord}_p \iota_p(\alpha_{\mathcal{P}}) = 0.$$

Remark 10.11. (1) The eigenvalues of $\mathbf{U}_p^{\mathcal{Q}}$ are $\alpha_{\mathcal{Q}}, \beta_{\mathcal{Q}}, p^{2\kappa-2}\alpha_{\mathcal{Q}}^{-1}, p^{2\kappa-2}\beta_{\mathcal{Q}}^{-1}$ and those of $\mathbf{U}_p^{\mathcal{P}}$ are $\alpha_{\mathcal{P}}, \beta_{\mathcal{P}}, p^{2\kappa-3}\alpha_{\mathcal{P}}^{-1}, p^{2\kappa-3}\beta_{\mathcal{P}}^{-1}$ by the proof of Proposition 7.4.

(2) The eigenvalue of f^\ddagger for $\mathbf{U}_p^{\mathcal{Q}}$ is a p -adic unit if and only if $(\mathcal{Q}\text{-ord})$ holds.

(3) The eigenvalue of f^\ddagger for $\mathbf{U}_p^{\mathcal{P}}$ is a p -adic unit if and only if $(\mathcal{P}\text{-ord})$ holds.

Lemma 10.12. *If π_p satisfies $(\mathcal{Q}\text{-ord})$ and A contains \mathfrak{o}_π and eigenvalues of $\mathbf{U}_p^{\mathcal{P}}$ and $\mathbf{U}_p^{\mathcal{Q}}$, then $\alpha_{\mathcal{P}}^3 \cdot f^\ddagger \in S_\kappa(\mathbf{K}_D(N, p), A)$.*

Proof. Remark 10.11(1) and Corollary 10.8 imply that

$$\beta_{\mathcal{Q}}, \beta_{\mathcal{P}}, p^{2\kappa-3}\alpha_{\mathcal{P}}^{-1}, p^{2\kappa-3}\beta_{\mathcal{P}}^{-1} \in \overline{\mathbb{Z}},$$

and the lemma follows from $(\mathcal{Q}\text{-ord})$ and Definition 10.10. \square

Let Γ^- be the maximal \mathbb{Z}_p -free quotient group of \mathcal{G}_∞ and Δ the torsion subgroup of \mathcal{G}_∞ . We have an exact sequence

$$1 \rightarrow \Delta \rightarrow \mathcal{G}_\infty \rightarrow \Gamma^- \rightarrow 1.$$

Fix a noncanonical isomorphism $\mathcal{G}_\infty \simeq \Delta \times \Gamma^-$ once and for all. If $n \geq 1$, then the map $\Delta \rightarrow \mathcal{G}_\infty \rightarrow \mathcal{G}_n$ is injective and hence

$$\mathcal{G}_n \simeq \Delta \times \Gamma_n^-, \quad \Gamma^- \twoheadrightarrow \Gamma_n^- = \mathcal{G}_n/\Delta.$$

Let $\chi : \Delta \rightarrow \overline{\mathbb{Q}}^\times$ be a branch character. Define the χ -branch of $\Theta_n^S(\phi_\pi)$ by

$$\Theta_n^S(\phi_\pi, \chi) = \chi(\Theta_n^S(\phi_\pi)) \in \mathbb{C}[\Gamma_n^-].$$

Enlarge \mathfrak{o}_π to a ring A so that A contains values of χ and eigenvalues of $\mathbf{U}_p^{\mathcal{P}}$ and $\mathbf{U}_p^{\mathcal{Q}}$. By Lemma 10.9, $\Theta_n^S(\phi_\pi, \chi)$ belongs to $A[\Gamma_n^-]$, and hence

$$\Theta^S(\phi_\pi, \chi) := \varprojlim_n \Theta_n^S(\phi_\pi, \chi) \in A[[\Gamma_\infty^-]].$$

Definition 10.13 (periods). Recall that we have normalized $f \in S_\kappa(\mathbf{K}_D(N), \mathfrak{o}_\pi)$ so that not all the Fourier coefficients vanish modulo the maximal ideal of the completion of \mathfrak{o}_π with respect to ι_p . Define a period Ω_{π, N^-} of π by

$$\Omega_{\pi, N^-} := \Lambda(1, \pi, \text{ad}) / \langle f, f \rangle_{\mathbf{K}_D(N)},$$

where we define the Petersson norm of f by

$$\langle f, f \rangle_{\mathbf{K}_D(N)} := \int_{\mathbf{K}_D(N) \backslash \mathfrak{H}_2^*} |f(Z)|^2 (\det Y)^{\kappa-3} dX dY.$$

Proposition 10.14. *Let $N = N^+ N^-$ be a square-free integer. Then*

$$\text{vol}(\mathbf{K}_D(N) \backslash \mathfrak{H}_2^*) = 2\xi_{\mathbb{Q}}(2)\xi_{\mathbb{Q}}(4) \prod_{q|N^+} (q^2 + 1) \prod_{\ell|N^-} (\ell^2 - 1).$$

Let $f \in S_\kappa(\mathbf{K}_D(N), \mathbb{C})$. Then

$$\langle f, f \rangle_{\mathbf{K}_D(N)} = \xi_{\mathbb{Q}}(2)\xi_{\mathbb{Q}}(4) \langle \phi_\pi, \pi(\tau_\infty)\phi_\pi \rangle_{\mathcal{K}_D(N)}.$$

Proof. Recall that a motive M of Artin-Tate type is attached to U_2^D in Section 1 of [19] and a canonical Haar measure $|\omega_v|$ on $U_2^D(\mathbb{Q}_v)$ is defined in Section 4 of [19]. For each rational prime q , let μ_q be the Haar measure $L_q(M^\vee(1))|\omega_q|$ on $U_2^D(\mathbb{Q}_q)$. Let μ_∞ be Euler-Poincaré measure on $U_2^D(\mathbb{R})$. Then $\mu = \otimes_v \mu_v$ defines a Haar measure on $U_2^D(\mathbb{A})$. Since the Tamagawa number of U_2^D is 1, we have

$$(10.4) \quad \int_{U_2^D(\mathbb{Q}) \backslash U_2^D(\mathbb{A})} \mu = L_\infty(M) / c(\text{Sp}_4(\mathbb{R}))$$

by Theorem 9.9 of [19], where $c(\text{Sp}_4(\mathbb{R}))$ is a cohomological invariant attached to the real symplectic group of rank 2.

Let $\mathbb{H} = D^2$ be a left D -vector space with Hermitian form

$$\langle (x, y), (x', y') \rangle = x\bar{y}' + y\bar{x}'.$$

Let $L = \mathfrak{o}_D \oplus \check{\mathfrak{o}}_D$ be a maximal lattice in $\mathbb{H}(\mathbb{Q})$, where

$$\check{\mathfrak{o}}_D = \{x \in D(\mathbb{Q}) \mid \text{Tr}_{\mathbb{Q}}^D(xy) \in \mathfrak{o}_D \text{ for every } y \in \mathfrak{o}_D\}.$$

Put $L_q = L \otimes_{\mathbb{Z}} \mathbb{Z}_q$. Define an open compact subgroup $\mathcal{K}(L)$ of $U_2^D(\widehat{\mathbb{Q}})$ by

$$\mathcal{K}(L) = \prod_q \mathcal{K}(L_q), \quad \mathcal{K}(L_q) = \{g \in \text{GU}_2^D(\mathbb{Q}_q) \mid L_q g = L_q\}.$$

Then $\mathcal{K}(L) \simeq \mathcal{K}_D(N^-)$. By the strong approximation property of U_2^D

$$U_2(\mathbb{Q}) \backslash U_2^D(\mathbb{A}) \simeq (\mathbf{K} \backslash \mathfrak{H}_2^*) \times U_2^{\mathbb{Z}} \times \mathcal{K}$$

for any open compact subgroup $\mathcal{K} = \prod_q \mathcal{K}_q$ of $U_2^D(\widehat{\mathbb{Q}})$, where we put $\mathbf{K} = \mathcal{K} \cap U_2^D(\mathbb{Q})$. Now we see from (10.4) that

$$\text{vol}(\mathbf{K} \backslash \mathfrak{H}_2^*) \text{vol}(U_2^{\mathbb{Z}}) \prod_q \mu_q(\mathcal{K}_q) = L_\infty(M) / c(\text{Sp}_4(\mathbb{R})).$$

Taking Lemma 3.3.3 of [31] into account, we get

$$\frac{\text{vol}(\mathbb{K}_D(N) \backslash \mathfrak{H}_2^*)}{\text{vol}(\mathbb{K}_D(N^-) \backslash \mathfrak{H}_2^*)} = \prod_{q|N^+} (q^2 + 1).$$

Proposition 9.3 of [17] says that $\mu_q(\mathcal{K}(L_q)) = 1$ or $q^2 - 1$ according to whether D is split over \mathbb{Q}_q or not. It is well-known that

$$\text{vol}(\text{Sp}_4(\mathbb{Z}) \backslash \mathfrak{H}_2) = 2\xi_{\mathbb{Q}}(2)\xi_{\mathbb{Q}}(4).$$

Combining these results, we obtain the first equality. Proposition 3.1 of [10] combined with this equality and Definition 9.3 gives the second identity. \square

Take θ so that $\mathfrak{o}_K = \mathbb{Z} + \mathbb{Z}\theta$. Recall the decomposition $N^+ \mathfrak{o}_K = \mathfrak{N}_0^+ \overline{\mathfrak{N}_0^+}$.

Theorem 10.15. *Let A be a subring of $\overline{\mathbb{Q}}_p$ which contains \mathfrak{o}_π , values of χ and eigenvalues of \mathbf{U}_p^P and \mathbf{U}_p^Q . If π_p satisfies (Q -ord), then*

$$\alpha_p^3 \cdot \Theta^S(\phi_\pi, \chi) \in A[\mathcal{G}_\infty].$$

Let $\hat{\nu} : \Gamma^- \rightarrow \overline{\mathbb{Q}}_p^\times$ be a p -adic character of finite order. Then

$$\begin{aligned} \frac{\hat{\nu}(\Theta^S(\phi_\pi, \chi))^2}{\langle f, f \rangle_{\mathbb{K}_D(N)}} &= w_K^2 \cdot 2^{2\kappa-3} \Delta_K^{\kappa-1} e(\pi_p, \chi_p \nu_p)^2 \frac{\Lambda(\frac{1}{2}, \text{Spn}(\pi)_K \otimes \chi \nu)}{N 2^{s_\pi} \Lambda(1, \pi, \text{ad})} \\ &\quad \times \epsilon_{N^+}(\pi)(\chi \nu)(\mathfrak{N}_0^+)^{-1} \prod_{\ell|(N^-, \Delta_K), \ell=\ell_K^2} (1 - \epsilon_\ell(\pi)(\chi \nu)(\ell_K)). \end{aligned}$$

Proof. By Definitions 7.2 and 10.10, $\alpha_p^3 \cdot e_{\text{ord}, p}^0 \phi_\pi$ is the adèlic lift of $\alpha_p^3 \cdot f^\ddagger$. In view of Lemmas 10.9 and 10.12, we conclude that

$$\alpha_p^3 \cdot p^{n\kappa} \mathbf{w}_S(e_{\text{ord}, p}^0 \phi_\pi, x_n(t)) \in A$$

for every $t \in \widehat{K}^\times$ and nonnegative integers n . Since $F = \mathbb{Q}$, we have $Q_K = 1$. We have $\det S = \frac{\Delta_K}{4}$ for our choice of θ . We finally get the stated formula by Propositions 10.4 and 10.14. \square

10.5. Reformulation in terms of optimal embeddings. We explain theta elements in Definition 10.2 agrees with the one given in the introduction. When \mathcal{O} is an order of \mathfrak{o} , an embedding $\iota : \mathcal{O} \hookrightarrow R$ is said to be optimal if $\iota(K) \cap R = \iota(\mathcal{O})$. Fix an optimal embedding $\Psi : \mathfrak{o}_K \hookrightarrow R$. Recall that $\mathcal{R} \subset R$ is the Eichler order of level pN^+ . For any positive integer n , write $\varsigma_p^{(n)} \in \gamma_n \widehat{\mathcal{R}}^\times$ for some $\gamma_n \in D^\times$. Then one verifies directly that the embedding $\Psi_n \in \text{Hom}(K, D)$ defined by $\Psi_n(x) = \gamma_n^{-1} \Psi(x) \gamma_n$ is an embedding from \mathcal{O}_{p^n} to \mathcal{R} of conductor p^n , namely an optimal embedding in $\text{Hom}(\mathcal{O}_{p^n}, \mathcal{R})$. For $\sigma \in \text{Gal}(K_{p^n}/K)$, write $\sigma = \text{rec}_K(t)|_{K_{p^n}}$ for some $t \in \widehat{K}^\times$. Write $\Psi_n(t) \in \gamma_n \widehat{\mathcal{R}}^\times$. By definition,

$$x_n(t) = \iota(t) \varsigma_p^{(n)} \in \gamma_n \Psi_n(t) \gamma_n^{-1} \varsigma_p^{(n)} \in \gamma_n \gamma \widehat{\mathcal{R}}^\times.$$

On the other hand, according to the recipe of the Galois action on Ψ_n ,

$$S_{\Psi_n \sigma} = \gamma^{-1} \Psi_n(p^n \sqrt{-\Delta_K/2}) \gamma = p^n (\gamma_n \gamma)^{-1} S_{\Psi} \gamma_n \gamma.$$

By Lemma 10.9, we find that

$$p^{n\kappa} \mathbf{w}_{S_\Psi}(x_n(t), f^\dagger) = \mathbf{c}_{p^n(\gamma_n\gamma)^{-1}S_\Psi\gamma_n\gamma}(f^\dagger) = \mathbf{c}_{S_\Psi^\sigma}(f^\dagger).$$

This shows that the theta element $\Theta_n^S(\phi_\pi, 1)$ with $S = S_\Psi$ agrees with the one described in the introduction.

REFERENCES

- [1] J. Arthur, The endoscopic classification of representations. Orthogonal and symplectic groups, Amer. Math. Soc. Colloq. Publ. **61**, xviii+590pp. Amer. Math. Soc., Providence, RI, 2013.
- [2] M. Asgari and R. Schmidt, Siegel modular forms and representations, *Manuscripta Math.* **104** (2001) 173–200.
- [3] M. Asgari and R. Schmidt, On the adjoint L -function of the p -adic $\mathrm{GSp}(4)$, *J. Number Theory* **128** (2008) 2340–2358.
- [4] M. Bertolini and H. Darmon, Heegner points on Mumford–Tate curves, *Invent. Math.* **126** (1996), no. 3, 413–456.
- [5] S. Böcherer, Bemerkungen über die Dirichletreihen von Koecher und Maass, *Mathematica Gottingensis* **68**, 36 pp., 1986.
- [6] A. Brumer and K. Kramer, Paramodular abelian varieties of odd conductor, *Trans. Amer. Math. Soc.* **366** (2014), 2463–2516.
- [7] D. Bump, S. Friedberg and M. Furusawa, Explicit formulas for the Waldspurger and Bessel models, *Israel J. Math.* **102** (1997) 125–177.
- [8] M. Chida and M.-L. Hsieh, Special values of anticyclotomic L -functions for modular forms, *J. Reine Angew. Math.* (2016) doi:10.1515/crelle-2015-0072
- [9] J. Coates, On p -adic L -functions attached to motives over \mathbb{Q} II, *Boletim da Sociedade Brasileira de Matemática - Bulletin/Brazilian Mathematical Society* **20**(1) (1989) 101–112.
- [10] M. Dickson, A. Pitale, A. Saha and R. Schmidt, Explicit refinements of Böcherer’s conjecture for Siegel modular forms of squarefree level, *J. Math. Soc. Japan* **72**(1) (2020) pp. 251–301 doi: 10.2969/jmsj/78657865
- [11] S. Friedberg and D. Goldberg, On local coefficients for non-generic representations of some classical groups, *Compos. Math.* **116** (1999), no. 2, 133–166.
- [12] G. Faltings and C.-L. Chai, Degeneration of abelian varieties. With an appendix by David Mumford. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* **22**. Springer-Verlag, Berlin, 1990. xii+316 pp.
- [13] M. Furusawa and K. Morimoto, On special Bessel periods and the Gross-Prasad conjecture for $\mathrm{SO}(2n+1) \times \mathrm{SO}(2)$, *Math. Ann.* **368** (2017) 561–586.
- [14] M. Furusawa and K. Morimoto, Refined global Gross-Prasad conjecture on special Bessel periods and Böcherer’s conjecture, to appear in *J. Eur. Math. Soc.*
- [15] M. Furusawa and K. Morimoto, On the Gross-Prasad conjecture with its refinement for $(\mathrm{SO}(5), \mathrm{SO}(2))$ and the generalized Böcherer’s conjecture, preprint. [arXiv: 2205.09503](https://arxiv.org/abs/2205.09503)
- [16] W. T. Gan, B. H. Gross and D. Prasad, Symplectic local root numbers, central critical L -values, and restriction problems, in the representation theory of classical groups, *Astérisque* **346** (2012) 1–109.
- [17] W. T. Gan, J. P. Hanke and J.-K. Yu, On an exact mass formula of Shimura, *Duke Math. J.* **107** (2001), no. 1, 103–133.
- [18] B. H. Gross, Heights and the special values of L -series, *Number theory, Montreal, Que.1985, CMS Conf. Proc.*, **7**, Amer. Math. Soc., Providence, RI, (1987), 115–187.
- [19] B. H. Gross, On the motive of a reductive group, *Invent. Math.* **130** (1997), no. 2, 287–313.

- [20] Y. Hirai, On Eisenstein series on quaternion unitary groups of degree 2, *J. Math. Soc. Japan* **51**(1) (1999) 93–128.
- [21] P.-C. Hung, On nonvanishing mod ℓ of central L -values with anticyclotomic twists for Hilbert modular forms, *J. Number Theory* **173** (2017) 170–209.
- [22] H. Jacquet, Automorphic forms on GL_2 , II, *Lecture Notes in Mathematics* **278**, Springer, Berlin, 1972.
- [23] E. Lapid and Z. Mao, A conjecture on Whittaker-Fourier coefficients of cusp forms, *J. Number Theory* **146** (2015) 448–505.
- [24] K.-W. Lan, Arithmetic compactifications of PEL-type Shimura varieties, *London Mathematical Society Monographs Series*, **36**, Princeton University Press, Princeton, NJ, 2013, xxvi+561.
- [25] G. Laumon, Fonctions zêtas des variétés de Siegel de dimension trois, *Formes automorphes. II. Le cas du groupe $GSp(4)$* , *Astérisque*, **302**, (2005) 1–66.
- [26] Y. Liu, Refined global Gan-Gross-Prasad conjecture for Bessel periods, *J. Reine Angew. Math.* **717** (2016) 133–194.
- [27] M.-L. Hsieh, Special values of anticyclotomic Rankin-Selberg L -functions, *Doc. Math.* **19** (2014) 709–767.
- [28] M. Miyauchi and T. Yamauchi, An explicit computation of p -stabilized vectors, *Journal de Théorie des Nombres de Bordeaux* **26** (2014) 531–558.
- [29] H. Narita, Jacquet–Langlands–Shimizu correspondence for theta lifts to $GSp(2)$ and its inner forms I: An explicit functorial correspondence, *J. Math. Soc. Japan*, **69**(4) (2017) 1443–1474.
- [30] A. Pitale and R. Schmidt, Bessel models for lowest weight representations of $GSp(4, \mathbb{R})$, *Int. Math. Res. Not. IMRN* 2009, no. **7**, 1159–1212.
- [31] B. Roberts and R. Schmidt, *Local newforms for $GSp(4)$* , *Lecture Notes in Mathematics*, 1918. Springer, Berlin, 2007. viii+307 pp.
- [32] P. Sally and M. Tadić, Induced representations and classifications for $GSp(2, F)$ and $Sp(2, F)$, *Bull. Soc. Math. France*, **121**, Mem. **52** (1993) 75–133.
- [33] R. Schmidt, Iwahori-spherical representations of $GSp(4)$ and Siegel modular forms of degree 2 with square-free level, *J. Math. Soc. Japan*, **57**(1) (2005) 259–293.
- [34] G. Shimura, Confluent hypergeometric functions on tube domains, *Math. Ann.* **260** (1982) 269–302.
- [35] E. Urban, Selmer groups and the Eisenstein-Klingen ideal, *Duke Math. J.*, **106**, No. 3 (2001) 485–525.
- [36] E. Urban, Sur les représentations p -adiques associées aux représentations cuspidales de $GSp_{4/\mathbb{Q}}$, *Formes automorphes. II. Le cas du groupe $GSp(4)$* , *Astérisque* **302** (2005) 151–176.
- [37] N. R. Wallach, Lie algebra cohomology and holomorphic continuation of generalized Jacquet integrals, *Representations of Lie Groups*, Kyoto, Hiroshima, 1986, *Adv. Stud. Pure Math.*, vol. **14**, Academic Press, Boston, MA, 1988, pp. 123–151.
- [38] R. Weissauer, Four dimensional Galois representations, *Formes automorphes. II. Le cas du groupe $GSp(4)$* , *Astérisque*, **302** (2005) 67–150.
- [39] M. Rösner and R. Weissauer, Global liftings between inner forms of $GSp(4)$, preprint. [arXiv: 2103.14715](https://arxiv.org/abs/2103.14715).
- [40] S. Yamana, L -functions and theta correspondence for classical groups, *Invent. Math.* **196**(3) (2014) 651–732.
- [41] S. Yamana, Siegel series for skew Hermitian forms over quaternion algebras, *Abh. Math. Sem. Univ. Hamburg* **87** (2017) 43–59.
- [42] H. Yoshida, On generalization of the Shimura-Taniyama conjecture I and II, *Siegel Modular Forms and Abelian Varieties*, *Proceedings of the 4-th Spring Conference on Modular Forms and Related Topics*, 2007, pp. 1–26.

DEPARTMENT OF MATHEMATICS AND NATIONAL CENTER FOR THEORETIC SCIENCE,
NATIONAL TAIWAN UNIVERSITY, NO. 1, SEC. 4, ROOSEVELT ROAD, TAIPEI 10617,
TAIWAN

E-mail address: mlhsieh@math.ntu.edu.tw

OSAKA METROPOLITAN UNIVERSITY, DEPARTMENT OF MATHEMATICS, GRADUATE
SCHOOL OF SCIENCE, 3-3-138 SUGIMOTO, SUMIYOSHI-KU, OSAKA 558-8585 JAPAN

E-mail address: yamana@omu.ac.jp