

INVARIANCE OF GROMOV-WITTEN THEORY UNDER A SIMPLE FLOP

Y. IWAO, Y.-P. LEE, H.-W. LIN, AND C.-L. WANG

ABSTRACT. We show that the generating functions of Gromov–Witten invariants *with ancestors* are invariant under a simple flop, for all genera, after an analytic continuation in the extended Kähler moduli space. This is a sequel to [14].

0. INTRODUCTION

0.1. Statement of the main results. Let X be a smooth complex projective manifold and $\psi : X \rightarrow \bar{X}$ a flopping contraction in the sense of minimal model theory, with $\bar{\psi} : Z \cong \mathbb{P}^r \rightarrow pt$ the restriction map to the extremal contraction. Assume that $N_{Z/X} \cong \mathcal{O}_{\mathbb{P}^r}(-1)^{\oplus(r+1)}$. It was shown in [14] that a simple \mathbb{P}^r flop $f : X \dashrightarrow X'$ exists and the graph closure $[\bar{\Gamma}_f] \in A^*(X \times X')$ induces a correspondence \mathcal{F} which identifies the Chow motives \hat{X} of X and \hat{X}' of X' . Furthermore, the big quantum cohomology rings, or equivalently genus zero Gromov–Witten invariants with 3 or more insertions, are invariant under a simple flop, after an analytic continuation in the extended Kähler moduli space.

The goal of the current paper is to extend the results of [14] to all genera. In the process we discovered the natural framework in the *ancestor potential*

$$\mathcal{A}_X(\bar{t}, s) := \exp \sum_{g=0}^{\infty} \hbar^{g-1} \bar{F}_g^X(\bar{t}, s),$$

which is a formal series in the Novikov variables $\{q^\beta\}_{\beta \in NE(X)}$ defined in the stable range $2g + n \geq 3$. See Section 1 for the definitions.

The main results of this paper are the following theorems.

Theorem 0.1. *The total ancestor potential \mathcal{A}_X (resp. $\mathcal{A}_{X'}$) is analytic in the extremal ray variable q^ℓ (resp. $q^{\ell'}$). They are identified via \mathcal{F} under a simple flop, after an analytic continuation in the extended Kähler moduli space $\omega \in H_{\mathbb{R}}^{1,1}(X) + i(\mathcal{K}_X \cup \mathcal{F}^{-1}\mathcal{K}_{X'})$ via*

$$q^\ell = e^{2\pi i(\omega \cdot \ell)},$$

where \mathcal{K}_X (resp. $\mathcal{K}_{X'}$) is the Kähler cone of X (resp. X').

There are extensive discussions of analytic continuation and the Kähler moduli in Section 3. We note that the *descendent potential* is in general *not*

invariant under \mathcal{F} (c.f. [14], §3). The descendants and ancestors are related via a simple transformation ([10, 8], c.f. Proposition 1.1), but the transformation is in general not compatible with \mathcal{F} . Nevertheless we do have

Theorem 0.2. *For a simple flop f , any generating function of mixed invariant of f -special type*

$$\langle \tau_{k_1, \bar{l}_1} \alpha_1, \dots, \tau_{k_n, \bar{l}_n} \alpha_n \rangle_g,$$

with $2g + n \geq 3$, is invariant under \mathcal{F} up to analytic continuation.

Here a mixed insertion $\tau_{k, \bar{l}} \alpha$ consists of descendants ψ^k and ancestors $\bar{\psi}^l$. Given $f : X \dashrightarrow X'$ with exceptional loci $Z \subset X$ and $Z' \subset X'$, a mixed invariant is of f -special type if for every insertion $\tau_{k, \bar{l}} \alpha$ with $k \geq 1$ we have $\alpha \cdot Z = 0$. Theorem 0.1 follows from an application of Theorem 0.2 when no descendent is present.

0.2. Outline of the contents. Section 1 contains some basic definitions as well as special terminologies in Gromov–Witten theory used in the article. One of the main ingredients of our proof of invariance in the higher genus theory is Givental’s quantization formalism [8] for *semisimple* Frobenius manifolds. This is reviewed in Section 2.

Another main ingredient, in comparing Gromov–Witten theory of X and X' , is the degeneration analysis. We generalize the genus zero results of the degeneration analysis in [14] to ancestor potentials in all genera. The analysis allows us to reduce the proofs of Theorem 0.1 (and 0.2) from flops of X to flops of the local model $\mathbb{P}(N_{Z/X} \oplus \mathcal{O})$.

To keep the main idea clear, we choose to work on local models first in Section 3 and postpone the degeneration analysis till section 4. The local models are semi-Fano toric varieties and localizations had been effectively used to solve the genus zero case. The idea is to utilize Givental’s quantization formalism on the local models to derive the invariance in higher genus, up to analytic continuation, from our results [14] in genus zero. In doing so, the key point is that local models have semisimple quantum cohomology, and we trace the effect of analytic continuation carefully during the process of quantization. The issues of the analyticity of the Frobenius manifolds and the analytic continuation involved in this study is discussed in the beginning Section 3.

The proofs of our main results Theorem 0.1 and 0.2, as well as the degeneration analysis, are presented in section 4.

In section 5 we include some discussions and calculations of the higher genus Gromov–Witten invariants attached to the extremal rays. Similar to the $g = 0$ case, there is also a classical defect occurring at $(g, n, d) = (1, 1, 0)$

$$-\frac{1}{24} [(c_{\text{top}-1}(X) \cdot \alpha)_X - (c_{\text{top}-1}(X') \cdot \mathcal{F}\alpha)_{X'}].$$

Our explicit formula in Theorem 6.12 for the $g = 1$ invariants attached to the extremal ray is seen to give quantum corrections to it.

The calculation of the explicit formula in genus one requires some elementary combinatorics, and is included in the appendix.

0.3. Some remarks on the crepant resolution conjecture. A morphism $\psi : X \rightarrow \bar{X}$ is called a *crepant resolution*, if X is smooth and \bar{X} is \mathbb{Q} -Gorenstein (e.g. an orbifold) such that $\psi^*K_{\bar{X}} = K_X$. In the case \bar{X} is an orbifold, there is a well-defined orbifold Gromov–Witten theory due to Chen–Ruan. The *crepant resolution conjecture* asserts a close relation between the Gromov–Witten theory of X and that of \bar{X} .

Crepant resolution conjecture, as formulated in [3], still uses descendent potentials rather than the ancestor potentials, as advocated in [11]. Yet ancestors often enjoy better properties than the corresponding descendents, as exploited by Getzler [6].

Since different crepant resolutions are related by a K -equivalent transformation, e.g. a flop, the conjecture must be consistent with a transformation under a flop. Although the descendent potentials can be obtained from ancestor potentials via a simple transformation, this very transformation actually spoils the invariance under \mathcal{F} . The insistence in the descendents may introduce unnecessary complication in the formulation of the conjecture. This is especially relevant in the stronger form of the conjecture when the orbifolds satisfy the Hard Lefschetz conditions.

Our result suggests that a more natural framework to study crepant resolution conjecture is to use ancestors rather than descendents. We leave the interested reader to consult [3] and references therein.

0.4. Acknowledgements. Part of this work was done during the second author’s visit to the NCU Center for Mathematics and Theoretic Physics (CMTP), Jhongli, Taiwan in November 2007. He is grateful to the Mathematics Department of National Central University for the hospitality during his stay.

1. DESCENDENT AND ANCESTOR POTENTIALS

1.1. The ancestor potential. For the stable range $2g + m \geq 3$, let

$$\pi := \text{ft} \circ \text{st} : \bar{M}_{g,m+1}(X, \beta) \rightarrow \bar{M}_{g,m}$$

be the composition of the *stabilization morphism* $\text{st} : \bar{M}_{g,m+1}(X, \beta) \rightarrow \bar{M}_{g,m+1}$ defined by forgetting the map and the *forgetful morphism* $\text{ft} : \bar{M}_{g,m+1} \rightarrow \bar{M}_{g,m}$ defined by forgetting the last l points. The *gravitational ancestors* are defined to be

$$(1.1) \quad \bar{\psi}_i := \pi^* \psi_i$$

for $i = 1, \dots, m$.

Let $\{T_\mu\}$ be a basis of $H^*(X, \mathbb{Q})$. Denote $\bar{t} = \sum_{\mu,k} \bar{t}_k^\mu \bar{\psi}^k T_\mu$, $s = \sum_\mu s^\mu T_\mu$, and let

$$\begin{aligned} \bar{F}_g^X(\bar{t}, s) &= \sum_{m,l,\beta} \frac{q^\beta}{m!l!} \langle \bar{F}^m, s^l \rangle_{g,m+l,\beta} \\ &= \sum_{m,l,\beta} \frac{q^\beta}{m!l!} \int_{[\bar{M}_{g,m+l}(X,\beta)]^{\text{vir}}} \prod_{i=1}^m \sum_{k,\mu} \bar{t}_k^\mu \bar{\psi}_i^k \text{ev}_i^* T_\mu \prod_{i=m+1}^{m+l} \sum_{\mu} s^\mu \text{ev}_i^* T_\mu \end{aligned}$$

be the generating function of genus g ancestor invariants. The ancestor potential is defined to be the formal expression

$$\mathcal{A}_X(\bar{t}, s) := \exp \sum_{g=0}^{\infty} \hbar^{g-1} \bar{F}_g^X(\bar{t}, s).$$

Note that \mathcal{A} depends on s (variables on the Frobenius manifold), in addition to $\bar{t} = \sum \bar{t}_k^\mu T_\mu z^k$ (variables on the ‘‘Fock space’’).

Let j be one of the first m marked points such that $\bar{\psi}_j$ is defined. Let D_j be the (virtual) divisor on $\bar{M}_{g,m+l}(X, \beta)$ defined by the image of the gluing morphism

$$\sum_{\beta'+\beta''=\beta} \sum_{l'+l''=l} \bar{M}_{0,\{j\}+l'+\bullet}(X, \beta') \times_X \bar{M}_{g,(m-1)+l''+\bullet}(X, \beta'') \rightarrow \bar{M}_{g,m+l}(X, \beta),$$

where \bullet represents the gluing point; $\bar{M}_{g,(m-1)+l''+\bullet}(X, \beta'')$ carries all first m marked points *except the j -th one*, which is carried by $\bar{M}_{0,\{j\}+l'+\bullet}(X, \beta')$. Ancestor and descendent invariants are related by the simple geometric equation

$$(1.2) \quad (\psi_j - \bar{\psi}_j) \cap [\bar{M}_{g,m+l}(X, \beta)]^{\text{vir}} = [D_j]^{\text{vir}}.$$

This can be easily seen from the definitions of ψ and $\bar{\psi}$. The morphism π in (1.1) contracts only rational curves during the processes of forgetful and stabilization morphisms. The (virtual) difference of ψ and $\bar{\psi}$ is exactly D_j .

1.2. The mixed invariants. We will consider more general *mixed invariants* with mixed ancestor and descendent insertions. Denote by

$$\langle \tau_{k_1, \bar{l}_1} \alpha_1, \dots, \tau_{k_n, \bar{l}_n} \alpha_n \rangle_{g,n,\beta}$$

the invariants with mixed descendent and ancestor insertion $\psi_i^{k_i} \bar{\psi}_i^{\bar{l}_i} \text{ev}_i^* \alpha_i$ at the i -th marked point and let

$$\begin{aligned} \langle \tau_{k_1, \bar{l}_1} \alpha_1, \dots, \tau_{k_n, \bar{l}_n} \alpha_n \rangle_g(s) &:= \sum_{l,\beta} \frac{q^\beta}{l!} \langle \tau_{k_1, \bar{l}_1} \alpha_1, \dots, \tau_{k_n, \bar{l}_n} \alpha_n, s^l \rangle_{g,n+l,\beta}, \\ \langle \tau_{k_1, \bar{l}_1} \alpha_1, \dots, \tau_{k_n, \bar{l}_n} \alpha_n \rangle_g(\bar{t}, s) &:= \sum_{m,l,\beta} \frac{q^\beta}{m!l!} \langle \tau_{k_1, \bar{l}_1} \alpha_1, \dots, \tau_{k_n, \bar{l}_n} \alpha_n, \bar{t}^m, s^l \rangle_{g,n+m+l,\beta}. \end{aligned}$$

to be the generating functions.

Equation (1.2) can be rephrased in terms of these generating functions.

Proposition 1.1. *In the stable range $2g + n \geq 3$,*

$$(1.3) \quad \begin{aligned} & \langle \tau_{k+1, \bar{l}} \alpha_1, \dots \rangle_g(\bar{t}, s) \\ &= \langle \tau_{k, \bar{l}+1} \alpha_1, \dots \rangle_g(\bar{t}, s) + \sum_v \langle \tau_k \alpha_1, T_v \rangle_0(s) \langle \tau_{\bar{l}} T^v, \dots \rangle_g(\bar{t}, s) \end{aligned}$$

where \dots denote the same mixed insertions.

In fact, only one special type of the mixed invariants will be needed. Let (X, E) be a smooth pair with $j : E \hookrightarrow X$ a smooth (infinity) divisor. At the i -th marked point, if $k_i \neq 0$, then we require that $\alpha_i = \varepsilon_i \in j_* H^*(E) \subset H^*(X)$. This type of invariants will be called *mixed invariants of special type* and the marked points with $k_i \neq 0$ will be called *marked points at infinity*.

For a birational map $f : X \dashrightarrow X'$ with exceptional loci $Z \subset X$, a mixed invariant is said to be of f -special type if $\alpha \cdot Z = 0$ for every insertion $\tau_{k, \bar{l}} \alpha$ with $k \neq 0$. When (X_{loc}, E) comes from the local model of (X, Z) , namely $X_{loc} := \tilde{E} = \mathbb{P}_Z(N_{Z/X} \oplus \mathcal{O})$ with E being the infinity divisor, these two notions of special type agree.

Proposition 1.1 will later be used (c.f. Theorem 3.7) in the following setting. Suppose that under a flop $f : X \dashrightarrow X'$ we have invariance of ancestor generating functions. To extend the invariance to allow also descendants we may reduce the problem to the $g = 0$ case and with at most one descendent insertion $\tau_k \alpha$. For local models, it is important that the invariants are of special type to ensure the invariance.

2. REVIEW OF GIVENTAL'S QUANTIZATION FORMALISM

2.1. Formal ingredients in the geometric Gromov–Witten theory. For a projective smooth variety X , Gromov–Witten theory of X consists of the following ingredients

- (i) $H := H^*(X, \mathbb{Q})$ is a \mathbb{Q} -vector space, assumed of rank N . Let $\{T_\mu\}_{\mu=1}^N$ be a basis of H and $\{s^\mu\}_{\mu=1}^N$ be the dual coordinates with $\partial/\partial s^\mu = T_\mu$. $\mathbf{1} \in H^0(X)$, the (dual of) fundamental class, is a special element. H carries a symmetric bilinear form, Poincaré pairing,

$$(\cdot, \cdot) : H \otimes H \rightarrow \mathbb{Q}.$$

Define

$$g_{\mu\nu} := (T_\mu, T_\nu)$$

and $g^{\mu\nu}$ to be the inverse matrix.

- (ii) Let $\mathcal{H}_t := \bigoplus_{k=0}^\infty H$ be the infinite dimensional complex vector space with basis $\{T_\mu \psi^k\}$. \mathcal{H}_t has a natural \mathbb{Q} -algebra structure:

$$T_\mu \psi^{k_1} \otimes T_\nu \psi^{k_2} \mapsto (T_\mu \cup T_\nu) \psi^{k_1+k_2}.$$

Let $\{t_k^\mu\}$, $\mu = 1, \dots, N$, $k = 0, \dots, \infty$, be the dual coordinates of the basis $\{T_\mu \psi^k\}$. We note that at each marked point, the insertion is

\mathcal{H}_t -valued. Let

$$t := \sum_{k,\mu} t_k^\mu T_\mu \psi^k$$

denote a general element in the vector space \mathcal{H}_t .

- (iii) The generating function of descendants $F_g^X(t)$ is a formal function on \mathcal{H}_t . The generating function of ancestors $\bar{F}_g^X(\bar{t}, s)$ is a formal function of $(\bar{t}, s) \in \mathcal{H}_t \times H$.
- (iv) H carries a (big quantum cohomology) ring structure. Let $s^\mu = t_0^\mu$ and $F_0(s) = F_0(t)|_{t_k=0, \forall k>0}$. The ring structure is defined by

$$T_{\mu_1} *_s T_{\mu_2} := \sum_{\nu, \nu'} \frac{\partial^3 F_0(s)}{\partial s^{\mu_1} \partial s^{\mu_2} \partial s^\nu} g^{\nu \nu'} T_{\nu'}.$$

$\mathbf{1}$ is the identity element of the ring.

- (v) The Dubrovin connection ∇_z on the tangent bundle TH is defined by

$$\nabla_z := d - z^{-1} \sum_{\mu} ds^\mu (T_\mu *).$$

The quantum cohomology differential equation

$$(2.1) \quad \nabla_z S = 0$$

has a fundamental solution $J = (J_{\mu, \nu}(s, z^{-1}))$, an $N \times N$ matrix-valued function, in (formal) power series of z^{-1} satisfying the conditions

$$(2.2) \quad J(s, z^{-1}) = Id + O(z^{-1}) \text{ and } J^*(s, -z^{-1})J(s, z^{-1}) = Id,$$

where $*$ denotes the adjoint with respect to (\cdot, \cdot) .

- (vi) The non-equivariant genus zero Gromov–Witten theory is graded, i.e. with a *conformal* structure. The grading is determined by an Euler field $E \in \Gamma(T_X)$,

$$(2.3) \quad E = \sum_{\mu} (1 - \frac{1}{2} \deg T_\mu) s^\mu \frac{\partial}{\partial s^\mu} + c_1(T_X).$$

2.2. Semisimple Frobenius manifolds. The concept of Frobenius manifolds was originally introduced by B. Dubrovin. We assume that the readers are familiar with the definitions of the Frobenius manifolds. See [13] Part I for an introduction. The quantum product $*$, together with Poincaré pairing, and the special element $\mathbf{1}$, defines on H a Frobenius manifold structure $(QH, *)$.

A point $s \in H$ is called a *semisimple* point if the quantum product at the tangent algebra $(T_s H, *_s)$ at $s \in H$ is isomorphic to $\bigoplus_1^N \mathbb{C}$ as an algebra. $(QH, *)$ is called *semisimple* if the semisimple points are dense in H . If $(QH, *)$ is semisimple, it has idempotents $\{\epsilon_i\}_1^N$

$$\epsilon_i * \epsilon_j = \delta_{ij} \epsilon_i.$$

defined up to S_N permutations. The *canonical coordinates* $\{u^i\}_1^N$ are defined by $\partial/\partial u^i = \epsilon_i$. When the Euler field is present, the canonical coordinates are also uniquely defined up to permutations. We will often use the *normalized form*

$$\tilde{\epsilon}_i = \frac{1}{\sqrt{(\epsilon_i, \epsilon_i)}} \epsilon_i.$$

Lemma 2.1. $\{\epsilon_i\}$ and $\{\tilde{\epsilon}_i\}$ form orthogonal bases.

Proof.

$$\begin{aligned} (\epsilon_i, \epsilon_j) &= (\epsilon_i * \epsilon_i, \epsilon_j) = (\epsilon_i, \epsilon_i * \epsilon_j) \\ &= (\epsilon_i, \delta_{ij} \epsilon_i) = \delta_{ij} (\epsilon_i, \epsilon_i). \end{aligned}$$

□

When the quantum cohomology is semisimple, the quantum differential equation (2.1) has a fundamental solution of the following type

$$\mathbf{R}(s, z) := \Psi(s)^{-1} R(s, z) e^{\mathbf{u}/z},$$

where $(\Psi_{\mu i}) := (T_\mu, \tilde{\epsilon}_i)$ is the transition matrix from $\{\tilde{\epsilon}_i\}$ to $\{T_\mu\}$; \mathbf{u} is the diagonal matrix $(\mathbf{u}_{ij}) = \delta_{ij} u^i$. The main information of \mathbf{R} is carried by $R(s, z)$, which is a (formal) power series in z . One notable difference between $J(s, z^{-1})$ and $R(s, z)$ is that the former is a (formal) power series in z^{-1} while the latter is a (formal) power series in z . See [8] and Theorem 1 in Chapter 1 of [13].

2.3. Preliminaries on quantization. Let $\mathcal{H}_q := H[z]$. Let $\{T_\mu z^k\}_{k=0}^\infty$ be a basis of \mathcal{H}_q , and $\{q_k^\mu\}$ the dual coordinates. We define an isomorphism of \mathcal{H}_q to \mathcal{H}_t as an affine vector space via a *dilaton shift* “ $t = q + z$ ”:

$$(2.4) \quad t_k^\mu = q_k^\mu + \delta^{\mu 1} \delta_{k1}.$$

The cotangent bundle $\mathcal{H} := T^* \mathcal{H}_q$ has a natural symplectic structure

$$\Omega = \sum_{k, \mu, \nu} g_{\mu\nu} dp_k^\mu \wedge dq_k^\nu$$

where $\{p_k^\mu\}$ are the dual coordinates in the fiber direction of \mathcal{H} in the natural basis $\{T_\mu z^{-k-1}\}_{k=0}^\infty$. \mathcal{H} is naturally isomorphic to the H -valued Laurent series in z^{-1} , $H[[z^{-1}]]$. In this way,

$$\Omega(f, g) = \text{Res}_{z=0}(f(-z), g(z)).$$

To quantize an infinitesimal symplectic transformation on (\mathcal{H}, Ω) , or its corresponding quadratic hamiltonians, we recall the standard Weyl quantization. An identification $\mathcal{H} = T^* \mathcal{H}_q$ of the symplectic vector space \mathcal{H} (the *phase space*) as a cotangent bundle of \mathcal{H}_q (the *configuration space*) is called a *polarization*. The “Fock space” will be a certain class of functions $f(\hbar, q)$ on \mathcal{H}_q (containing at least polynomial functions), with additional formal

variable \hbar (“Planck’s constant”). The classical observables are certain functions of p, q . The quantization process is to find for the classical mechanical system on (\mathcal{H}, Ω) a “quantum” system on the Fock space such that the classical observables, like the hamiltonians $h(q, p)$ on \mathcal{H} , are quantized to become operators $\widehat{h}(q, \partial/\partial q)$ on the Fock space.

Let $A(z)$ be an $\text{End}(H)$ -valued Laurent formal series in z satisfying

$$\Omega(Af, g) + \Omega(f, Ag) = 0,$$

for all $f, g \in \mathcal{H}$. That is, $A(z)$ defines an infinitesimal symplectic transformation. $A(z)$ corresponds to a quadratic polynomial ¹ $P(A)$ in p, q

$$P(A)(f) := \frac{1}{2}\Omega(Af, f).$$

Choose a *Darboux coordinate system* $\{q_k^i, p_k^i\}$ so that $\Omega = \sum dp_k^i \wedge dq_k^i$. The quantization $P \mapsto \widehat{P}$ assigns

$$(2.5) \quad \begin{aligned} \widehat{1} &= 1, \quad \widehat{p}_k^i = \sqrt{\hbar} \frac{\partial}{\partial q_k^i}, \quad \widehat{q}_k^i = q_k^i / \sqrt{\hbar}, \\ \widehat{p}_k^i p_l^j &= \widehat{p}_k^i \widehat{p}_l^j = \hbar \frac{\partial}{\partial q_k^i} \frac{\partial}{\partial q_l^j}, \\ \widehat{p}_k^i \widehat{q}_l^j &= q_l^j \frac{\partial}{\partial q_k^i}, \\ \widehat{q}_k^i \widehat{q}_l^j &= q_k^i q_l^j / \hbar, \end{aligned}$$

In summary, the quantization is the process

$$\begin{array}{ccccc} A & \mapsto & P(A) & \mapsto & \widehat{P(A)} \\ \text{inf. sympl. transf.} & \mapsto & \text{quadr. hamilt.} & \mapsto & \text{operator on Fock sp.} \end{array}$$

It can be readily checked that the first map is a Lie algebra isomorphism: The Lie bracket on the left is defined by $[A_1, A_2] = A_1 A_2 - A_2 A_1$ and the Lie bracket in the middle is defined by Poisson bracket

$$\{P_1(p, q), P_2(p, q)\} = \sum_{k,i} \frac{\partial P_1}{\partial p_k^i} \frac{\partial P_2}{\partial q_k^i} - \frac{\partial P_2}{\partial p_k^i} \frac{\partial P_1}{\partial q_k^i}.$$

The second map is close to be a Lie algebra homomorphism. Indeed

$$[\widehat{P}_1, \widehat{P}_2] = \{\widehat{P}_1, \widehat{P}_2\} + \mathcal{C}(P_1, P_2),$$

where the cocycle \mathcal{C} , in orthonormal coordinates, vanishes except

$$\mathcal{C}(p_k^i p_l^j, q_k^i q_l^j) = -\mathcal{C}(q_k^i q_l^j, p_k^i p_l^j) = 1 + \delta^{ij} \delta_{kl}.$$

¹Due to the nature of the infinite dimensional vector spaces involved, the “polynomials” here might have infinite many terms, but the degrees remain finite.

Example 2.2. Let $\dim H = 1$ and $A(z)$ be multiplication by z^{-1} . It is easy to see that $A(z)$ is infinitesimally symplectic.

$$(2.6) \quad \begin{aligned} P(z^{-1}) &= -\frac{q_0^2}{2} - \sum_{m=0}^{\infty} q_{m+1} p_m \\ \widehat{P(z^{-1})} &= -\frac{q_0^2}{2} - \sum_{m=0}^{\infty} q_{m+1} \frac{\partial}{\partial q_m}. \end{aligned}$$

Note that one often has to quantize the symplectic instead of the infinitesimal symplectic transformations. Following the common practice in physics, define

$$(2.7) \quad \widehat{e^{A(z)}} := e^{\widehat{A(z)}},$$

for $A(z)$ an infinitesimal symplectic transformation.

2.4. Ancestor potentials via quantization. Let N be the rank of $H = H^*(X)$ and $\mathcal{D}_N(\mathbf{t}) = \prod_{i=1}^N \mathcal{D}_{pt}(t^i)$ be the descendent potential of N points, where

$$\mathcal{D}_{pt}(t^i) \equiv \mathcal{A}_{pt}(t^i) := \exp \sum_{g=0}^{\infty} \hbar^{g-1} F_g^{pt}(t^i)$$

is the total descendent potential on a point and $t^i = \sum_k t_k^i z^k$.

Suppose that $(QH, *)$ is semisimple, then the ancestor potential can be reconstructed from the $\mathcal{D}_N(\mathbf{t})$ via the the quantization formalism.

First of all, $\{\tilde{\epsilon}_i\}$ define an *orthonormal basis* for H with canonical coordinates $\{u^i\}_{i=1}^N$. Therefore, the dual coordinates (p_k^i, q_k^i) of the basis $\{\tilde{\epsilon}_i z^k\}_{k \in \mathbb{Z}}$ for \mathcal{H} form a Darboux coordinate system. The coordinate system $\mathbf{t} = \{t_k^i\}$ is related to $\mathbf{q} = \{q_k^i\}$ by the dilaton shift (2.4). Note that $\partial/\partial q_k^i = \partial/\partial t_k^i$.

The following beautiful formula was first formulated by Givental [8]. Many special cases have since been solved by Givental and others [2], [12]. It was completely established by C. Teleman in a recent preprint [17].

Theorem 2.3 ([8, 17]).

$$(2.8) \quad \mathcal{A}_X(\bar{t}, s) = e^{\bar{c}(s)} \widehat{\Psi}^{-1}(s) \widehat{R}_X(s, z) e^{\widehat{\mathbf{u}/z}(s)} \mathcal{D}_N(\mathbf{t}),$$

where $\bar{c}(s) = \frac{1}{48} \ln \det(\epsilon_i, \epsilon_j)$.

Note that it is not very difficult to check that $\ln R_X(s, z)$ defines an infinitesimal symplectic transformation. See e.g. [8, 13]. $\widehat{R}_X(s, z)$ is therefore well-defined. By Example 2.2, $e^{\widehat{\mathbf{u}/z}}$ is also well-defined. Since the quantization involves only the z variable, $\widehat{\Psi}^{-1}(s)$ really is the induced transformation from canonical coordinates to flat coordinates. No quantization is needed.

Remark 2.4. The operator $e^{\widehat{\mathbf{u}/z}}$ can be removed from the above expression. It is shown in [8] that the string equation implies that $e^{\widehat{\mathbf{u}/z}} \mathcal{D}_N = \mathcal{D}_N$.

3. ANALYTIC CONTINUATION AND LOCAL MODELS

In the first part of this section, we discuss the issues of the analyticity of the Frobenius manifolds and the analytic continuation involved in the study of the flops $f : X \dashrightarrow X'$. We then move to the study of the local models. There the semisimplicity of the Frobenius manifolds and the quantization formalism are used to reduce the invariance of Gromov–Witten theory to the semi-classical (genus zero) case.

3.1. Review of the genus zero theory. Let $f : X \dashrightarrow X'$ be a simple \mathbb{P}^r flop with \mathcal{F} being the graph correspondence. This subsection rephrases the analytic continuation of big quantum rings proved in [14] in more algebraic terms.

Let NE_f be the cone of curve classes $\beta \in NE(X)$ with $\mathcal{F}\beta \in NE(X')$, i.e. the classes which are effective on both sides. Let

$$\mathbf{f}(q) = \frac{q}{1 - (-1)^{r+1}q}$$

be the rational function coming from the generating function of three points Gromov–Witten invariants attached to the extremal ray $\ell \subset Z \cong \mathbb{P}^r$ with positive degrees. Namely for any $i, j, k \in \mathbb{N}$ with $i + j + k = 2r + 1$,

$$\mathbf{f}(q^\ell) = \sum_{d \geq 1} \langle h^i, h^j, h^k \rangle_{0,3,d,\ell} q^{d\ell},$$

where h denotes a class in X which restricts to the hyperplane class of Z .

Define the ring

$$(3.1) \quad \mathcal{R} = \widehat{\mathbb{C}[NE_f]}[\mathbf{f}],$$

which can be regarded as certain algebraization of the Novikov ring $\widehat{NE}(X)$ in the q^ℓ variable. Notice that \mathcal{R} is canonically identified with its counterpart $\mathcal{R}' = \widehat{\mathbb{C}[NE'_f]}[\mathbf{f}']$ under \mathcal{F} since $\mathcal{F}NE_f = NE'_f$ and

$$(3.2) \quad \mathcal{F}\mathbf{f}(q^\ell) = (-1)^r - \mathbf{f}(q^\ell)$$

(via $\mathbf{f}(q) + \mathbf{f}(q^{-1}) = (-1)^r$).

Theorem 3.1. *The genus zero n -point functions with $n \geq 3$ lie in \mathcal{R} :*

$$\langle \alpha \rangle^X \in \mathcal{R}$$

for all $\alpha \in H^*(X)^{\oplus n}$. Moreover $\mathcal{F}\langle \alpha \rangle^X = \langle \mathcal{F}\alpha \rangle^{X'}$ in \mathcal{R}' .

Proof. This is the main result of [14] except the statement that $\langle \alpha \rangle^X \in \mathcal{R}$. For this, the degeneration analysis in § 4 of [14] implies

$$\langle \alpha \rangle^{\bullet X} = \sum_{\mu} m(\mu) \sum_I \langle \alpha_1 \mid \varepsilon_I, \mu \rangle^{\bullet(Y,E)} \langle \alpha_2 \mid \varepsilon^I, \mu \rangle^{\bullet(\tilde{E},E)}.$$

(A generalization to all genera is presented in the next section.) Here $Y = \text{Bl}_Z X = \bar{\Gamma}_f \subset X \times X'$; $\langle \cdot \rangle^{\bullet}$ denote invariants with possibly disconnected

domain curves. Under the projections $\phi : Y \rightarrow X$ and $\phi' : Y \rightarrow X'$, the variable q^{β_1} for $\beta_1 \in NE(Y)$ is identified with $q^{\phi_*\beta_1} \in NE(X)$. If q^{β_1} appears in the sum of contact type μ , then $(E.\beta_1) = |\mu| \geq 0$. Also

$$\mathcal{F}\phi_*\beta_1 = \phi'_*\beta_1 + |\mu|\ell' \in NE(X').$$

Hence $\langle \alpha_1 \mid \varepsilon_I, \mu \rangle^{\bullet(Y,E)} \in \widehat{\mathbb{C}[NE_f]}$.

For the local model $\tilde{E} := \mathbb{P}_Z(N_{Z/X} \oplus \mathcal{O})$, the process in [14] (§5, Theorem 5.6) via reconstruction theorem shows that there are indeed only two generators of the functional equations. One is (3.2), which is the source of analytic continuation. Another one is the quasi-linearity ([14], Lemma 5.4), which is an identity in $\mathbb{C}[NE_f]$ where no analytic continuation is needed.

Denote

$$(3.3) \quad \delta = \delta_h = q^\ell \frac{d}{dq^\ell}.$$

Then all other analytic continuation come from $\delta^m \mathbf{f}$'s with $m \geq 0$. It remains to show that $\delta^m \mathbf{f}$ is a polynomial in \mathbf{f} . This follows easily from $\delta \mathbf{f} = \mathbf{f} + (-1)^{r+1} \mathbf{f}^2$ and $\delta(\mathbf{f}_1 \mathbf{f}_2) = (\delta \mathbf{f}_1) \mathbf{f}_2 + \mathbf{f}_1 \delta \mathbf{f}_2$ by induction on m . \square

3.2. Integral structure on local models. For $X = \tilde{E}$, the above proof shows that

$$(3.4) \quad \langle \alpha \rangle \in \mathbb{C}[NE_f][\mathbf{f}] =: \mathcal{R}_{loc}$$

without the need of taking completion, where $NE_f = \mathbb{Z}_+ \gamma + \mathbb{Z}_+(\gamma + \ell)$. In fact for a given set of insertions α and genus g , the virtual dimension count shows that the *contact weight* $d_2 := (E.\beta)$ is fixed among all $\beta = d_1 \ell + d_2 \gamma$ in the series $\langle \tau_{k,l} \alpha \rangle_g^X$. Hence for $g = 0$ we must have

$$\langle \alpha \rangle^X = q^{d_2 \gamma} (p_0(\mathbf{f}) + q^\ell p_1(\mathbf{f}) + \cdots + q^{d_2 \ell} p_{d_2}(\mathbf{f}))$$

for certain polynomials $p_i(\mathbf{f}) \in \mathbb{Z}[\mathbf{f}]$.

In particular $\langle \alpha \rangle$ is an analytic function over the extended Kähler moduli $\omega \in H_{\mathbb{R}}^{1,1}(X) + i(\mathcal{K}_X \cup \mathcal{F}^{-1} \mathcal{K}_{X'})$ via the identification

$$(3.5) \quad q^\beta = e^{2\pi i(\omega.\beta)}.$$

Thus analytic continuation can be taken in the traditional complex analytic sense or as isomorphisms in the ring $\mathcal{R}_{loc} \cong \mathcal{R}'_{loc}$.

3.3. Analytic structure on the Frobenius manifolds. The Frobenius manifold corresponding to X is a priori a formal scheme, given by the formal completing \widehat{H}_X of $H^*(X, \mathbb{C})$ at the origin. The big quantum product takes values in the *Novikov ring*, or equivalently the *formal Kähler moduli* $\widehat{\mathbb{C}[NE(X)]}$. The divisor axiom implies that one may combine $H^2(X, \mathbb{C})$ directions of the Frobenius manifold and the formal Kähler moduli into a

formal completion at $q = 0$ of the complex torus

$$(3.6) \quad q \in \frac{H^2(X, \mathbb{C})}{H^2(X, \mathbb{Z})}.$$

Indeed, let $s = s' + s_1$ be a point in the Frobenius manifold with $s_1 \in H^2(X, \mathbb{C})$. The divisor axiom says that

$$\langle \alpha \rangle_\beta(s) q^\beta = \langle \alpha \rangle_\beta(s') q^\beta e^{(s_1, \beta)}.$$

Compared with (3.5), this suggests an identification of the formal Kähler moduli and the corresponding underlying Frobenius manifold in the $H^2(X)$ direction into the complex torus (3.6).² Note that the “origin” of $s_1 = 0$ is moved to the origin of $q = 0$ under this identification. In the case ℓ is the only primitive numerical class of curves and h an ample class such that $(h, \ell) = 1$, one may set $s_1 = th$. Thus, we have the familiar identification

$$q = q^\ell = e^t,$$

which will be used in the appendix. This identification can be done at the analytic level in some directions of $H^2(X)$ when the convergence is known.

Let $f : X \dashrightarrow X'$ be a simple \mathbb{P}^r flop and h be an ample divisor class dual to the extremal ray ℓ , i.e. $(h, \ell) = 1$. Then $H^2(X, \mathbb{Z}) = \mathbb{Z}h \oplus H^2(X, \mathbb{Z})^{\perp \ell}$. Theorem 3.1 gives an analytic structure on \widehat{H}_X in the h -direction:

Corollary 3.2. (i) *The Frobenius manifold structure on \widehat{H}_X can be extended to*

$$H_X := \widehat{H}_X^{\perp \ell} \times (\mathbb{P}_{q^\ell}^1 \setminus (-1)^{r+1}).$$

(ii) $H_X \cong H_{X'}$.

(iii) *If X is the local model, H_X is an analytic manifold.*

Proof. Indeed, Theorem 3.1 says that, as functions of \mathbf{f} , all invariants are defined on $\mathbf{f} \in \mathbb{C}$. Equivalently, as function of q^ℓ , all invariants are defined on $\mathbb{P}^1 \setminus (-1)^{r+1}$. This proves (i). (ii) follows from (i), and (iii) from Section 3.2 \square

Corollary 3.2 and results of the previous subsections show that the Frobenius manifold structures on the quantum cohomology of X and X' are isomorphic. The former is a series expansion of analytic functions at $q^\ell = 0$, and the latter at $q^\ell = \infty$. Considered as a one-parameter family

$$H_X \rightarrow \mathbb{P}_{q^\ell}^1 \setminus (-1)^{r+1},$$

it produces a family of product structure on $\widehat{H}_X^{\perp \ell} \otimes \widehat{\mathbb{C}}[\widehat{NE}_f]$. At two special points 0 and ∞ , the Frobenius structure specializes to the big quantum cohomology modulo extremal rays of X and of X' respectively. The term

²In string theory, the identification of weights $q^\beta = e^{2\pi i(\omega, \beta)}$ is essential in matching the A model and B model moduli spaces in mirror symmetry (cf. [4]). It is generally believed that the GW theory converges in the “large radius limit”, i.e. when $\text{Im } \omega$ is large.

“analytic continuation” used in this paper can thus be understood in this way.

3.4. Semisimplicity of big quantum ring for local models. Toric varieties admits a nice big torus action and its equivariant cohomology ring is always semisimple, hence as a deformation the equivariant big quantum cohomology ring (the Frobenius manifold) is also semisimple. Givental’s quantization formalism works in the equivariant setting, hence one way to prove the higher genus invariance for local models is to extend results in [14] to the equivariant setting. This can in principle be done, but here we take a direct approach which requires no more work.

Lemma 3.3. *For $X = \mathbb{P}_{\mathbb{P}^r}(\mathcal{O}(-1)^{r+1} \oplus \mathcal{O})$, $QH^*(X)$ is semisimple.*

Proof. By [4], the proof of Proposition 11.2.17 and [14], Lemma 5.2, the small quantum cohomology ring is given by Batyrev’s ring (though X is only semi-Fano). Namely for $q_1 = q^\ell$ and $q_2 = q^\gamma$,

$$QH_{small}^*(X) \cong \mathbb{C}[h, \zeta][q_1, q_2] / (h^{r+1} - q_1(\zeta - h)^{r+1}, (\zeta - h)^{r+1}\zeta - q_2).$$

Solving the relations, we get the eigenvalues of the quantum multiplications h^* and ζ^* :

$$(3.7) \quad h = \eta^j \omega^i q_1^{\frac{1}{r+1}} q_2^{\frac{1}{r+2}} (1 + \omega^i q_1^{\frac{1}{r+1}})^{-\frac{1}{r+2}}, \quad \zeta = \eta^j q_2^{\frac{1}{r+2}} (1 + \omega^i q_1^{\frac{1}{r+1}})^{\frac{r+1}{r+2}}$$

for $i = 0, 1, \dots, r$ and $j = 0, 1, \dots, r+1$. where ω and η are the $(r+1)$ -th and the $(r+2)$ -th root of unity respectively. As these eigenvalues of h^* (resp. ζ^*) are all different, we see that h^* and ζ^* are semisimple operators, hence $QH_{small}^*(X)$ is semisimple.

This proves that the formal Frobenius manifold $(QH^*, *)$ is semisimple at the origin $s = 0$. Since semisimplicity is an open condition, the formal Frobenius manifold $QH^*(X)$ is also semisimple. \square

Remark 3.4. The Batyrev ring for any toric variety, whether or not equal to the small quantum ring, is always semisimple.

3.5. Invariance of mixed invariants of special type.

Proposition 3.5. *For the local models, the correspondence \mathcal{F} for a simple flop induces, after the analytic continuation, an isomorphism of the ancestor potentials.*

Proof. Since a flop induces K -equivalence, by (2.3) the Euler vector fields of X and X' are identified under \mathcal{F} . By Theorem 3.1 and Lemma 3.3, X and X' give rise to isomorphic semisimple conformal formal Frobenius manifolds over \mathcal{R} (or rather \mathcal{R}_{loc}):

$$QH^*(X) \cong QH^*(X')$$

under \mathcal{F} . The first statement then follows from Theorem 2.3, the quantization formula, since all the quantities involved are uniquely determined by the underlying abstract Frobenius structure.

To be more explicit, to compare $\mathcal{F}\mathcal{A}_X$ with $\mathcal{A}_{X'}$ is equivalent to compare $\mathcal{F}(\widehat{\Psi}_X^{-1}\widehat{R}_X)e^{\widehat{\mathbf{u}}/z}$ with $\widehat{\Psi}_{X'}^{-1}\widehat{R}_{X'}e^{\widehat{\mathbf{u}}/z}$, and $\mathcal{F}\bar{c}$ with \bar{c}' . Recall that

$$\epsilon_i := \partial_{u^i}, \quad \tilde{\epsilon}_i := \frac{\epsilon_i}{\sqrt{(\epsilon_i, \epsilon_i)}}.$$

Lemma 3.6. \mathcal{F} sends canonical coordinates on X to canonical coordinates on X' : $\mathcal{F}\epsilon_i = \epsilon'_i$, $\mathcal{F}\tilde{\epsilon}_i = \tilde{\epsilon}'_i$. Moreover, \bar{c} , Ψ and \mathbf{u} transform covariantly under \mathcal{F} .

Proof. As \mathcal{F} preserves the big quantum product, \mathcal{F} sends idempotents $\{\epsilon_i\}$ to idempotents $\{\epsilon'_i\}$. Since the canonical coordinates are uniquely defined for conformal Frobenius manifolds (up to S_N permutation which is fixed by \mathcal{F}), \mathcal{F} takes canonical coordinates on X to those on X' . Furthermore, \mathcal{F} preserves the Poincaré pairing [14], hence that $\mathcal{F}\tilde{\epsilon}_i = \tilde{\epsilon}'_i$.

The \mathcal{F} covariance of $\bar{c}(s) = \frac{1}{48} \ln \det(\epsilon_i, \epsilon_j)$, the matrix $\mathbf{u}_{ij} = (\delta_{ij}u^i)$ and the matrix $\Psi_{\mu i} = (T_{\mu}, \tilde{\epsilon}_i)$ also follow immediately. For example,

$$\mathcal{F}\Psi_{\mu i} = (\mathcal{F}T_{\mu}, \mathcal{F}\tilde{\epsilon}_i).$$

□

The lemma implies that the Darboux coordinate systems on X and X' defined by canonical coordinates are compatible under \mathcal{F} . By the definition of the quantization process (2.5), which assigns differential operators $\partial/\partial q_k^i$'s in an universal manner under a Darboux coordinate system, it clearly commutes with \mathcal{F} . It is thus enough to prove the invariance of the semi-classical counterparts, or equivalently the ‘‘covariance’’ of the corresponding matrix functions, under \mathcal{F} . Note that all the invariance and covariance are up to an analytic continuation.

Therefore, one is left with the proof of the covariance of the R matrix under \mathcal{F} , after analytic continuation. Namely $\mathcal{F}R(s) = R'(\mathcal{F}s)$.

This follows from the uniqueness of R for semisimple formal conformal Frobenius manifolds. To be explicit, recall that in the proof of [13], Theorem 1, the formal series $R(s, z) = \sum_{n=0}^{\infty} R_n(s)z^n$ of the R matrix is recursively constructed by $R_0 = \text{Id}$ and the following relation in canonical coordinates:

$$(3.8) \quad (R_n)_{ij}(du^i - du^j) = [(\Psi d\Psi^{-1} + d)R_{n-1}]_{ij}.$$

Applying \mathcal{F} to it, we get $\mathcal{F}R_n = R'_n$ by induction on n . □

In order to generalize Proposition 3.5 to simple flops of general smooth varieties, which will be carried out in the next section by degeneration analysis, we have to allow descendent insertions at the infinity marked points, i.e. those marked points where the cohomology insertions come from $j_*H^*(E) \subset H^*(X)$.

Theorem 3.7. *For the local models, the correspondence \mathcal{F} for a simple flop induces, after the analytic continuation, an isomorphism of the generating functions of mixed invariants of special type in the stable range.*

Proof. Using Proposition 3.5 and 1.1 and by induction on the power k of descendent, the theorem is reduced to the case of $g = 0$ and with exactly one descendent insertion. It is of the form $\langle \tau_k \alpha, T_\nu \rangle_0(s)$ with $k \geq 0$ and by our assumption $\alpha \in j_* H^*(E)$. (Notice that for $s = 0$ this is not in the stable range.) This series is a formal sum of subseries

$$\langle \tau_k \alpha, T_\nu, T_{\mu_1}, \dots, T_{\mu_l} \rangle_{0,2+l}$$

which are sums over $\beta \in NE(\tilde{E})$. Each such series supports a unique $d_2 \geq 0$ in $\beta = d_1 \ell + d_2 \gamma$. If $d_2 = 0$ then the series and its counterpart in $X' = \tilde{E}'$ (which supports the same d_2) are both trivial since α is supported in E . If $d_2 > 0$, then the invariance follows from [14], Theorem 5.6. \square

We will generalize the theorem into the form of Theorem 0.2 by removing the local model condition after we discuss the degeneration formula.

Remark 3.8. By section 3.4 and the proof of Proposition 3.5, the canonical coordinates u_i 's, idempotents ϵ_i 's, hence the transition matrix Ψ and the R matrix can all be solved in some integral extension $\tilde{\mathcal{H}}_{loc}$ of \mathcal{H}_{loc} . It is interesting to know whether all genus g ancestor n -point generating functions take value in $\tilde{\mathcal{H}}_{loc}$ and $\mathcal{F} \langle \tau_{\bar{l}} \alpha \rangle_g^X = \langle \tau_{\bar{l}} \mathcal{F} \alpha \rangle_g^{X'}$ in $\tilde{\mathcal{H}}_{loc}$. This is plausible from Theorem 2.3 since the quantization process requires no further extensions. In fact the calculation in genus one in the Appendix suggests that $\langle \tau_{\bar{l}} \alpha \rangle_g^X$ might belong to \mathcal{H}_{loc} .

4. DEGENERATION ANALYSIS

Let $f : X \dashrightarrow X'$ be a simple \mathbb{P}^r flop with \mathcal{F} being the graph correspondence. To prove Theorem 0.2, we need to show that, up to analytic continuation,

$$\mathcal{F} \langle \tau_{k,\bar{l}} \alpha \rangle_g^X = \langle \tau_{k,\bar{l}} \mathcal{F} \alpha \rangle_g^{X'}$$

for all $\tau_{k,\bar{l}} \alpha = (\tau_{k_1,\bar{l}_1} \alpha_1, \dots, \tau_{k_n,\bar{l}_n} \alpha_n)$ being of f -special type and $g \geq 0$ with $2g + n \geq 3$.

We follow the same strategy employed in Section 4 of [14]. The two minor changes are

- (1) to generalize primary invariants to ancestors (and descendants);
- (2) to generalize genus zero invariants to arbitrary genus.

Since it is quite straightforward to make the necessary modifications, we will simply comment on the necessary changes and ask the interested readers to consult Section 4 of [14] for further details.

The first step is to apply degeneration to the normal cone

$$W = \text{Bl}_{Z \times \{0\}} X \times \mathbb{A}^1 \rightarrow \mathbb{A}^1.$$

$W_0 = Y_1 \cup Y_2$, $Y_1 = Y = \text{Bl}_Z X \xrightarrow{\phi} X$ and $Y_2 = \tilde{E} = \mathbb{P}_Z(N_{Z/X} \oplus \mathcal{O}) \xrightarrow{p} Z$. $E = Y \cap \tilde{E}$ is the ϕ exceptional divisor as well as the infinity divisor of \tilde{E} .

Define the generating series for genus g (connected) relative invariants

$$(4.1) \quad \langle A \mid \varepsilon, \mu \rangle_g^{(\tilde{E}, E)} := \sum_{\beta_2 \in NE(\tilde{E})} \frac{1}{|\text{Aut } \mu|} \langle A \mid \varepsilon, \mu \rangle_{g, \beta_2}^{(\tilde{E}, E)} q^{\beta_2}$$

and the similar one with possibly disconnected domain curves

$$(4.2) \quad \langle A \mid \varepsilon, \mu \rangle^{\bullet(\tilde{E}, E)} := \sum_{\Gamma; \mu_\Gamma = \mu} \frac{1}{|\text{Aut } \Gamma|} \langle A \mid \varepsilon, \mu \rangle_\Gamma^{\bullet(\tilde{E}, E)} q^{\beta^\Gamma} \hbar^{g^\Gamma - |\Gamma|}.$$

Here for connected invariants of genus g we assign the \hbar -weight \hbar^{g-1} , while for disconnected ones we simply assign the product weights.

Since the degeneration formula is really about the degeneration of the virtual cycles, the ancestors and descendants obey the same formula. Therefore, Proposition 4.6 of [14] can be generalized into the following form:

Proposition 4.1 (Reduction to relative local models). *To prove*

$$\mathcal{F} \langle \tau_{k, \bar{l}} \alpha \rangle_g^X = \langle \tau_{k, \bar{l}} \mathcal{F} \alpha \rangle_g^{X'}$$

for all α and k, l , it suffices to show

$$\mathcal{F} \langle \tau_{k, \bar{l}} A \mid \varepsilon, \mu \rangle_h^{(\tilde{E}, E)} = \langle \tau_{k, \bar{l}} \mathcal{F} A \mid \varepsilon, \mu \rangle_h^{(\tilde{E}', E)}$$

for all $A \in H^*(\tilde{E})^n$, $k, l \in \mathbb{Z}_+^n$, $\varepsilon \in H^*(E)^\rho$, contact type μ , and all $h \leq g$.

Proof. For the n -point mixed generating function

$$\langle \tau_{k, \bar{l}} \alpha \rangle^X = \sum_g \langle \tau_{k, \bar{l}} \alpha \rangle_g^X \hbar^{g-1} = \sum_{g; \beta \in NE(X)} \langle \tau_{k, \bar{l}} \alpha \rangle_{g, \beta}^X q^\beta \hbar^{g-1},$$

the degeneration formula gives (let $m(\mu) = \prod \mu_i$, $C_\eta = m(\mu) / |\text{Aut } \eta|$):

$$\begin{aligned} & \langle \tau_{k, \bar{l}} \alpha \rangle^X \\ &= \sum_{\beta \in NE(X)} \sum_{\eta \in \Omega_\beta} \sum_I C_\eta \langle \tau_{k_1, \bar{l}_1} \alpha_1 \mid \varepsilon_I, \mu \rangle_{\Gamma_1}^{\bullet(Y_1, E)} \langle \tau_{k_2, \bar{l}_2} \alpha_2 \mid \varepsilon^I, \mu \rangle_{\Gamma_2}^{\bullet(Y_2, E)} q^{\phi^* \beta} \hbar^{g-1} \\ &= \sum_\mu \sum_I \sum_{\eta \in \Omega_\mu} C_\eta \left(\langle \tau_{k_1, \bar{l}_1} \alpha_1 \mid \varepsilon_I, \mu \rangle_{\Gamma_1}^{\bullet(Y_1, E)} q^{\beta_1} \hbar^{g^{\Gamma_1} - |\Gamma_1|} \right) \\ & \quad \times \left(\langle \tau_{k_2, \bar{l}_2} \alpha_2 \mid \varepsilon^I, \mu \rangle_{\Gamma_2}^{\bullet(Y_2, E)} q^{\beta_2} \hbar^{g^{\Gamma_2} - |\Gamma_2|} \right) \hbar^\rho, \end{aligned}$$

where we have used $g-1 = \sum_i (g^{\Gamma_i} - |\Gamma_i|) + \rho$ with ρ being the number of contact points. Notice that $\beta = \phi_* \beta_1 + p_* \beta_2$ and we identify $q^{\beta_1} = q^{\phi_* \beta_1}$, $q^{\beta_2} = q^{p_* \beta_2}$ throughout our degeneration analysis.

We consider also absolute invariants $\langle \tau_{k, \bar{l}} \alpha \rangle^{\bullet X}$ with product weights in \hbar . Then by comparing the order of automorphisms,

$$(4.3) \quad \langle \tau_{k, \bar{l}} \alpha \rangle^{\bullet X} = \sum_\mu m(\mu) \sum_I \langle \tau_{k_1, \bar{l}_1} \alpha_1 \mid \varepsilon_I, \mu \rangle_{\Gamma_1}^{\bullet(Y_1, E)} \langle \tau_{k_2, \bar{l}_2} \alpha_2 \mid \varepsilon^I, \mu \rangle_{\Gamma_2}^{\bullet(Y_2, E)} \hbar^\rho.$$

To compare $\mathcal{F} \langle \tau_{k, \bar{l}} \alpha \rangle^{\bullet X}$ and $\langle \tau_{k, \bar{l}} \mathcal{F} \alpha \rangle^{\bullet X'}$, by [14], Proposition 4.4, we may assume that $\alpha_1 = \alpha'_1$ and $\alpha'_2 = \mathcal{F} \alpha_2$. This choice of cohomology liftings

identifies the relative invariants of (Y_1, E) and those of (Y'_1, E') with the same topological types. It remains to compare

$$\langle \tau_{k_2, \bar{l}_2} \alpha_2 \mid \varepsilon^I, \mu \rangle^{\bullet(\tilde{E}, E)} \quad \text{and} \quad \langle \tau_{k_2, \bar{l}_2} \alpha_2 \mid \varepsilon^I, \mu \rangle^{\bullet(\tilde{E}', E)}.$$

We further split the sum into connected invariants. Let Γ^π be a connected part with the contact order μ^π induced from μ . Denote $P : \mu = \sum_{\pi \in P} \mu^\pi$ a partition of μ and $P(\mu)$ the set of all such partitions. Then

$$\langle A \mid s, \mu \rangle^{\bullet(\tilde{E}, E)} = \sum_{P \in P(\mu)} \prod_{\pi \in P} \sum_{\Gamma^\pi} \frac{1}{|\text{Aut } \mu^\pi|} \langle A^\pi \mid s^\pi, \mu^\pi \rangle_{\Gamma^\pi}^{\bullet(\tilde{E}, E)} q^{\beta^{\Gamma^\pi}} \hbar^{g^{\Gamma^\pi} - 1}.$$

In the summation over Γ^π , the only index to be summed over is β^{Γ^π} on \tilde{E} and the genus. This reduces the problem to $\langle A^\pi \mid s^\pi, \mu^\pi \rangle_g^{\bullet(\tilde{E}, E)}$.

Instead of working with all genera, the proposition follows from the same argument by reduction modulo \hbar^{g+1} . \square

Proposition 4.2 (Relative to absolute). *For a simple flop $\tilde{E} \dashrightarrow \tilde{E}'$, to prove*

$$\mathcal{F} \langle \tau_{\bar{l}} A \mid \varepsilon, \mu \rangle_g^{\bullet(\tilde{E}, E)} = \langle \tau_{\bar{l}} \mathcal{F} A \mid \varepsilon, \mu \rangle_g^{\bullet(\tilde{E}', E)}$$

for all A, l, ε, μ , it suffices to show for mixed invariants of special type

$$\mathcal{F} \langle \tau_{\bar{l}} A, \tau_k \varepsilon \rangle_h^{\tilde{E}} = \langle \tau_{\bar{l}} \mathcal{F} A, \tau_k \varepsilon \rangle_h^{\tilde{E}'}$$

for all A, l, ε and $k \in \mathbb{Z}_+^\rho$, and all $h \leq g$.

Sketch of Proof. We apply degeneration to the normal cone for $Z \hookrightarrow \tilde{E}$ to get $W \rightarrow \mathbb{A}^1$. Then $W_0 = Y_1 \cup Y_2$ with $\pi : Y_1 \cong \mathbb{P}_E(\mathcal{O}_E(-1, -1) \oplus \mathcal{O}) \rightarrow E$ a \mathbb{P}^1 bundle and $Y_2 \cong \tilde{E}$.

By induction on g and then on $(|\mu|, n, \rho)$ with ρ in the reverse ordering, the same procedure used in the proof of [14], Proposition 4.8 leads to

$$\begin{aligned} \langle \tau_{\bar{l}} A, \tau_{\mu_1-1} \varepsilon_{i_1}, \dots, \tau_{\mu_\rho-1} \varepsilon_{i_\rho} \rangle_g^{\bullet \tilde{E}} &= \sum_{\mu'} m(\mu') \times \\ &\sum_{l'} \langle \tau_{\mu_1-1} \varepsilon_{i_1}, \dots, \tau_{\mu_\rho-1} \varepsilon_{i_\rho} \mid \varepsilon^{l'}, \mu' \rangle_0^{\bullet(Y_1, E)} \langle \tau_{\bar{l}} A \mid \varepsilon_{l'}, \mu' \rangle_g^{\bullet(\tilde{E}, E)} + R, \end{aligned}$$

where R denotes the remaining terms which either have lower genus or have total contact order smaller than $d_2 = |\mu| = |\mu'|$ or have number of insertions fewer than n on the (\tilde{E}, E) side or the invariants on (\tilde{E}, E) are disconnected ones.

For the main terms, the integrals on (Y_1, E) are all fiber integrals and this allows to conclude that there is a single top order term in the sum given by

$$C(\mu) \langle \tau_{\bar{l}} A \mid \varepsilon_I, \mu \rangle^{\bullet(\tilde{E}, E)}$$

with $C(\mu) \neq 0$. Thus the proposition follows by induction. \square

Proof of Main Theorems. We only need to prove Theorem 0.2.

By Proposition 4.1, the theorem is reduced to the relative local case. Moreover, for any insertion $\tau_{k,\bar{l}}\alpha$ with nontrivial descendent ($k \geq 1$), we may select the cohomology lifting of α to be $(\alpha, 0)$. To avoid trivial invariants this insertion must go to the (Y_1, E) side in the degeneration formula. Hence the theorem is reduced to the case of relative local models $X = \tilde{E} = \mathbb{P}_{\mathbb{P}^r}(\mathcal{O}(-1)^{r+1} \oplus \mathcal{O})$ with at most ancestor insertions.

Now by Propositions 4.2, the proof is further reduced to the case for absolute invariants of the form

$$\langle \tau_{\bar{l}} A, \tau_k \varepsilon \rangle_g^{\tilde{E}}$$

which are mixed invariants of special type. But this is exactly the content of Theorem 3.7. The proof is complete. \square

Remark 4.3. The proof also shows that nontrivial descendent invariants of f -special type without $2g + n \geq 3$, that is $(g, n) = (0, 1)$ or $(0, 2)$, are also invariant under \mathcal{F} .

5. EXPLICIT FORMULAE FOR PRIMARY INVARIANTS ATTACHED TO THE EXTREMAL RAY

In this section we specialize our curve classes to the extremal ray and investigate the invariance in more explicit terms. Note that all Gromov–Witten invariants discussed here are primary.

5.1. Generalities concerning flopping curves. Let $\dim X \geq 3$. In general, for ℓ being a curve class with $(K_X \cdot \ell) = 0$, the virtual dimension of $\overline{M}_{g,n}(X, d\ell)$ is given by

$$(5.1) \quad D_{g,n} = (\dim X - 3)(1 - g) + n$$

which is independent of d . If moreover ℓ is a log-extremal ray of flopping type (e.g. in our case ℓ is the line class of Z), $\langle \alpha \rangle_{g,n,d\ell}$ depends only on the local geometry of $(Z, N_{Z/X})$ for $d \geq 1$. But for $d = 0$ it depends on the global geometry of X .

If $D_{g,n}$ is negative, all Gromov–Witten invariants must vanish. From fundamental class axiom, all primary invariants $\langle 1, \dots \rangle_{g,n,d\ell}$ must vanish if $\overline{M}_{g,n-1}(X, d\ell)$ exists. We are therefore left with 3 cases: $g = 0$, $g = 1$, $\dim X = 3$ (and $g \geq 2$).

$g = 0$ then $D_{g,3} = \dim X$ and the 3-point functions with $d \geq 1$ are expected to correct the classical cubic product corresponding to $d = 0$, which is indeed the case for simple flops. There is no n -point invariant with $n \geq 4$ and $d = 0$. In fact for simple flops the n -point functions with $n \geq 4$, $d \geq 1$ are invariant under \mathcal{F} .

If $g = 1$ then $D_{1,n} = n$. By the fundamental class axiom each cohomology insertion must be a divisor. Hence if $d \geq 1$ by the divisor axiom the n -point

invariants are determined by the “partition function”

$$\int_{[\overline{M}_{1,0}(X,d\ell)]^{\text{vir}}} 1.$$

For $d = 0$ and $n \geq 2$, the divisor axiom shows that $\langle \alpha \rangle_{1,n,0} = 0$. $n = 1$ case requires different consideration.

Indeed it is well known that $\overline{M}_{g,n}(X,0) \cong X \times \overline{M}_{g,n}$ and the virtual fundamental class is given by

$$(5.2) \quad [\overline{M}_{g,n}(X,0)]^{\text{vir}} = e(\mathcal{E}) \cap [X \times \overline{M}_{g,n}]$$

where the obstruction bundle is given by $\mathcal{E} = \pi_1^* T_X \otimes \pi_2^* \mathcal{H}_g^\vee$ with \mathcal{H}_g being the Hodge bundle. Let $\lambda_i = c_i(\mathcal{H}_g)$.

For $(g,n) = (1,1)$ we clearly have

$$c(\mathcal{E}) = c_{\text{top}}(X) - c_{\text{top}-1}(X) \cdot \lambda_1.$$

Thus for one point invariants we get a (semi-)classical term

$$(5.3) \quad \langle \alpha \rangle_{1,0}^X = -(c_{\text{top}-1}(X) \cdot \alpha)_X \cdot \int_{\overline{M}_{1,1}} \lambda_1 = -\frac{1}{24} (c_{\text{top}-1}(X) \cdot \alpha)_X,$$

where the basic Hodge integral $\int_{\overline{M}_{1,1}} \lambda_1 = 1/24$ is used.

For simple flops, we will verify that $\mathcal{F} \langle \alpha \rangle_1^X = \langle \mathcal{F} \alpha \rangle_1^{X'}$ in the next subsection by showing that the genus one invariants with $d \geq 1$ correct the semi-classical defect $\langle \alpha \rangle_{1,0}^X - \langle \mathcal{F} \alpha \rangle_{1,0}^{X'}$.

For $\dim X = 3$ (and $g \geq 2$), $D_{g,n} = n$. As in the $g = 1$ case we are reduced to consider the case $n = 0$. For simple \mathbb{P}^1 flop, the extremal invariants with $d \geq 1$ are determined by Faber and Pandharipande [5] to be

$$\langle - \rangle_{g,d} := \int_{[\overline{M}_{g,0}(X,d\ell)]^{\text{vir}}} 1 = C_g d^{2g-3}$$

where $C_g = |\chi(M_g)| / (2g-3)!$. We claim that the generating function

$$(5.4) \quad \langle - \rangle_g := \sum_{d=0}^{\infty} \langle - \rangle_{g,d} q^d = \langle - \rangle_{g,0} + C_g \delta^{2g-3} \mathbf{f},$$

is invariant under \mathcal{F} (since $2g-3 \geq 1$), where the operator δ is defined in (3.3). The second term is invariant following the analysis in Section 3. For degree zero term, it is not difficult to see $\langle - \rangle_{g,0}^X = \langle - \rangle_{g,0}^{X'}$: The degeneration analysis in Section 4 reduces the proof to a corresponding statement for local models. The local models of X and X' are both isomorphic to $\mathbb{P}_{\mathbb{P}^1}(\mathcal{O}(-1)^2 \oplus \mathcal{O})$, and hence have the same degree zero invariants. In fact it is not hard to see from (5.2) that

$$(5.5) \quad e(\mathcal{E}) = \frac{(-1)^g}{2} (c_3(X) - c_2(X) \cdot c_1(X)) \lambda_{g-1}^3.$$

The invariance of these Chern numbers can be directly verified for \mathbb{P}^1 flops.

5.2. The genus one case. Let G be the genus one potential of twisted GW invariants on $\mathcal{O}_{\mathbb{P}^r}(-1)^{r+1}$ without marked points. Namely, for ℓ being the line class in \mathbb{P}^r ,

$$G(q) := \langle - \rangle_1 = \sum_{d \geq 0} \langle - \rangle_{1,d\ell} q^d.$$

For $r = 1, d \geq 1$, the formula

$$\langle - \rangle_{1,d} = \frac{1}{12d}$$

was first obtained by physical consideration in [1] and later mathematically justified in [9]. Here, by using the theory of semisimple Frobenius manifolds, we generalize it to all $r \in \mathbb{N}$ and get

Proposition 5.1. For $d \in \mathbb{N}$,

$$(5.6) \quad \langle - \rangle_{1,d} = (-1)^{d(r+1)} \frac{r+1}{24d}.$$

In fact the $g = 1$ case can be achieved without using the machinery of quantization. Givental has shown in [7], Theorem 4.1 that the total differential dG can be expressed, up to a constant, in terms of the canonical coordinates u_i 's as

$$(5.7) \quad dG = \sum_i \left[d \log \Delta_i / 48 - c_{-1}^i du_i / 24 + R_{ii}^1 du_i / 2 \right],$$

where the RHS is determined by the *equivariant genus zero theory*.

In the appendix we will determine all the terms $u_i, \Delta_i := 1/(\epsilon_i \cdot \epsilon_i)$ and R_{ii}^1 step by step. The final result (Theorem 6.12) is equivalent to (5.6):

$$(5.8) \quad \delta_h G = (-1)^{r+1} \frac{r+1}{24} \mathbf{f}$$

in the *non-equivariant limit*, where $\delta_h := qd/dq$.

We use it in the following setting: Let $f : X \dashrightarrow X'$ be a simple \mathbb{P}^r flop. Under the canonical correspondence $\mathcal{F} = [\bar{\Gamma}_f]$ we have $\mathcal{F}\ell = -\ell'$ and we identify $q' = q^{-1}$. Then $\delta_{h'} = -\delta_h$ and by (3.2), $\delta_h \mathbf{f}(q) = \delta_{h'} \mathbf{f}(q')$. Hence $\delta_h^2 G(q) = \delta_{h'}^2 G'(q')$ and by the divisor axiom

$$\langle h, \dots, h \rangle_{1,n}^X = \delta_h^n G(q) = (-1)^{n-2} \delta_{h'}^n G'(q') = (-1)^n \langle h', \dots, h' \rangle_{1,n}^{X'}$$

Since $\mathcal{F}h^k = (-1)^k h'^k$, this implies the invariance of n -point functions for all $n \geq 2$. (The invariance for $g \geq 2, n \geq 1$ is proved in the same way.)

For $n = 1$, to prove the invariance we may assume that X and X' are local models. We compute via (5.3), (5.8) and (3.2) that

$$\langle h \rangle_1^X - \langle \mathcal{F}h \rangle_1^{X'} = -\frac{1}{24} ((c_{2r}(X) \cdot h) - (c_{2r}(X') \cdot (\xi' - h'))) - \frac{r+1}{24}.$$

Since $X \cong X'$, the invariance will follow from

Proposition 5.2.

$$(c_{2r}(X) \cdot (2h - \xi)) = -(r+1).$$

Proof. Since $X = \mathbb{P}_Z(N_{Z/X} \oplus \mathcal{O}) \xrightarrow{p} Z$, by the Leray-Hirsh theorem,

$$H^*(X) = \mathbb{Z}[h, \xi] / (h^{r+1}, \xi(\xi - h)^{r+1}).$$

From $0 \rightarrow \mathcal{O} \rightarrow \mathcal{O}(1) \otimes p^*(N_{Z/X} \oplus \mathcal{O}) \rightarrow T_{X/Z} \rightarrow 0$ we get $c(X) = p^*c(Z).c(T_{X/Z}) = (1+h)^{r+1}(1+\xi)(1+\xi-h)^{r+1}$. So we need to calculate the degree $2r+1$ terms in

$$c(X).(2h - \xi) = (1+h)^{r+1}(1+\xi)(1+\xi-h)^{r+1}(2h - \xi)$$

under the additional relation $h^r \xi^{r+1} = 1$.

Since $\xi(\xi - h)^{r+1} = 0$, $(1+\xi)(1+(\xi-h))^{r+1}$ start with

$$(\xi - h)^{r+1} + C_1^{r+1} \xi(\xi - h)^r + C_1^{r+1}(\xi - h)^r + C_2^{r+1} \xi(\xi - h)^{r-1} + \text{lower terms.}$$

Thus in $(1+h)^{r+1} = h^{r+1} + C_1^{r+1}h^r + \dots$ only h^r and h^{r-1} contribute to terms with total degree $2r+1$. There are four such terms:

$$C_1^{r+1}C_1^{r+1}h^r(\xi - h)^r(2h - \xi) = -(C_1^{r+1})^2,$$

$$C_1^{r+1}C_2^{r+1}h^r\xi(\xi - h)^{r-1}(2h - \xi) = -C_1^{r+1}C_2^{r+1},$$

$$C_2^{r+1}h^{r-1}(\xi - h)^{r+1}(2h - \xi) = 2C_2^{r+1}$$

and (using the Chern relation $0 = \xi(\xi - h)^{r+1} = \xi^{r+1} - C_1^{r+1}\xi^r h + \dots$)

$$\begin{aligned} C_2^{r+1}C_1^{r+1}h^{r-1}\xi(\xi - h)^r(2h - \xi) &= C_2^{r+1}C_1^{r+1}(2h^r\xi(\xi - h)^r - h^{r-1}\xi^2(\xi - h)^r) \\ &= C_2^{r+1}C_1^{r+1}(2 - C_1^{r+1} + C_1) = C_2^{r+1}C_1^{r+1}. \end{aligned}$$

The sum is $-(r+1)^2 + (r+1)r = -(r+1)$ as expected. \square

6. APPENDIX: THE CALCULATIONS FOR $g = 1$

We calculate the genus one twisted Gromov–Witten invariants on the bundle $\mathcal{O}_{\mathbb{P}^r}(-1)^{r+1}$ by using Givental's work [7].

6.1. The Frobenius structure and the canonical coordinates. This calculation follows the general scheme outlined in [7].

Let E be the total space of $\mathcal{O}_{\mathbb{P}^r}(-1)^{r+1}$ over \mathbb{P}^r and consider the torus action of $\mathbb{C}^* \times \mathbb{C}^*$ on E such that the first \mathbb{C}^* acts trivially on \mathbb{P}^r and by scalar multiplication on the fiber of E and the second one acts by

$$\alpha \cdot [x_0 : x_1 : \dots : x_r] = [\alpha^{l_0}x_0 : \alpha^{l_1}x_1 : \dots : \alpha^{l_r}x_r].$$

Let λ and λ' denote the characters of these two actions respectively. Let $i : E_{\text{fixed}} \hookrightarrow E$ be the injection, where E_{fixed} denotes the fixed loci. Denote by p the equivariant hyperplane class of \mathbb{P}^r for the first action.

Proposition 6.1. *The characteristic polynomial of p^* in equivariant small quantum cohomology is given by*

$$(6.1) \quad q(\lambda - p)^{r+1} = p^{r+1}.$$

Proof. Formally this follows from the formula for small quantum ring of local models in Lemma 3.3, with h, ξ being replaced by p, λ .

Alternatively, by [7] Corollary 4.4 (in the limit $\lambda' \rightarrow 0$ which exists),

$$\text{Quantum } (\lambda - p)^{r+1} = \text{Classical } \frac{1}{1 + (-1)^r q} (\lambda - p)^{r+1}.$$

Using the fact that $p^{r+1} = 0$ in the classical product and the quantum p^k coincides with the classical one for $k \leq r$, we get

$$(1 + (-1)^r q)(\lambda - p)^{r+1} = (\lambda - p)^{r+1} - (-1)^{r+1} p^{r+1}$$

in small quantum product. The proposition follows. \square

Solving this formally in q or locally analytically in the Kähler moduli coordinate t with $q = e^t$, we get (with $\xi := e^{2\pi i/(r+1)}$):

$$(6.2) \quad p_i = \frac{\lambda}{1 + (-1)^r \xi^i q^{-\frac{1}{r+1}}}, \quad i = 0, 1, \dots, r.$$

Using the *standard basis* $1, p, \dots, p^r$ of $H_{\mathbb{C}^\times}^*(\mathbb{P}^r)$, it is easy to see that p_i 's are the eigenvalues of the quantum multiplication operator p^* . It follows that for $k \leq r$, p_i^k 's are the eigenvalues of $p^{k*} = (p^*)^{\circ k}$. The common eigenvectors ϵ_i 's which simultaneously diagonalize the quantum product $p^k * \epsilon_i = p_i^k \epsilon_i$ are known as the *canonical basis*. Let t_i 's be the *standard (flat) coordinates* dual to $\{1, p, \dots, p^r\}$ and u_i 's be the *canonical coordinates* dual to $\{\epsilon_0, \dots, \epsilon_r\}$. The canonical basis $\{\epsilon_i\}$ is *orthogonal* with respect to the Poincaré pairing since quantum multiplications are self adjoint.

In practice we may simply define u_i and then ϵ_i by the relation

$$(6.3) \quad du_i = \sum_{k=0}^r p_i^k dt_k.$$

Calculations in canonical coordinates are thus essentially *formal linear algebra* which in our case are reduced to the *roots-coefficients* relation

$$(6.4) \quad (-1)^r p^{r+1} - \mathbf{f} C_1^{r+1} \lambda p^r + \mathbf{f} C_2^{r+1} \lambda^2 p^{r-1} - \dots + (-1)^{r+1} \mathbf{f} \lambda^{r+1} = 0.$$

Remark 6.2. It is important to point out that all the coefficients involve \mathbf{f} only! Thus *in principle* everything *canonically determined* by the Frobenius structure should be invariant under flops after one more differentiation by $\delta_h = qd/dq \equiv d/dt$. This note provides a demonstration in this direction.

We start with determining $\sum_i c_{-1}^i du_i/24$. Recall that c_{-1}^i is the localization of $c_{\dim-1}/c_{\dim}$ at the i -th fixed point $[0 : \dots : 1 : \dots : 0]$. Since the Chern roots at there are given by

$$\underbrace{l_i \lambda' + \lambda, \dots, l_i \lambda' + \lambda}_{r+1}, \quad \underbrace{(l_j - l_i) \lambda'}_r \quad (j \neq i),$$

we see that

$$c_{-1}^i = \frac{r+1}{l_i \lambda' + \lambda} + \sum_{j \neq i} \frac{1}{(l_j - l_i) \lambda'}.$$

In the limit $\lambda' \rightarrow 0$, the trouble terms with $1/\lambda'$ must be canceled out with the corresponding terms in R_{ii}^1 's via (5.7). So $c_{-1}^i = (r+1)/\lambda$ and

$$(6.5) \quad \sum_i c_{-1}^i du_i / 24 = \frac{r+1}{24\lambda} \sum_i du_i = \frac{r+1}{24\lambda} \sum_{k=0}^r \left(\sum_i p_i^k \right) dt_k.$$

Since we are only interested in the non-equivariant limit $\lambda \rightarrow 0$, by (6.4) the only terms remaining are with $k = 1$, hence we obtain

Proposition 6.3.

$$\sum_i c_{-1}^i du_i / 24 = (-1)^r \frac{(r+1)^2}{24} \mathbf{f} dt_1.$$

In Givental's formulation, we will need to identify $t = t_1$ in the sequel. But we will keep $q = e^t$ independent of t_1 whenever possible.

6.2. The Poincaré pairing and the formula for $\Delta_i := \langle \epsilon_i, \epsilon_i \rangle^{-1}$.

Lemma 6.4. *The equivariant Poincaré pairing is given by*

$$\langle p^k \cdot p^l \rangle = C_{r-d}^{2r-d} \frac{1}{\lambda^{2r+1-d}}$$

where $d = k + l$. It vanishes if $d > r$.

Proof. The localization formula for the first \mathbf{C}^\times action reads as:

$$\int_E \omega = \int_{\mathbb{P}^r} \frac{i^* \omega}{(\lambda - p)^{r+1}} = \int_{\mathbb{P}^r} \frac{i^* \omega}{\lambda^{r+1}} \left(1 - \frac{p}{\lambda}\right)^{-(r+1)}.$$

Since p^k vanishes if $k > r$, we get by Taylor expansion that

$$\left(1 - \frac{p}{\lambda}\right)^{-(r+1)} = 1 + C_1^{r+1} \frac{p}{\lambda} + C_2^{r+2} \left(\frac{p}{\lambda}\right)^2 + \dots + C_r^{r+r} \left(\frac{p}{\lambda}\right)^r.$$

So the Poincaré pairing is given by

$$\langle p^k \cdot p^l \rangle = \int_E p^d = C_{r-d}^{r+r-d} \frac{1}{\lambda^{r+1+(r-d)}} = C_{r-d}^{2r-d} \frac{1}{\lambda^{2r+1-d}}.$$

□

Define $a_i := 1 + (-1)^r \zeta^i q^{-\frac{1}{r+1}}$ so $p_i = \lambda/a_i$. For convenience we also denote by $c_i = (-1)^r \zeta^i q^{-\frac{1}{r+1}}$, so $a_i = 1 + c_i$ and $c_j = c_0 \zeta^j = c_i \zeta^{j-i}$.

Proposition 6.5. *The canonical basis ϵ_i 's are given by*

$$\epsilon_i = \frac{q c_i}{r+1} a_i^r \prod_{l \neq i} \left(1 - a_l \frac{p}{\lambda}\right).$$

They are vector fields along the Kähler moduli variable q .

Proof. Call the RHS ε_i and it suffices to show that $du_j(\varepsilon_i) = \delta_{ji}$. By (6.3), the effect of du_j is simply the replacement of p by $p_j = \lambda/a_j$. Hence

$$\begin{aligned} du_j(\varepsilon_i) &= \frac{qc_i}{r+1} \left(\frac{a_i}{a_j}\right)^r \prod_{l \neq i} (a_j - a_l) \\ &= \frac{qc_i}{r+1} \left(\frac{a_i}{a_j}\right)^r \prod_{l \neq i} (-1)^r (\zeta^j - \zeta^l) q^{-\frac{1}{r+1}} \\ &= \frac{\zeta^{jr+i}}{r+1} \left(\frac{a_i}{a_j}\right)^r \prod_{l \neq i} (1 - \zeta^{l-j}) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases}. \end{aligned}$$

□

Denote by $S_k^l(x)$ the k -th elementary symmetric polynomial in x_j 's with $0 \leq j \leq r$ and $j \neq l$. We often need some basic formulae on roots of unity whose verifications are elementary and omitted.

Lemma 6.6. *The following formulae for ζ hold:*

- (1) $S_k^i(\zeta) := S_k^i(\zeta^0, \dots, \zeta^r) = (-1)^k \zeta^{ki}$ for $k = 0, 1, \dots, r$.
- (2) $\sum_{i=0}^r \zeta^{ki} = 0$ for $k = 1, 2, \dots, r$.

Clearly the lemma applies to c_j 's as well. Then

$$\begin{aligned} \varepsilon_i &= \frac{qc_i a_i^r}{r+1} \prod_{j \neq i} \left[\left(1 - \frac{p}{\lambda}\right) - c_j \frac{p}{\lambda} \right] \\ (6.6) \quad &= \frac{qc_i a_i^r}{r+1} \sum_{k=0}^r (-1)^k S_k^i(c) \left(\frac{p}{\lambda}\right)^k \left(1 - \frac{p}{\lambda}\right)^{r-k} \\ &= \frac{qc_i a_i^r}{r+1} \sum_{k=0}^r c_i^k \left(\frac{p}{\lambda}\right)^k \left(1 - \frac{p}{\lambda}\right)^{r-k}. \end{aligned}$$

It is convenient to denote $\langle p \rangle^{k+l} = \langle p^k, p^l \rangle = \langle p^{k+l} \rangle$.

Lemma 6.7. *For $k = 0, 1, \dots, r-1$,*

$$\left\langle \left(\frac{p}{\lambda}\right)^k \left(1 - \frac{p}{\lambda}\right)^{2r-k} \right\rangle = 0.$$

Proof. By Lemma 6.4, it equals

$$\begin{aligned} &\left\langle \frac{p}{\lambda} \right\rangle^k - C_1^{2r-k} \left\langle \frac{p}{\lambda} \right\rangle^{k+1} + \dots + (-1)^{r-k} C_{r-k}^{2r-k} \left\langle \frac{p}{\lambda} \right\rangle^{k+(r-k)} \\ &= \frac{1}{\lambda^{2r+1}} \left(C_{r-k}^{2r-k} - C_1^{2r-k} C_{r-(k+1)}^{2r-(k+1)} + \dots + (-1)^{r-k} C_{r-k}^{2r-k} C_{r-(k+(r-k))}^{2r-(k+(r-k))} \right) \\ &= \frac{C_r^{2r-k}}{\lambda^{2r+1}} \left(1 - C_1^{r-k} + \dots + (-1)^{r-k} C_{r-k}^{r-k} \right) \\ &= \frac{C_r^{2r-k}}{\lambda^{2r+1}} (1-1)^{r-k} = 0. \end{aligned}$$

□

Using (6.6), Lemma 6.6 and Lemma 6.7, we compute

$$\begin{aligned}
\Delta_i^{-1} &= \langle \epsilon_i, \epsilon_i \rangle \\
&= \frac{q^2 c_i^2 a_i^{2r}}{(r+1)^2} (-1)^r (r+1) c_i^r \left\langle \left(\frac{p}{\lambda} \right)^r \left(1 - \frac{p}{\lambda} \right)^r \right\rangle \\
&= \frac{q^2 c_i^2 a_i^{2r}}{r+1} c_i^r \left\langle \frac{p}{\lambda} \right\rangle^r \quad (c_i^{r+1} = q^{-1}) \\
&= \frac{q c_i a_i^{2r}}{(r+1) \lambda^{2r+1}}.
\end{aligned}$$

Lemma 6.8. *As a function in q , the norm square inverse of ϵ_i equals*

$$\Delta_i = (r+1) \lambda q^{-1} c_i^{-1} p_i^{2r}.$$

Proposition 6.9.

$$d \log(\Delta_0 \Delta_1 \cdots \Delta_r) = r \frac{1 - (-1)^r q}{1 + (-1)^r q} d \log q = r(1 - 2(-1)^r \mathbf{f}) d \log q.$$

Proof. Simply take log differentiation of

$$\Delta_0 \Delta_1 \cdots \Delta_r = (r+1)^{r+1} \lambda^{(2r+1)(r+1)} \xi^{-\frac{r(r+1)}{2}} q^{-r} \mathbf{f}^{2r}.$$

□

6.3. The transition matrix Ψ and the connection one form $\Psi d\Psi^{-1}$.

The matrix Ψ is defined by

$$p^\mu = \sum_i \Psi_\mu^i \tilde{\epsilon}_i = \sum_i \langle p^\mu, \tilde{\epsilon}_i \rangle \tilde{\epsilon}_i$$

relative to the orthonormal frame $\{\tilde{\epsilon}_i := \sqrt{\Delta_i} \epsilon_i\}$. From (6.3), we get the dual relation

$$(6.7) \quad p^k = \sum_i p_i^k \epsilon_i.$$

Hence by Lemma 6.8,

$$(6.8) \quad \Psi_\mu^i = p_i^\mu / \sqrt{\Delta_i} = \sqrt{\frac{q c_i}{r+1}} \lambda^{-\frac{1}{2}} p_i^{\mu-r} = \sqrt{\frac{q c_i}{r+1}} \lambda^{\mu-r-\frac{1}{2}} a_i^{r-\mu}.$$

The inverse Ψ^{-1} has already been determined in Proposition 6.5 up to a normalization factor. Indeed, $(\Psi^{-1})_j^\mu$ is the coefficient of p^μ in the expression of $\sqrt{\Delta_j} \epsilon_j$, hence we get

$$(6.9) \quad (\Psi^{-1})_j^\mu = (-1)^\mu \sqrt{\frac{q c_j}{r+1}} \lambda^{r-\mu+\frac{1}{2}} S_\mu^j(a).$$

To proceed, notice that by Lemma 6.6

$$\begin{aligned} S_\mu^j(a) &= \sum_{k_1 < \dots < k_\mu, k_s \neq j} (1 + c_{k_1}) \cdots (1 + c_{k_\mu}) \\ &= \sum \left(1 + (c_{k_1} + \dots + c_{k_\mu}) + (c_{k_1}c_{k_2} + \dots) + \dots + (c_{k_1} \cdots c_{k_\mu}) \right) \\ &= C_\mu^r - C_{\mu-1}^{r-1}c_j + C_{\mu-2}^{r-2}c_j^2 - \dots + (-1)^\mu c_j^\mu. \end{aligned}$$

We regard $q = e^t$ and take differentiation in $t = \log q$. Then

$$(d\Psi^{-1})_j^\mu = \frac{r}{2(r+1)} (\Psi^{-1})_j^\mu dt + (-1)^\mu \sqrt{\frac{qc_j}{r+1}} \lambda^{r-\mu+\frac{1}{2}} dS_\mu^j(a),$$

and we compute

$$\begin{aligned} (\Psi d\Psi^{-1})_j^i &= \sum_{\mu=0}^r \Psi_\mu^i (d\Psi^{-1})_j^\mu = \frac{r\delta_{ij}}{2(r+1)} dt - \frac{q\sqrt{c_i c_j}}{(r+1)^2} dt \\ &\quad \times \sum_{\mu=1}^r (-1)^\mu a_i^{r-\mu} \left(-C_{\mu-1}^{r-1}c_j + 2C_{\mu-2}^{r-2}c_j^2 - \dots + (-1)^\mu \mu c_j^\mu \right). \end{aligned}$$

The last sum equals

$$\begin{aligned} &c_j \left(a_i^{r-1} - C_1^{r-1} a_i^{r-2} + \dots + (-1)^{r-2} C_{r-2}^{r-1} a_i + (-1)^{r-1} \right) \\ &\quad + 2c_j^2 \left(a_i^{r-2} - C_1^{r-2} a_i^{r-3} + \dots + (-1)^{r-3} C_{r-3}^{r-2} a_i + (-1)^{r-2} \right) + \dots \\ &= c_j (a_i - 1)^{r-1} + 2c_j^2 (a_i - 1)^{r-2} + \dots + rc_j^r \\ &= c_j c_i^{r-1} + 2c_j^2 c_i^{r-2} + \dots + rc_j^r. \end{aligned}$$

If $i = j$, using $qc_j^{r+1} = 1$ we get

$$(\Psi d\Psi^{-1})_i^i = \frac{r}{2(r+1)} dt - \frac{r(r+1)}{2} \frac{qc_j^{r+1}}{(r+1)^2} dt = 0.$$

This agrees with the well-known fact that $\Psi d\Psi^{-1}$ is skew-symmetric.

If $i \neq j$, using $c_j = c_i \zeta^{j-i}$ we get the connection matrix to be

$$(6.10) \quad (\Psi d\Psi^{-1})_j^i = \frac{\zeta^{\frac{1}{2}(j-i)}}{(r+1)^2} \sum_{\mu=1}^r \mu \zeta^{\mu(j-i)} d \log q.$$

6.4. The first asymptotic matrix R^1 and the final computation.

Now we identify the Kähler moduli coordinate $t = t_1$. Recall the defining relation of R^1 :

$$(6.11) \quad (\Psi d\Psi^{-1})_j^i = R_{ij}^1 (du_i - du_j).$$

The off-diagonal part of R^1 is uniquely solvable from (6.11). Since the LHS involves only $d \log q = dt$ and $du_i - du_j = \sum_{k=0}^r (p_i^k - p_j^k) dt_k$, we get

$$(6.12) \quad R_{ij}^1 = \frac{(-1)^r \zeta^{\frac{1}{2}(j-i)}}{(r+1)^2 \lambda (\zeta^j - \zeta^i)} q^{\frac{1}{r+1}} a_i a_j \sum_{\mu=1}^r \mu \zeta^{\mu(j-i)}.$$

Denote by

$$g_k(\zeta) := \frac{1}{\zeta^k - 1} \sum_{\mu=1}^r \mu \zeta^{\mu k} \sum_{\mu=1}^r \mu \zeta^{-\mu k}.$$

The diagonal part is determined by the *flatness relation* up to a constant:

$$(6.13) \quad dR_{ii}^1 + \sum_j R_{ij}^1 R_{ji}^1 (du_i - du_j) = 0.$$

Hence by (6.10), (6.12) and writing out $a_i a_j$ we get

$$\begin{aligned} dR_{ii}^1 &= \sum_{j \neq i} \frac{(-1)^r q^{\frac{1}{r+1}} a_i a_j}{(r+1)^4 \lambda (\zeta^j - \zeta^i)} dt \sum_{\mu=1}^r \mu \zeta^{\mu(j-i)} \sum_{\mu=1}^r \mu \zeta^{\mu(i-j)} \\ &= \frac{(-1)^r \zeta^{-i} q^{\frac{1}{r+1}}}{(r+1)^4 \lambda} dt \sum_{k=1}^r g_k(\zeta) + \frac{1}{(r+1)^4 \lambda} dt \sum_{k=1}^r (\zeta^k + 1) g_k(\zeta) \\ &\quad + \frac{(-1)^r \zeta^i q^{-\frac{1}{r+1}}}{(r+1)^4 \lambda} dt \sum_{k=1}^r \zeta^k g_k(\zeta). \end{aligned}$$

Lemma 6.10.

$$\sum_{k=1}^r (\zeta^k + 1) g_k(\zeta) = 0.$$

Proof. Let $f(\zeta)$ be the expression, then $f(\zeta) = f(\zeta^{-1}) = -f(\zeta)$. \square

Let

$$\Xi_r := \sum_{k=1}^r g_k(\zeta).$$

By the lemma,

$$dR_{ii}^1 = \frac{(-1)^r \Xi_r}{(r+1)^4 \lambda} (\zeta^{-i} q^{\frac{1}{r+1}} - \zeta^i q^{-\frac{1}{r+1}}) dt.$$

So by integration in t ,

$$R_{ii}^1 = \frac{(-1)^r \Xi_r}{(r+1)^3 \lambda} (\zeta^{-i} q^{\frac{1}{r+1}} + \zeta^i q^{-\frac{1}{r+1}}).$$

Since we are only interested in the non-equivariant limit $\lambda \rightarrow 0$, by (6.3) the only terms remaining in $\sum_{i=0}^r R_{ii}^1 du_i$ is in the dt_1 direction. Thus

$$\begin{aligned} \sum_{i=0}^r R_{ii}^1 du_i &= \frac{\Xi_r}{(r+1)^3 \lambda} \sum_{i=0}^r (c_i^{-1} + c_i) \frac{\lambda}{a_i} dt \\ &= \frac{\Xi_r}{(r+1)^3} \sum_{i=0}^r \left((c_i^{-1} + c_i) \prod_{j \neq i} a_j \right) \frac{dt}{1 + (-1)^r q^{-1}}. \end{aligned}$$

The sum can be computed as (using $a_j = 1 + c_j$ and Lemma 6.6)

$$\begin{aligned} &\sum_{i=0}^r \left((c_i^{-1} - 1) \prod_{j \neq i} a_j + \prod a_j \right) \\ &= \sum_{i=0}^r \left((c_i^{-1} - 1)(1 - c_i + c_i^2 - \cdots + (-1)^r c_i^r) + (1 + (-1)^r q^{-1}) \right) \\ &= (r+1)(-2 + 1 + (-1)^r q^{-1}), \end{aligned}$$

and we get

$$(6.14) \quad \sum_{i=0}^r R_{ii}^1 du_i = \frac{\Xi_r}{(r+1)^2} \frac{1 - (-1)^r q}{1 + (-1)^r q} dt.$$

So by (5.7), Proposition 6.3, Proposition 6.9 and (6.14),

$$dG = \left\{ \left[\frac{r}{48} + \frac{\Xi_r}{2(r+1)^2} \right] \frac{1 - (-1)^r q}{1 + (-1)^r q} - \frac{(-1)^r (r+1)^2}{24} \frac{q}{1 + (-1)^r q} \right\} dt.$$

Thus $dG = (a + b\mathbf{f}) dt$ for some a and b . In Givental's formula a should be ignored since dG has no constant terms. Nevertheless the constant Ξ_r can be explicitly computed:

Lemma 6.11.

$$\Xi_r = -\frac{(r+2)(r+1)^2 r}{24}.$$

We leave the interesting proof to the readers. With it, a simple substitution leads to the final result (a redundant constant $-r(r+1)/48$ has been removed):

Theorem 6.12.

$$dG = \left[\frac{(-1)^{r+1}(r+1)}{24} \frac{q}{1 - (-1)^{r+1}q} \right] d \log q.$$

REFERENCES

- [1] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa; *Holomorphic anomalies in topological field theories*, Nuclear Physics **B405** (1993), 279–304.
- [2] A. Bertram, I. Ciocan-Fontanine, B. Kim; *Two proofs of a conjecture of Hori and Vafa*, Duke Math. J. 126 (2005), no. 1, 101–136.
- [3] T. Coates and Y. Ruan; *Quantum Cohomology and Crepant Resolutions: A Conjecture*, arXiv:0710.5901.

- [4] D.A. Cox and S. Katz; *Mirror Symmetry and Algebraic Geometry*, Math. Surv. Mono. **68**, Amer. Math. Soc. 1999.
- [5] C. Faber and R. Pandharipande; *Hodge integrals and Gromov-Witten theory*, Invent. Math. **139** (2000), 173–199.
- [6] E. Getzler; *The jet-space of a Frobenius manifold and higher-genus Gromov-Witten invariants*, Frobenius manifolds, 45–89, Aspects Math., E36, Vieweg, Wiesbaden, 2004.
- [7] A. Givental; *Elliptic Gromov-Witten invariants and the generalized mirror conjecture*, in “Integrable Systems and Algebraic Geometry”, World Sci., 1998, 105 – 155.
- [8] —; *Gromov-Witten invariants and quantization of quadratic Hamiltonians*, Dedicated to the memory of I. G. Petrovskii on the occasion of his 100th anniversary. Mosc. Math. J. **1** (2001), no. 4, 551–568, 645.
- [9] T. Graber and R. Pandharipande; *Localizations of virtual classes*, Invent. Math. **135** (1999), 487–518.
- [10] M. Kontsevich and Yu. Manin; *Relations between the correlators of the topological sigma-model coupled to gravity*, Comm. Math. Phys. **196** (1998), no. 2, 385–398.
- [11] Y.-P. Lee; *Notes on axiomatic Gromov–Witten theory and applications*, arXiv:0710.4349v2.
- [12] —; *Witten’s conjecture and the Virasoro conjecture for genus up to two*, Gromov-Witten theory of spin curves and orbifolds, 31–42, Contemp. Math., 403, Amer. Math. Soc., Providence, RI, 2006.
Invariance of tautological equations II: Gromov–Witten theory, (Appendix with Y. Iwao) math.AG/0605708.
- [13] — and R. Pandharipande; *Frobenius manifolds, Gromov–Witten theory, and Virasoro constraints*, a book in preparation. Material needed for this paper is available from <http://www.math.princeton.edu/~rahulp/>.
- [14] Y.-P. Lee, H.-W. Lin and C.-L. Wang; *Flops, motives and invariance of quantum rings*, Ann. of Math., to appear.
- [15] J. Li; *A degeneration formula for GW-invariants*, J. Diff. Geom. **60** (2002), 199–293.
- [16] D. Maulik and R. Pandharipande; *A topological view of Gromov-Witten theory*, Topology **45** (2006), no. 5, 887–918.
- [17] C. Teleman; *The structure of 2D semi-simple field theories*, arXiv:0712.0160v1.
- [18] C.-L. Wang; *K-equivalence in birational geometry*, in “Proceeding of the Second International Congress of Chinese Mathematicians (Grand Hotel, Taipei 2001)”, International Press 2003, math.AG/0204160.

Y. IWAO AND Y.-P. LEE: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH 84112-0090, U.S.A.

E-mail address: yshr@math.utah.edu

E-mail address: yplee@math.utah.edu

H.-W. LIN AND C.-L. WANG: DEPARTMENT OF MATHEMATICS, NATIONAL CENTRAL UNIVERSITY, JHONGLI 32001, TAIWAN; NATIONAL CENTER FOR THEORETICAL SCIENCES, HSINCHU 30013, TAIWAN.

E-mail address: linhw@math.ncu.edu.tw

E-mail address: dragon@math.ncu.edu.tw; dragon@math.cts.nthu.edu.tw